

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 February 2001 (15.02.2001)

PCT

(10) International Publication Number  
**WO 01/10903 A2**

(51) International Patent Classification<sup>7</sup>: C07K 14/00

(21) International Application Number: PCT/US00/21878

(22) International Filing Date: 9 August 2000 (09.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/147,986 9 August 1999 (09.08.1999) US  
60/160,807 21 October 1999 (21.10.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:  
US 60/147,986 (CIP)  
Filed on 9 August 1999 (09.08.1999)  
US 60/160,807 (CIP)  
Filed on 21 October 1999 (21.10.1999)

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043

(US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). YANG, Junming [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

## Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/10903 A2

(54) Title: PROTEASES AND PROTEASE INHIBITORS

(57) Abstract: The invention provides human proteases and protease inhibitors (PPIM) and polynucleotides which identify and encode PPIM. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of PPIM.

## PROTEASES AND PROTEASE INHIBITORS

### TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of proteases and protease  
5 inhibitors and to the use of these sequences in the diagnosis, treatment, and prevention of cell  
proliferative and autoimmune/inflammatory disorders.

### BACKGROUND OF THE INVENTION

Proteolytic processing is an essential component of normal cell growth, differentiation,  
10 remodeling, and homeostasis. The cleavage of peptide bonds within cells is necessary for the maturation  
of precursor proteins to their active forms, the removal of signal sequences from targeted proteins, the  
degradation of incorrectly folded proteins, and the controlled turnover of peptides within the cell.  
Proteases participate in apoptosis, inflammation, and tissue remodeling during embryonic development,  
wound healing, and normal growth. They are necessary components of bacterial, parasitic, and viral  
15 invasion and replication within a host. Four principal categories of mammalian proteases have been  
identified based on active site structure, mechanism of action, and overall three-dimensional structure.  
(See Beynon, R.J. and J.S. Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University  
Press, New York NY, pp. 1-5.)

The serine proteases (SPs) are a large family of proteolytic enzymes that include the digestive  
20 enzymes, trypsin and chymotrypsin; components of the complement cascade and of the blood-clotting  
cascade; and enzymes that control the degradation and turnover of macromolecules of the extracellular  
matrix. SPs are so named because of the presence of a serine residue found in the active catalytic site for  
protein cleavage. The active site of all SPs is composed of a triad of residues including the  
aforementioned serine, an aspartate, and a histidine residue. SPs have a wide range of substrate  
25 specificities and can be subdivided into subfamilies on the basis of these specificities. The main sub-  
families are trypases which cleave after arginine or lysine; aspases which cleave after aspartate;  
chymases which cleave after phenylalanine or leucine; metases which cleavage after methionine; and  
serases which cleave after serine. Clp protease is a unique member of the serine protease family as its  
activity is controlled by a regulatory subunit that binds and hydrolyzes ATP. Clp protease was  
30 originally found in plant chloroplasts but is believed to be widespread in both prokaryotic and eukaryotic  
cells (Maurizi, M.R. et al. (1990) J. Biol. Chem. 266:12546-12552). SKD3, a mammalian homolog of  
the bacterial Clp regulatory subunit, has recently been identified in mouse (Perier, F. et al. (1995) Gene  
152:157-163).

Cysteine proteases are involved in diverse cellular processes ranging from the processing of  
35 precursor proteins to intracellular degradation. Mammalian cysteine proteases include lysosomal

cathepsins and cytosolic calcium activated proteases, calpains. Of particular note, cysteine proteases are produced by monocytes, macrophages and other cells of the immune system which migrate to sites of inflammation and in their protective role secrete various molecules to repair damaged tissue. These cells may overproduce the same molecules and cause tissue destruction in certain disorders. In autoimmune diseases such as rheumatoid arthritis, the secretion of the cysteine protease, cathepsin C, degrades collagen, laminin, elastin and other structural proteins found in the extracellular matrix of bones. The cathepsin family of lysosomal proteases includes the cysteine proteases: cathepsins B, H, K, L, O<sub>2</sub>, and S; and the aspartyl proteases; cathepsins D and G. Various members of this endosomal protease family are differentially expressed. Some, such as cathepsin D, have a ubiquitous tissue distribution while others, such as cathepsin L, are found only in monocytes, macrophages, and other cells of the immune system.

Aspartic proteases include bacterial penicillopepsin, mammalian pepsin, renin, chymosin, and certain fungal proteases. The characteristic active site residues of aspartic proteases are a pair of aspartic acid residues, for example, Asp33 and Asp213 in penicillopepsin. Aspartic proteases are also called acid proteases because the optimum pH for their activity is between 2 and 3. In this pH range, one of the aspartate residues is ionized and the other is neutral. A potent inhibitor of aspartic proteases is the hexapeptide pepstatin which, in the transition state, resembles normal substrates.

Carboxypeptidases A and B are the principal mammalian representatives of the metallo-protease family. Both are exopeptidases of similar structure and active site configuration. Carboxypeptidase A, like chymotrypsin, prefers C-terminal aromatic and aliphatic side chains of hydrophobic nature, whereas carboxypeptidase B is directed toward basic arginine and lysine residues. Active site components include zinc, which coordinates two glutamic acid and one histidine residues in the protein.

Ubiquitin proteases are associated with the ubiquitin conjugation system (UCS), a major pathway for the degradation of cellular proteins in eukaryotic cells and some bacteria. The UCS mediates the elimination of abnormal proteins and regulates the half-lives of important regulatory proteins that control cellular processes such as gene transcription and cell cycle progression. In the UCS pathway, proteins targeted for degradation are conjugated to a ubiquitin, a small heat stable protein. The ubiquitinated protein is then recognized and degraded by proteasome, a large, multisubunit proteolytic enzyme complex, and ubiquitin is released for reutilization by ubiquitin protease. The UCS is implicated in the degradation of mitotic cyclic kinases, oncoproteins, tumor suppressor genes such as p53, viral proteins, cell surface receptors associated with signal transduction, transcriptional regulators, and mutated or damaged proteins (Ciechanover, A. (1994) *Cell* 79:13-21). A murine proto-oncogene, Unp, encodes a nuclear ubiquitin protease whose overexpression leads to oncogenic transformation of NIH3T3 cells, and the human homolog of this gene is consistently elevated in small cell tumors and adenocarcinomas of the lung (Gray, D.A. (1995) *Oncogene* 10:2179-2183).

Protease inhibitors and other regulators of protease activity control the activity and effects of proteases. Protease inhibitors have been shown to control pathogenesis in animal models of proteolytic disorders (Murphy, G. (1991) Agents Actions Suppl. 35:69-76). Low levels of the cystatins, low molecular weight inhibitors of the cysteine proteases, correlate with malignant progression of tumors  
5 (Calkins, C. et al (1995) Biol. Biochem. Hoppe Seyler 376:71-80).

The plasma inter- $\alpha$ -trypsin inhibitor family molecules are serine protease inhibitors (serpins) composed of a 240 kDa plasma protein complex of at least five different types of glycoproteins. These glycoproteins consist of four heavy (H) chains and one 30 kDa light (L) chain named H1, H2, H3, H4, and L, and are independently synthesized and proteolytically processed from precursor proteins (Daveau,  
10 M. et al. (1998) Arch. Biochem. Biophys. 350:315-323; and Salier, J.P. et al. (1992) Mamm. Genome 2:233-239). The plasma inter- $\alpha$ -trypsin inhibitor light chains have sequence similarity to the Kunitz trypsin inhibitors which appear to be present in all vertebrates (Salier, J.P. (1990) Trends Biochem. Sci. 15:435-439). Some examples of the Kunitz trypsin inhibitors are tissue factor pathway inhibitor, which regulates tissue factor-induced coagulation, and protease nexin-2, which regulates serum coagulation  
15 factor XIa. (Broze, G.J. (1995) Annu. Rev. Med. 46:103-112; and Wagner, S.L. et al. (1993) Brain Res. 626:90-98). The heavy chain precursors encode a signal peptide sequence and the mature chain. Other plasma inter- $\alpha$ -trypsin inhibitor heavy chains have been described in human and rodents (Bourguignon, J. et al. (1993) Eur. J. Biochem. 212:771-776; Salier, 1992, supra; and Salier, J.P. (1996) Biochem. J. 315:1-9). The expression of the rat plasma inter- $\alpha$ -trypsin inhibitor genes is regulated by inflammation  
20 in vivo. The genes are predominantly expressed in the rat liver, but H2 and H3 mRNA is also present in brain, intestine, and stomach (Daveau, supra).

Kallistatins are members of the serine protease inhibitor family. Kallistatin forms a specific and covalently-linked complex with tissue kallikrein, which is a serine proteinase capable of cleaving kininogen to release vasoactive kinin. Components of the tissue kallikrein-kinin system include tissue  
25 kallikrein, kallistatin, kininogen, kinin, bradykininB1 and B2 receptors, and kininases (Chao, J. and L. Chao (1995) Biol. Chem. Hoppe Seyler 376:705-713).

Proteases and protease inhibitory molecules may contain amino acid sequence motifs which determine protein-protein interactions, such as the potential metal-binding site of von Willebrand factor type A3 (vWFA3) motif, glycine-amino acid-serine-amino acid-serine. This motif is also required for  
30 ligand interaction in the homologous I-type domains of integrins CR3 and LFA-1 (Huizinga, E.G. (1997) Structure 5:1147-1156).

Protease inhibitors play a major role in the regulation of the activity and effect of proteases. They have been shown to control pathogenesis in animal models of proteolytic disorders and in the treatment of HIV (Murphy, G. (1991) Agents Actions Suppl. 35:69-76; and Pakyz, A. and D. Israel  
35 (1997) J. Am. Pharm. Assoc. (Wash.) NS37:543-551).

The discovery of new proteases and protease inhibitors and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cell proliferative and autoimmune/inflammatory disorders.

5

### SUMMARY OF THE INVENTION

The invention features purified polypeptides, proteases and protease inhibitors, referred to collectively as "PPIM" and individually as "PPIM-1," "PPIM-2," "PPIM-3," "PPIM-4," "PPIM-5," "PPIM-6," "PPIM-7," "PPIM-8," "PPIM-9," "PPIM-10," "PPIM-11," "PPIM-12," "PPIM-13," "PPIM-14," "PPIM-15," "PPIM-16," "PPIM-17," "PPIM-18," "PPIM-19," "PPIM-20," "PPIM-21," 10 "PPIM-22," "PPIM-23," "PPIM-24," "PPIM-25," "PPIM-26," and "PPIM-27." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic 15 fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-27.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the 20 group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting 25 of SEQ ID NO:1-27. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-27. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:28-54.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence 30 selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1- 35 27. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In

another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions

whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

- 5           The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, c) a
- 10   polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

- The invention further provides a composition comprising an effective amount of a polypeptide
- 15   comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected
- 20   from the group consisting of SEQ ID NO:1-27, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional PPIM, comprising administering to a patient in need of such treatment the composition.

- 25           The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected
- 30   from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a
- 35   method of treating a disease or condition associated with decreased expression of functional PPIM,

comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring  
5 amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in  
10 the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional PPIM, comprising administering to a patient in need of such treatment the composition.

15 The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino acid sequence selected  
20 from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

25 The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-27, c) a biologically active fragment of an amino  
30 acid sequence selected from the group consisting of SEQ ID NO:1-27, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-27. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the  
35 test compound with the activity of the polypeptide in the absence of the test compound, wherein a



change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:28-54, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, ii) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:28-54, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

### BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding PPIM.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of PPIM.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones  
5 encoding PPIM were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

## DESCRIPTION OF THE INVENTION

10 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

15 It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings  
20 as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in  
25 connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

## DEFINITIONS

"PPIM" refers to the amino acid sequences of substantially purified PPIM obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human,  
30 and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of PPIM. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PPIM either by directly interacting with PPIM or by acting on components of the biological pathway in which PPIM participates.

35 An "allelic variant" is an alternative form of the gene encoding PPIM. Allelic variants may

result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides.

- 5 Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

- “Altered” nucleic acid sequences encoding PPIM include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as PPIM or a polypeptide with at least one functional characteristic of PPIM. Included within this definition are
- 10 polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding PPIM, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding PPIM. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent PPIM. Deliberate
- 15 amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of PPIM is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include:
- 20 asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

- The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic
- 25 molecules. Where “amino acid sequence” is recited to refer to a sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

- “Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in
- 30 the art.

- The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of PPIM. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of PPIM either by directly interacting with PPIM or by acting on components of the biological pathway in which PPIM
- 35 participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant.

Antibodies that bind PPIM polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic PPIM, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution.

Compositions comprising polynucleotide sequences encoding PPIM or fragments of PPIM may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (PE Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
20	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
25	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
30	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
35	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the

side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

10 A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of PPIM or the polynucleotide encoding PPIM which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:28-54 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:28-54, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:28-54 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:28-54 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:28-54 and the region of SEQ ID NO:28-54 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-27 is encoded by a fragment of SEQ ID NO:28-54. A fragment of SEQ ID NO:1-27 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-27. For example, a fragment of SEQ ID NO:1-27 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-27. The precise length of a fragment of SEQ ID NO:1-27 and the region of SEQ ID NO:1-27 to which the fragment corresponds

are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

35        *Matrix: BLOSUM62*

*Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

5 *Expect: 10*

*Word Size: 11*

*Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the  
10 length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

15 Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the  
20 percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

25 Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with  
30 polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

35 *Matrix: BLOSUM62*



*Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

5 *Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150  
10 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for  
15 chromosome replication, segregation and maintenance.

The term “humanized antibody” refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a  
20 complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e.,  
25 binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v)  
30 SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target  
35 sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for

nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C<sub>0</sub>t or R<sub>0</sub>t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of PPIM which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of PPIM which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of PPIM. For example, modulation may

cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of PPIM.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an PPIM may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of PPIM.

"Probe" refers to nucleic acid sequences encoding PPIM, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example

Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived  
5 from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000  
10 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program  
15 (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public  
20 from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful  
25 in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This  
30 artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant  
35 nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding PPIM, or fragments thereof, or PPIM itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels

and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

"Transformation" describes a process by which exogenous DNA is introduced into a recipient  
5 cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells  
10 includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid  
15 introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or *in vitro* fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic  
20 organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook, J. et al. (1989), *supra*.

25 A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater  
30 sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species  
35 variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides

generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

### THE INVENTION

The invention is based on the discovery of new human proteases and protease inhibitors (PPIM), the polynucleotides encoding PPIM, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative and autoimmune/inflammatory disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding PPIM. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each PPIM were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each PPIM and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding PPIM. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:28-54 and

to distinguish between SEQ ID NO:28-54 and related polynucleotide sequences. The polypeptides encoded by the selected fragments of SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, 5 SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, and SEQ ID NO:54 are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express PPIM as a fraction of total tissues expressing PPIM. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing PPIM as a fraction of total tissues expressing PPIM. Column 5 lists the vectors used to subclone each cDNA 10 library. Of particular note is the expression of SEQ ID NO:28 in gastrointestinal tissue. Of particular note is the tissue-specific expression of SEQ ID NO:51. Over 83% of the tissues expressing SEQ ID NO:51 are derived from gastrointestinal tissue, particularly the liver.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding PPIM were isolated. Column 1 references the nucleotide SEQ ID 15 NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:30 maps to chromosome 9 within the interval from 78.4 to 90.6 centiMorgans. This interval also contains a gene associated with cell proliferation.

SEQ ID NO:37 maps to chromosome 12 within the interval from 116.6 to 118.9 centiMorgans. 20 This interval also contains a gene associated with a neurological disorder.

SEQ ID NO:47 maps to chromosome 4 within the interval from 99.2 to 105.2 centiMorgans. This interval also contains a gene associated with cardiovascular disease.

The invention also encompasses PPIM variants. A preferred PPIM variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence 25 identity to the PPIM amino acid sequence, and which contains at least one functional or structural characteristic of PPIM.

The invention also encompasses polynucleotides which encode PPIM. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:28-54, which encodes PPIM. The polynucleotide sequences of SEQ 30 ID NO:28-54, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding PPIM. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least 35 about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence



encoding PPIM. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:28-54 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:28-54. Any one of the  
5 polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of PPIM.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding PPIM, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the  
10 invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring PPIM, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode PPIM and its variants are generally capable of  
15 hybridizing to the nucleotide sequence of the naturally occurring PPIM under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding PPIM or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons  
20 are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PPIM and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode PPIM and PPIM  
25 derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding PPIM or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing  
30 to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:28-54 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) Methods Enzymol. 152:399-407; Kimmel, A.R. (1987) Methods Enzymol. 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the  
35 embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA

polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (PE Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (PE Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (PE Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding PPIM may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library

does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, PE Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled.

10 Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode PPIM may be cloned in recombinant DNA molecules that direct expression of PPIM, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express PPIM.

15

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter PPIM-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

20

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of PPIM, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be

30

35 recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively,

fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding PPIM may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, PPIM itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431 A peptide synthesizer (PE Biosystems). Additionally, the amino acid sequence of PPIM, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active PPIM, the nucleotide sequences encoding PPIM or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding PPIM. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding PPIM. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding PPIM and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding PPIM and appropriate transcriptional and translational control

elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

- 5 A variety of expression vector/host systems may be utilized to contain and express sequences encoding PPIM. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco
- 10 mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; Scorer, C.A. et al. (1994) Bio/Technology 12:181-184; Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945; Takamatsu, N. (1987) EMBO
- 15 J. 6:307-311; Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659; and Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia
- 20 viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.) The invention is not limited by the host cell
- 25 employed.

- In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding PPIM. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding PPIM can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid
- 30 (Life Technologies). Ligation of sequences encoding PPIM into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of PPIM are needed, e.g. for the production of antibodies,
- 35

vectors which direct high level expression of PPIM may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of PPIM. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, supra; and Scorer, supra.)

Plant systems may also be used for expression of PPIM. Transcription of sequences encoding PPIM may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, supra; Broglie, supra; and Winter, supra.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding PPIM may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses PPIM in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of PPIM in cell lines is preferred. For example, sequences encoding PPIM can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue

culture techniques appropriate to the cell type.

- Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk<sup>-</sup>* and *apr<sup>-</sup>* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.)
- Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

- Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding PPIM is inserted within a marker gene sequence, transformed cells containing sequences encoding PPIM can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PPIM under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

- In general, host cells that contain the nucleic acid sequence encoding PPIM and that express PPIM may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

- Immunological methods for detecting and measuring the expression of PPIM using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PPIM is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. 1V; Coligan, J.E. et al.

(1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization  
5 or PCR probes for detecting sequences related to polynucleotides encoding PPIM include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding PPIM, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6  
10 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

15 Host cells transformed with nucleotide sequences encoding PPIM may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode PPIM may be designed to contain signal sequences which direct secretion  
20 of PPIM through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the  
25 protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

30 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding PPIM may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric PPIM protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of PPIM activity. Heterologous protein and peptide moieties  
35 may also facilitate purification of fusion proteins using commercially available affinity matrices. Such



moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the PPIM encoding sequence and the heterologous protein sequence, so that PPIM may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled PPIM may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

PPIM of the present invention or fragments thereof may be used to screen for compounds that specifically bind to PPIM. At least one and up to a plurality of test compounds may be screened for specific binding to PPIM. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of PPIM, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which PPIM binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express PPIM, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing PPIM or cell membrane fractions which contain PPIM are then contacted with a test compound and binding, stimulation, or inhibition of activity of either PPIM or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with PPIM, either in solution or affixed to a solid support, and detecting the binding of PPIM to the compound. Alternatively, the

assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

PPIM of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of PPIM. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for PPIM activity, wherein PPIM is combined with at least one test compound, and the activity of PPIM in the presence of a test compound is compared with the activity of PPIM in the absence of the test compound. A change in the activity of PPIM in the presence of the test compound is indicative of a compound that modulates the activity of PPIM. Alternatively, a test compound is combined with an in vitro or cell-free system comprising PPIM under conditions suitable for PPIM activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of PPIM may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding PPIM or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding PPIM may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding PPIM can also be used to create "knockin" humanized animals

(pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding PPIM is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with  
 5 potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress PPIM, e.g., by secreting PPIM in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) *Biotechnol. Annu. Rev.* 4:55-74).

### THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between  
 10 regions of PPIM and proteases and protease inhibitors. In addition, the expression of PPIM is closely associated with cell proliferation, inflammation, the immune response, and gastrointestinal, neurological, and reproductive tissue. Therefore, PPIM appears to play a role in cell proliferative and autoimmune/inflammatory disorders. In the treatment of disorders associated with increased PPIM expression or activity, it is desirable to decrease the expression or activity of PPIM. In the treatment of  
 15 disorders associated with decreased PPIM expression or activity, it is desirable to increase the expression or activity of PPIM.

Therefore, in one embodiment, PPIM or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PPIM. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic  
 20 keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle,  
 25 ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis,  
 30 cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis,  
 35 psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic

anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.

In another embodiment, a vector capable of expressing PPIM or a fragment or derivative thereof  
5 may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PPIM including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified PPIM in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PPIM including, but not limited to, those provided  
10 above.

In still another embodiment, an agonist which modulates the activity of PPIM may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PPIM including, but not limited to, those listed above.

In a further embodiment, an antagonist of PPIM may be administered to a subject to treat or  
15 prevent a disorder associated with increased expression or activity of PPIM. Examples of such disorders include, but are not limited to, those cell proliferative and autoimmune/inflammatory disorders described above. In one aspect, an antibody which specifically binds PPIM may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express PPIM.

20 In an additional embodiment, a vector expressing the complement of the polynucleotide encoding PPIM may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PPIM including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate  
25 therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

30 An antagonist of PPIM may be produced using methods which are generally known in the art. In particular, purified PPIM may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind PPIM. Antibodies to PPIM may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab  
35 expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally

preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with PPIM or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PPIM have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of PPIM amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to PPIM may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce PPIM-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for PPIM may also be generated. For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion

of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

- 5           Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between PPIM and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two  
10 non-interfering PPIM epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

- Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for PPIM. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of PPIM-antibody complex divided by the  
15 molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple PPIM epitopes, represents the average affinity, or avidity, of the antibodies for PPIM. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular PPIM epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$   
20 L/mole are preferred for use in immunoassays in which the PPIM-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of PPIM, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical  
25 Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

- The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of PPIM-antibody  
30 complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

- In another embodiment of the invention, the polynucleotides encoding PPIM, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene  
35 expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA,

PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding PPIM. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PPIM. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

- 5 In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 10 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

- In another embodiment of the invention, polynucleotides encoding PPIM may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked 20 inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial 25 hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C 30 virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in PPIM expression or regulation causes disease, the expression of PPIM from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic 35 deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in PPIM are treated by constructing mammalian expression vectors encoding PPIM and introducing these vectors by mechanical means into PPIM-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle  
 5 delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of PPIM include, but are not limited to, the pCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA),  
 10 PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). PPIM may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-  
 15 5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, supra)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene  
 20 encoding PPIM from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and  
 25 A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to PPIM expression are treated by constructing a retrovirus vector consisting of (i) the  
 30 polynucleotide encoding PPIM under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) *Proc. Natl. Acad. Sci.*  
 35 *USA* 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate



vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) *J. Virol.* 61:1647-1650; Bender, M.A. et al. (1987) *J. Virol.* 61:1639-1646; Adam, M.A. and A.D. Miller (1988) *J. Virol.* 62:3802-3806; Dull, T. et al. (1998) *J. Virol.* 72:8463-8471; Zufferey, R. et al. (1998) *J. Virol.* 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) *J. Virol.* 71:7020-7029; Bauer, G. et al. (1997) *Blood* 89:2259-2267; Bonyhadi, M.L. (1997) *J. Virol.* 71:4707-4716; Ranga, U. et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:1201-1206; Su, L. (1997) *Blood* 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding PPIM to cells which have one or more genetic abnormalities with respect to the expression of PPIM. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) *Transplantation* 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) *Annu. Rev. Nutr.* 19:511-544; and Verma, I.M. and N. Somia (1997) *Nature* 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding PPIM to target cells which have one or more genetic abnormalities with respect to the expression of PPIM. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing PPIM to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction

and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding PPIM to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for PPIM into the alphavirus genome in place of the capsid-coding region results in the production of a large number of PPIM-coding RNAs and the synthesis of high levels of PPIM in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of PPIM into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme

molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding PPIM.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by  
5 scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using  
10 ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences  
15 encoding PPIM. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of  
20 the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

25 An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding PPIM. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular  
30 chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased PPIM expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding PPIM may be therapeutically useful, and in the treatment of disorders associated with  
35 decreased PPIM expression or activity, a compound which specifically promotes expression of the

polynucleotide encoding PPIM may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in  
5 altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding PPIM is exposed to at least one test compound thus obtained. The sample may comprise, for  
10 example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding PPIM are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding PPIM. The amount of hybridization may be quantified, thus forming the  
15 basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al.  
20 (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al.  
25 (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using  
30 methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

35 An additional embodiment of the invention relates to the administration of a composition which

generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of PPIM, antibodies to PPIM, and mimetics, agonists, antagonists, or inhibitors of PPIM.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising PPIM or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, PPIM or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example PPIM or fragments thereof, antibodies of PPIM, and agonists, antagonists or inhibitors of PPIM, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by

standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

10 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu$ g to 100,000  $\mu$ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

**DIAGNOSTICS**

In another embodiment, antibodies which specifically bind PPIM may be used for the diagnosis of disorders characterized by expression of PPIM, or in assays to monitor patients being treated with PPIM or agonists, antagonists, or inhibitors of PPIM. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for PPIM include methods which utilize the antibody and a label to detect PPIM in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring PPIM, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PPIM expression. Normal or standard values for PPIM expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to PPIM under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by

various methods, such as photometric means. Quantities of PPIM expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding PPIM may be used for  
5 diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of PPIM may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of PPIM, and to monitor regulation of PPIM levels during therapeutic intervention.

10 In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding PPIM or closely related molecules may be used to identify nucleic acid sequences which encode PPIM. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the  
15 probe identifies only naturally occurring sequences encoding PPIM, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the PPIM encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:28-54 or from genomic sequences including promoters, enhancers, and introns of the PPIM gene.

20 Means for producing specific hybridization probes for DNAs encoding PPIM include the cloning of polynucleotide sequences encoding PPIM or PPIM derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups,  
25 for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding PPIM may be used for the diagnosis of disorders associated with expression of PPIM. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed  
30 connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate,  
35 salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and an autoimmune/inflammatory

disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anenmia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic  
5 dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid  
10 arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma. The polynucleotide sequences encoding PPIM may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR  
15 technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered PPIM expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding PPIM may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide  
20 sequences encoding PPIM may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding PPIM in the sample indicates the  
25 presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of PPIM, a normal or standard profile for expression is established. This may be accomplished by combining body  
30 fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PPIM, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from  
35 patients who are symptomatic for a disorder. Deviation from standard values is used to establish the



presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays  
5 may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ  
10 preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding PPIM may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding PPIM, or  
15 a fragment of a polynucleotide complementary to the polynucleotide encoding PPIM, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences  
20 encoding PPIM may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding PPIM are used to amplify DNA using the polymerase chain reaction  
25 (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally,  
30 sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass  
35 spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San

Diego CA).

Methods which may also be used to quantify the expression of PPIM include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for PPIM, or PPIM or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies,

or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of  
5 pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to  
10 that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The  
15 normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at  
20 <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present  
25 invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present  
30 invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and analyzing the  
35 polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-

dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as

5 Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard

10 methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for PPIM to quantify the

15 levels of PPIM expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-

20 reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the

25 analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological

30 sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

35 In another embodiment, the toxicity of a test compound is assessed by treating a biological

sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding PPIM may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, e.g., Landcr, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding PPIM on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal

associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to  
5 that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, PPIM, its catalytic or immunogenic fragments, or  
10 oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between PPIM and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds  
15 having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with PPIM, or fragments thereof, and washed. Bound PPIM is then detected by methods well known in the art. Purified PPIM can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively,  
20 non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PPIM specifically compete with a test compound for binding PPIM. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PPIM.

25 In additional embodiments, the nucleotide sequences which encode PPIM may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding  
30 description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, in particular U.S. Ser. No. 60/147,986 and U.S. Ser. No. 60/160,807, are hereby expressly incorporated by  
35 reference.

## EXAMPLES

### I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERSCRIP<sup>T</sup> plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

### II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled

water and stored, with or without lyophilization, at 4 °C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

### III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows.

Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (PE Biosystems) thermal cyclor or the PTC-200 thermal cyclor (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (PE Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences



and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation

5 using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the

10 GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and

15 amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:28-54. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

#### IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene

20 and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much

25 faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum} \{ \text{length}(\text{Seq. 1}), \text{length}(\text{Seq. 2}) \}}$$

30

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a

35 score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every

mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding PPIM occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

#### **V. Chromosomal Mapping of PPIM Encoding Polynucleotides**

The cDNA sequences which were used to assemble SEQ ID NO:28-54 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:28-54 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO., to that map location.

The genetic map locations of SEQ ID NO:30, SEQ ID NO:37, and SEQ ID NO:47 are described in The Invention as ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. A disease associated with the public and Incyte sequences located within the indicated interval is also reported in the Invention.

## VI. Extension of PPIM Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:28-54 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other  
5 primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

10 Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and  $\beta$ -  
15 mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2:  
20 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar,  
25 Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested  
30 with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with  
35 Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E.*

coli cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems).

In like manner, the polynucleotide sequences of SEQ ID NO:28-54 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### 15 VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:28-54 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

#### VIII. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical

microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schna (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate

5 using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schna, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may

10 comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence

15 scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### Tissue or Cell Sample Preparation

20 Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse

25 transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified

30 using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

#### 35 Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 µg.

5 Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated  
10 with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 µl of the array element DNA, at an average concentration of 100 ng/µl, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

15 Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

#### 20 Hybridization

Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly  
25 larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

#### Detection

30 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-  
35 scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a

resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate  
5 filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a  
10 cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed,  
15 the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a  
20 linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot  
25 is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### **IX. Complementary Polynucleotides**

Sequences complementary to the PPIM-encoding sequences, or any parts thereof, are used to  
30 detect, decrease, or inhibit expression of naturally occurring PPIM. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of PPIM. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent  
35 promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is

designed to prevent ribosomal binding to the PPIM-encoding transcript.

#### X. Expression of PPIM

Expression and purification of PPIM is achieved using bacterial or virus-based expression systems. For expression of PPIM in bacteria, cDNA is subcloned into an appropriate vector containing  
5 an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express PPIM upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of  
10 PPIM in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding PPIM by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA  
15 transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, PPIM is synthesized as a fusion protein with, e.g., glutathione S-  
20 transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from PPIM at  
25 specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified PPIM obtained by these methods can be used directly in the assays shown in Examples XI and  
30 XV.

#### XI. Demonstration of PPIM Activity

Protease activity of PPIM is measured by the hydrolysis of appropriate synthetic peptide substrates conjugated with various chromogenic molecules. The degree of hydrolysis is quantified by spectrophotometric (or fluorometric) absorption of the released chromophore (Beynon, R.J. and J.S.



Bond (1994) Proteolytic Enzymes: A Practical Approach, Oxford University Press, New York NY, pp.25-55). Peptide substrates are designed according to the category of protease activity as endopeptidase (serine, cysteine, aspartic proteases), aminopeptidase (leucine aminopeptidase), or carboxypeptidase (Carboxypeptidase A and B, procollagen C-proteinase). Chromogens commonly used  
5 are 2-naphthylamine, 4-nitroaniline, and furylacrylic acid. Assays are performed at ambient temperature using an aliquot of PPIM and the appropriate substrate in a suitable buffer. Reactions are carried out in an optical cuvette and followed by the measurement of increase/decrease in absorbance of the chromogen released during hydrolysis of the peptide substrate. The change in absorbance is proportional to PPIM activity in the assay.

## 10 XII. Functional Assays

PPIM function is assessed by expressing the sequences encoding PPIM at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which  
15 contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector.  
20 Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA  
25 content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the  
30 cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of PPIM on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding PPIM and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin  
35 G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads

coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding PPIM and other genes of interest can be analyzed by northern analysis or microarray techniques.

### 5 XIII. Production of PPIM Specific Antibodies

PPIM substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the PPIM amino acid sequence is analyzed using LASERGENE software  
10 (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A  
15 peptide synthesizer (PE Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide and anti-PPIM activity by, for example, binding the peptide or PPIM to a substrate, blocking with 1% BSA, reacting  
20 with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

### XIV. Purification of Naturally Occurring PPIM Using Specific Antibodies

Naturally occurring or recombinant PPIM is substantially purified by immunoaffinity chromatography using antibodies specific for PPIM. An immunoaffinity column is constructed by covalently coupling anti-PPIM antibody to an activated chromatographic resin, such as CNBr-activated  
25 SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing PPIM are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PPIM (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/PPIM binding (e.g.,  
30 a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and PPIM is collected.

### XV. Identification of Molecules Which Interact with PPIM

PPIM, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules

previously arrayed in the wells of a multi-well plate are incubated with the labeled PPIM, washed, and any wells with labeled PPIM complex are assayed. Data obtained using different concentrations of PPIM are used to calculate values for the number, affinity, and association of PPIM with the candidate molecules.

- 5           Alternatively, molecules interacting with PPIM are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, Nature 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

PPIM may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions  
10 between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.

- 15 Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	28	088718	LIVRNOT01	088718H1 (LIVRNOT01), 151754F1 (FIBRAGT01), 151754R1 (FIBRAGT01), SCEA00861V1, SCEA01403V1, SCEA03107V1, SCEA01683V1
2	29	114551	TESTNOT01	1273531F1 (TESTTUT02), 1498122F1 (SINTBST01), 1686926F6 (PROSNOT15), 1922870R6 (BRSTTUT01), 2270121R6 (UTRSNOT02), 3227104F6 (COTRNOT01)
3	30	1261376	SYNORAT05	428341R6 (BLADNOT01), 488402R6 (HNT2AGT01), 1261376H1 (SYNORAT05), 1261376T6 (SYNORAT05), 1413230F6 (BRAINOT12), 1448134F1 (PLACNOT02), 1869342F6 (SKINBIT01), 2263303H1 (UTRSNOT02), 2365444T6 (ADRENOT07), 2875019H1 (THYRNOT10), 2908347H1 (THYMNOT05), 3818352H1 (BONSTUT01), g3840298, g1965665, g848456
4	31	1299481	BRSTNOT07	1299481H1 (BRSTNOT07), 1302262F6 (PLACNOT02), 1596742X330D1 (BRAINOT14), 1725693F6 (PROSNOT14), 2125677X306D3 (BRSTNOT07), SCHAO2258V1, SCHAO0613V1, g1477302
5	32	1873139	LEUKNOT02	003818R1 (HMCINOT01), 1873139F6 (LEUKNOT02), 1873139X325D1 (LEUKNOT02), 1873139X326V1 (LEUKNOT02), 1899870F6 (BLADTUT06), 2510118F6 (CONUTUT01)
6	33	1903112	OVARNOT07	1903112H1 (OVARNOT07), 1905330T6 (OVARNOT07), 2509325H1 (CONUTUT01), 2621121R6 (KERANOT02)
7	34	1993044	CORPNOT02	1858513F6 (PROSNOT18), 1993044H1 (CORPNOT02), 3733554F6 (SMCCNOS01), 4749046H1 (SMCRUNT01), 4960159H1 (TLYMNOT05), 5397428H1 (LIVRTUT13), SBCAO7095F3
8	35	2292182	BRAINON01	2199554H1 (SPINFET02), 2199554X305B1 (SPINFET02), 2292182R6 (BRAINON01), 3480414T6 (OVARNOT11), 5427954H1 (THYMTUT03)
9	36	2331301	COLNNOT11	1253717H1 (LUNGFET03), 2331301H1 (COLNNOT11), 2331301R6 (COLNNOT11)
10	37	2517512	BRAITUT21	1222614R1 (COLNUTUT02), 1486943F6 (UCMCL5T01), 1486943T6 (UCMCL5T01), 1569195F1 (UTRSNOT05), 1813007F6 (PROSTUT12), 2517512H1 (BRAITUT21), 5847584H1 (BRAENOT04)
11	38	3489039	EPIGNOT01	2541141F6 (BONRTUT01), 3489039H1 (EPIGNOT01), 4871852H1 (COLDNOT01)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
12	39	5432879	SPLNOT17	1429082F6 (SINTBST01), 1807480F6 (SINTNOT13), 2303440H1 (BRSTNOT05), 2669584F6 (ESOGTUT02), 3073745H1 (BONEUNT01), 3190142R6 (THYMNON04), 4693457H2 (BRAENOT02), 4774453H1 (BRAQNOT01), 5432879H1 (SPLNOT17), 9836070
13	40	5853753	FIBAUNT02	834033T1 (PROSNOT07), 1521711F6 (BLADTUT04), 1757751R6 (PITUNOT03), 2161634F6 (ENDCNOT02), SAEA01666R1, SCGA11716V1, SCGA05971V1, SCGA07285V1
14	41	411344	BRSTNOT01	411344F1 (BRSTNOT01), 411344H1 (BRSTNOT01), 411344R1 (BRSTNOT01), 1859850F6 (PROSNOT18), 2183379F6 (SININOT01), 2474963H1 (SMCANOT01), 2546619X300D1 (UTRSNOT11), 3728811H1 (SMCCNON03), 3932959H1 (PROSTUT09)
15	42	1256390	MENITUT03	1256390H1 (MENITUT03), SBAA04311F1, SBAA04104F1, SBAA03263F1, SBAA01188F1
16	43	1786774	BRAINOT10	857246H1 (NGANNOT01), 1786774F6 (BRAINOT10), 1786774H1 (BRAINOT10), 1810671T6 (PROSTUT12), 5202653H1 (STOMNOT08)
17	44	1911808	CONNTUT01	1255942F6 (MENITUT03), 1354692F6 (LUNGNOT09), 1354692T1 (LUNGNOT09), 1418156T1 (KIDNNOT09), 1436123F6 (PANCNOT08), 1498302T1 (SINTBST01), 1735923X304D1 (COLNNOT22), 1735923X318D4 (COLNNOT22), 1834236R6 (BRAINON01), 1911808F6 (CONNTUT01), 1911808H1 (CONNTUT01), 2360308H1 (LUNGFET05), 3075823H1 (BONEUNT01), 4106766H1 (BRSTTUT17), 5713020H1 (MASTXT01)
18	45	1973875	UCMCL5T01	1220149R6 (NEUTGMT01), 1377281F1 (LUNGNOT10), 1377281T1 (LUNGNOT10), 1508602F6 (LUNGNOT14), 1973875H1 (UCMCL5T01), 5098879F6 (EPIMNON05)
19	46	2323917	OVARNOT02	1609987F6 (COLNTUT06), 2012426R6 (TESTNOT03), 2012426T6 (TESTNOT03), 2323917H1 (OVARNOT02), 2323917T6 (OVARNOT02), 4851027H1 (TESTNOT10)
20	47	2754960	THPIAZS08	039061R6 (HUVENOB01), 580098H1 (BRAVTXT05), 2025465H1 (KERANOT02), 2754960H1 (THPIAZS08), 2754960R6 (THPIAZS08), 2754960X11F1 (THPIAZS08), 2754960X15F1 (THPIAZS08), 2754960X310U1 (THPIAZS08), 2754960X50F1 (THPIAZS08), 3821989T6 (BONSTUT01), 93736615

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
21	48	3092341	BRSTNOT19	3092341H1 (BRSTNOT19), 3092341T6 (BRSTNOT19)
22	49	3658034	ENDPNOT02	2623516R6 (KERANOT02), 3658034F6 (ENDPNOT02), 3658034H1 (ENDPNOT02), 3658034T6 (ENDPNOT02), 5216522H1 (BRSTNOT35), 5590053H1 (ENDINOT02)
23	50	3883861	UTRSNOT05	858111H1 (NGANNOT01), 858233H1 (NGANNOT01), 1364808R1 (SCORNON02), 1861181F6 (PROSNOT19), 1906985T6 (OVARNOT07), 2687868H1 (LJUNGNOT23), 2687868X366D1 (LJUNGNOT23), 2721116X369D1 (LJUNGTUT10), 3883861H1 (UTRSNOT05), 5217169H1 (BRSTNOT35)
24	51	4993873	LIVRTUT11	4987943H1 (LIVRTUT10), 4993873H1 (LIVRTUT11), SCEA01665V1, SCEA00232V1, SXBC01625V1, SXBC01802V1, SCSA03627V1
25	52	5208004	BRAFNOT02	4696870F6 (BRALNOT01), 5208004H1 (BRAFNOT02)
26	53	5267783	BRAFDIT02	220636R1 (STOMNOT01), 679457R6 (UTRSNOT02), 1330537F6 (PANCNOT07), 1808720F6 (PROSTUT12), 1969475H1 (BRSTNOT04), 2697426F6 (UTRSNOT12), 2991180H1 (KIDNFET02), 3532849H1 (KIDNNOT25), 4992376F6 (LIVRTUT11), 5004695F6 (PROSTUT21), 5267783H1 (BRAFDIT02)
27	54	5583922	FIBAUNT01	726878R1 (SYNOOAT01), 956818X11 (KIDNNOT05), 1658964X12 (URETTUT01), 1658964X13 (URETTUT01), 2544879F6 (UTRSNOT11), 3748858H1 (UTRSNOT18), 4761921H1 (PLACNOT05), 5043801H1 (PLACFER01)

Table 2

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences	Homologous Sequence	Analytical Methods and Databases
1	444	S91 T244 T251 S277 T386 T38 S182 T263 T373 Y346	N36 N180 N197 N295	Signal_peptide: M1-A23 Serpins (serine protease inhibitors): M1-P441, L68-L444	g1397241 RASPI	Motifs BLAST-GenBank HMMER SPScan HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST_PRODOR BLAST_DOMO
2	565	S9 S19 T343 T458 S5 S58 S82 S114 S184 S185 S295 T382 T432 T476 T495 T543 S2 S5 S12 S25 S42 T169 S307 T337 S352 T357 T426 S513 T523 Y220 Y514	N112 N494	Ubiquitin carboxyl-terminal hydrolases family 2: G226-L243, Y235-I549	g2746775 Similar to peptidase family C19 (ubiquitin carboxyl-terminal peptidase)	Motifs BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS BLAST_PRODOR BLAST_DOMO
3	589	T43 S71 S181 S200 S260 S304 S312 T506 T572 T40 S66	N55 N126 N136 N164 N167 N302 N501	Ubiquitin family signature: M37-K107 Ubiquitin-associated domain: Q541-S586	g3873621 Similar to ubiquitin family	Motifs BLAST-GenBank HMMER-PFAM
4	775	T305 T2 S27 S43 S67 S392 S611 S615 T647 S665 S710 S729 S759 S96 T106 S217 S288 S301 S316 S432 S438 T443 S575 T719 S723 Y334	N49 N215 N322 N387 N468 N487 N497 N504 N508 N568 N600	Ubiquitin carboxyl-terminal hydrolases family 2: G112-L129, G193-L202, V230-C244, Y354-V391, N380-S401 Ubiquitin hydrolase carboxyl-terminal thiolesterase: G112-K211	g2739431 Hematopoietic-specific IL-2 deubiquitinating enzyme	Motifs BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS BLAST_PRODOR BLAST_DOMO

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences	Homologous Sequence	Analytical Methods and Databases
5	351	S9 S41 S48 S194 S201 T203 T257 S278 T322 T324 S129 S162 S181 S194 S225 T226 S348 Y271	N46 N123 N317	Ubiquitin carboxyl-terminal hydrolases family 2: L49-L337	g5410230 Ubiquitin-specific protease 3	Motifs BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS BLAST-PRODOM BLAST-DOMO
6	136	T30 S104 Y98		Dipeptidyl serine protease iv: I9-S128 Serine family prolyl endopeptidase: M4-I136	g577284 Dipeptidyl peptidase IV	Motifs BLAST-GenBank BLAST-PRODOM BLAST-DOMO
7	396	S24 S139 T168 T177 S198 S223 S279 T369 S26 S60 S223 S292	N166	Ubiquitin carboxyl-terminal hydrolase: E74-I283	g2854121 BRCA1 associated protein 1	Motifs BLAST-GenBank BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO HMMER-PFAM
8	246	S87 Y65	N94 N156 N195 N225	Zinc-binding metalloprotease domain: R121-H133		Motifs HMMER-PFAM
9	262	T32 S78 S85 T89 S125 S26 S170 S244	N168	Inter-alpha-trypsin glycoprotein inhibitor precursor: T32-T197		Motifs BLAST-PRODOM
10	406	S18 S37 T80 S98 S112 S178 T292 S298 T320 T391 S105 S212 S220 Y213	N14 N56 N176 N318		g3309170 COP9 complex subunit 4	Motifs BLAST-GenBank



Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences	Homologous Sequence	Analytical Methods and Databases
11	172	T117 S135 S146		signal peptide motif: M1-G13 ATP-binding kinase: I6-E164 AAA-protein family: P4-M69	g3875433 Similar to ATP binding protein	Motifs BLAST-GenBank SPScan BLAST-PRODOM BLAST-DOMO
12	517	S485 S4 T11 S128 T133 S155 S156 S171 S172 S278 T288 S485 S3 T57 T199 T204 S278 T455 S462 T480	N286	Ubiquitin carboxyl-terminal hydrolases family 2: K61-P256, F436-V481, S470-S491	g2459395 Ubiquitin protease	Motifs BLAST-GenBank HMMER-PFAM BLIMPS-BLOCKS BLAST-PRODOM
13	346	T237 S12 T64 T72 T124 T236 T261 S319 S150 T194 S226 T251 S319		Ubiquitin-activating enzyme signature: S297-344, 11-163, 19-189, 7-174, 9-192, R35-G170 Membrane protein: 11-249	g3647283 Ubiquitin activating enzyme	Motifs BLAST-GenBank HMMER BLAST-PRODOM BLAST-DOMO
14	151	T24 T47 S118 S61 Y131		Ubiquitin-conjugating enzymes: M1-D148 Active site: F58-M115	g4090259 Ubiquitin-conjugating enzyme E2	Motifs BLAST-GenBank HMMER-PFAM ProfileScan BLAST-PRODOM BLAST-DOMO
15	362	S199 S208 S212 S270 S281 T317 S327 S52 S122 T149	N120 N162 N175 N239	Signal peptide: M1-S26 Zinc carboxypeptidase: Y38-E320 Zinc binding region: E202-L258	g6013463 Carboxypeptidase homolog	Motifs BLAST-GenBank SPScan HMMER-PFAM ProfileScan BLAST-PRODOM BLAST-DOMO

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences	Homologous Sequence	Analytical Methods and Databases
16	123	S2 S9 T37 T46 T60 S112 T53 S112	N104	Kunitz type protease inhibitor active site region: C70-C120	g512802 Kunitz type protease inhibitor	Motifs BLAST-GenBank HMMER-PFAM ProfileScan BLAST-DOMO
17	983	S87 S461 S531 T761 T123 T143 S191 S445 S634 S660 T789 T820 S879 S886 S888 T890 T17 S158 T280 T398 T549 S598 S601 S687 Y268 Y688	N278 N427 N625 N884 N922	Ubiquitin carboxyl-terminal hydrolases family 2: G90-w107, Y336-I374	gl429371 Ubiquitin-specific protease	Motifs BLAST-GenBank HMMER-PFAM BLAST-DOMO
18	227	S49 T101 T131 T157 S166 S49 S144 S194 T199		Ubiquitin signature: K159-H179, A180-D200 (P value = 0.00032)	g9372 Ubiquitin (P value = 1.7e <sup>08</sup> )	Motifs BLAST-GenBank BLIMPS-PRINTS
19	403	T47 S146 T261 T352 T381 S4 T119 S234 S291 S313	N117 N145 N232 N260 N289 N317	Ubiquitin carboxyl-terminal hydrolases family 2: G221-L238	g4731026 Nod1 activator of caspase-9 and NFKB	Motifs BLAST-GenBank HMMER-PFAM
20	372	T87 S291 S22 S197 T208 S343 T169 S185 S223 S260 T266	N188 N335	Ubiquitin carboxyl-terminal hydrolases family 2: A166-Q348 Active site: Y302-C320	g4469352 Ubiquitin specific protease UBP43	Motifs BLAST-GenBank HMMER-PFAM BLAST-DOMO
21	94	T9	N50	Signal peptidase: V41-R55	g3687497 Putative mitochondrial inner membrane protease subunit	Motifs BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences	Homologous Sequence	Analytical Methods and Databases
22	248	S77 S135 S156 S183 S205 T3 S71 S72 T139	N47 N158	Alpha-2-macroglobulin family: T3-Y198 Complement precursor: E4-S206	g2073373 Alpha-2-macroglobulin protease inhibitor	Motifs BLAST-GenBank HMMER-PFAM BLAST-PRODOM
23	520	S166 S272 T301 S326 S379 S455 S56 T82 S136 S227 S498	N164 N355	Signal peptide: M1-R27 Peptidase M10: F39-S225 Matrixin domain: F128-G288 Neutral zinc metalloproteinase binding region: V237-L246 Hemopexin domain: I341-K400	g1731986 MMP-19 matrix metalloproteinase	Motifs BLAST-GenBank SIGPEPT SPScan HMMER-PFAM ProfileScan BLAST-PRODOM BLAST-DMO
24	422	T188 S156 S306 T386 S130 T176 T226 T295 S357 S365	N94 N106 N169 N350	Signal peptide: M1-G26 Transmembrane domain: F398-N418 Serpins (serine protease inhibitors): P43-V420 Protease "bait" region: A371-G422	g425146 Kallistatin	Motifs BLAST-GenBank SIGPEPT SPScan HMMER HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DMO
25	114	S74 S16		Eukaryotic thiol (cysteine) protease active site: R71-S114		Motifs BLAST-GenBank ProfileScan

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences	Homologous Sequence	Analytical Methods and Databases
26	742	T167 S186 S308 S337 S343 T360 S439 S578 S92 S172 S239 T256 T278 S329 T414 S504 S633 T656 T708 Y28 Y107 Y356		Zinc carboxypeptidases, zinc-binding regions signatures: H32-W42		Motifs BLAST-GenBank
27	734	T83 S128 S151 S223 S233 T523 S574 T616 T665 T688 T34 S122 S203 S340 T546 S547 T703	N57 N210 N220 N318 N428 N472	Signal peptide: M1-G20 Zinc carboxypeptidases: H299-Y412, W421-Y678 Enkephalin convertase: P458-V687 Zinc binding region: E478-F529	g4322263 Metallocarboxy-peptidase CPX-1	Motifs BLAST-GenBank SIGPEPT SPScan HMMER-PFAM ProfileScan BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO

Table 3

Nucleotide SEQ ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
28	164-208	Gastrointestinal (1.000)	Inflammation (0.500)	PBLUESCRIPT
29	57-101	Reproductive (0.274) Nervous (0.202) Cardiovascular (0.119) Gastrointestinal (0.119)	Cancer (0.524) Inflammation (0.273) Cell Proliferation(0.190)	PBLUESCRIPT
30	111-155	Nervous (0.222) Reproductive (0.194) Gastrointestinal (0.139)	Cancer (0.403) Inflammation (0.361)	PSPORT1
31	921-965	Nervous (0.300) Reproductive (0.200) Cardiovascular (0.100) Dermatologic (0.100) Developmental (0.100) Gastrointestinal (0.100) Hematopoietic/Immune (0.100)	Cancer (0.400) Cell Proliferation(0.300) Neurological (0.100)	pINCY
32	809-853	Hematopoietic/Immune (0.194) Reproductive (0.239) Gastrointestinal (0.164)	Cancer (0.403) Inflammation (0.269) Cell Proliferation(0.134)	pINCY
33	273-317	Reproductive (0.500) Cardiovascular (0.125) Dermatologic (0.125) Gastrointestinal (0.125) Hematopoietic/Immune (0.125)	Cancer (0.625) Cell Proliferation(0.125) Inflammation (0.125)	pINCY
34	55-99	Nervous (0.185) Cardiovascular (0.111) Gastrointestinal (0.111)	Cancer (0.352) Inflammation (0.204) Cell Proliferation(0.204)	pINCY
35	218-262	Gastrointestinal (0.313) Hematopoietic/Immune (0.250) Reproductive (0.188)	Cancer (0.630) Cell Proliferation(0.250)	PSPORT1
36	325-369	Developmental (0.500) Gastrointestinal (0.500)	Cancer (0.500) Cell Proliferation(0.500)	PSPORT1
37	99-143	Nervous (0.198) Reproductive (0.165) Cardiovascular (0.154)	Cancer (0.374) Inflammation (0.374) Cell Proliferation(0.154)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
38	1-46	Reproductive (0.278) Gastrointestinal (0.208) Cardiovascular (0.125)	Cancer (0.347) Inflammation (0.306) Cell Proliferation (0.153)	pINCY
39	109-153	Gastrointestinal (0.280) Hematopoietic/Immune (0.200) Musculoskeletal (0.120)	Inflammation (0.440) Cancer (0.280) Cell Proliferation (0.160)	pINCY
40	489-533	Nervous (0.209) Reproductive (0.203) Gastrointestinal (0.135)	Cancer (0.473) Cell Proliferation (0.243) Inflammation (0.264)	pINCY
41	589-633	Reproductive (0.229) Cardiovascular (0.200) Gastrointestinal (0.171) Nervous (0.171)	Cancer (0.314) Cell Proliferation (0.314) Inflammation/Trauma (0.372)	PBLUESCRIPT
42	649-693	Nervous (0.250) Reproductive (0.214) Cardiovascular (0.143)	Cancer (0.500) Inflammation/Trauma (0.321) Cell Proliferation (0.179)	pINCY
43	164-208	Nervous (0.333) Gastrointestinal (0.333) Reproductive (0.333)	Cancer (0.444) Inflammation/Trauma (0.444) Neurological (0.111)	pINCY
44	271-208	Reproductive (0.226) Developmental (0.151) Nervous (0.151)	Cancer (0.377) Inflammation/Trauma (0.358) Cell Proliferation (0.321)	pINCY
45	784-828	Reproductive (0.257) Hematopoietic/Immune (0.171) Nervous (0.171)	Cancer (0.486) Inflammation/Trauma (0.486) Cell Proliferation (0.143)	PBLUESCRIPT
46	219-263	Reproductive (0.444) Gastrointestinal (0.222) Nervous (0.222)	Inflammation/Trauma (0.666) Cancer (0.222)	PSPORT1
47	597-641	Reproductive (0.364) Cardiovascular (0.212) Nervous (0.152)	Cancer (0.545) Cell Proliferation (0.242) Inflammation/Trauma (0.273)	PSPORT1

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
48	271-315	Gastrointestinal (0.278) Reproductive (0.278) Cardiovascular (0.111) Hematopoietic/Immune (0.111) Nervous (0.111)	Cancer (0.444) Inflammation/Trauma (0.555) Cell Proliferation (0.167)	pINCY
49	217-261	Hematopoietic/Immune (0.364) Reproductive (0.273)	Cell Proliferation (0.364) Inflammation/Trauma (0.364) Cancer (0.182)	pINCY
50	164-208	Reproductive (0.333) Nervous (0.222) Gastrointestinal (0.167)	Cancer (0.611) Inflammation/Trauma (0.223)	pINCY
51	388-432	Gastrointestinal (0.833) Reproductive (0.166)	Cancer (0.666) Cell Proliferation (0.166)	pINCY
52	218-262	Nervous (0.750) Hematopoietic/Immune (0.250)	Inflammation/Trauma (0.500) Neurological (0.250)	pINCY
53	325-369	Reproductive (0.289) Nervous (0.253) Gastrointestinal (0.120)	Cancer (0.410) Inflammation/Trauma (0.386) Cell Proliferation (0.145)	pINCY
54	165-209	Reproductive (0.352) Urologic (0.185) Developmental (0.130)	Cancer (0.630) Cell Proliferation (0.167) Inflammation/Trauma (0.204)	pINCY

Table 4

Polynucleotide SEQ ID NO:	Library	Library Comment
28	LIVRNOT01	Library was constructed at Stratagene, using RNA isolated from the liver tissue of a 49-year-old male.
29	TESTNOT01	Library was constructed using RNA isolated from the testicular tissue of a 37-year-old Caucasian male, who died from liver disease. Patient history included cirrhosis, jaundice, and liver failure.
30	SYNORAT05	Library was constructed using RNA isolated from the knee synovial tissue of a 62-year-old female with rheumatoid arthritis.
31	BRSTNOT07	Library was constructed using RNA isolated from diseased breast tissue removed from a 43-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated mildly proliferative fibrocystic changes with epithelial hyperplasia, papillomatosis, and duct ectasia. Pathology for the associated tumor tissue indicated invasive grade 4, nuclear grade 3 mammary adenocarcinoma with extensive comedo necrosis. Family history included epilepsy, cardiovascular disease, and type II diabetes.
32	LEUKNOT02	Library was constructed using RNA isolated from white blood cells of a 45-year-old female with blood type O+. The donor tested positive for cytomegalovirus (CMV).
33	OVARNOT07	Library was constructed using RNA isolated from left ovarian tissue removed from a 28-year-old Caucasian female during a vaginal hysterectomy and removal of the fallopian tubes and ovaries. The tissue was associated with multiple follicular cysts, endometrium in a weakly proliferative phase, and chronic cervicitis of the cervix with squamous metaplasia. Family history included benign hypertension, hyperlipidemia, and atherosclerotic coronary artery disease.
34	CORPNOT02	Library was constructed using RNA isolated from diseased corpus callosum tissue removed from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
35	BRAINON01	Library was constructed and normalized from 4.88 million independent clones from a brain tissue library. RNA was made from brain tissue removed from a 26-year-old Caucasian male during cranioplasty and excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated a grade 4 oligoastrocytoma in the right fronto-parietal part of the brain.
36	COLNNOT11	Library was constructed using RNA isolated from colon tissue removed from a 60-year-old Caucasian male during a left hemicolectomy.



Table 4 (cont.)

Polynucleotide SEQ ID NO:	Library	Library Comment
37	BRAITUT21	Library was constructed using RNA isolated from brain tumor tissue removed from the midline frontal lobe of a 61-year-old Caucasian female during excision of a cerebral meningial lesion. Pathology indicated subfrontal meningothelial meningioma with no atypia. One ethmoid and mucosal tissue sample indicated meningioma. Family history included cerebrovascular disease, senile dementia, hyperlipidemia, benign hypertension, atherosclerotic coronary artery disease, congestive heart failure, and breast cancer.
38	EPIGNOT01	Library was constructed using RNA isolated from epiglottic tissue removed from a 71-year-old male during laryngectomy with right parathyroid biopsy. Pathology for the associated tumor tissue indicated recurrent grade 1 papillary thyroid carcinoma.
39	SPLNNOT17	Library was constructed using polyA RNA isolated from the spleen tissue of a 2-year-old Hispanic male who died from cerebral anoxia.
40	FIBAUNT02	Library was constructed using RNA isolated from untreated aortic adventitial fibroblasts removed from a 65-year-old Caucasian female.
41	BRSTNOT01	Library was constructed using RNA isolated from the breast tissue of a 56-year-old Caucasian female who died in a motor vehicle accident.
42	MENITUT03	Library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningial lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
43	BRAINOT10	Library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 74-year-old Caucasian male, who died from Alzheimer's disease.
44	CONNTUT01	Library was constructed using RNA isolated from a soft tissue tumor removed from the clival area of the skull of a 30-year-old Caucasian female. Pathology indicated chondroid chordoma with neoplastic cells reactive for keratin.
45	UCMCL5T01	Library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of 12 individuals. The cells were cultured for 12 days with IL-5 before RNA was obtained from the pooled lysates.
46	OVARNOT02	Library was constructed using RNA isolated from ovarian tissue removed from a 59-year-old Caucasian female who died of a myocardial infarction. Patient history included cardiomyopathy, coronary artery disease, previous myocardial infarctions, hypercholesterolemia, hypertension, and arthritis.

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Library	Library Comment
47	THPIAZS08	This subtracted THP-1 promonocyte cell line library was constructed using 5.76 million clones from a 5-aza-2'-deoxycytidine (AZ) treated THP-1 cell library. Starting RNA was made from THP-1 promonocyte cells treated for three days with 0.8 micromolar AZ. The hybridization probe for subtraction was derived from a similarly constructed library, made from RNA isolated from untreated THP-1 cells. 5.76 million clones from the AZ-treated THP-1 cell library were then subjected to two rounds of subtractive hybridization with 5 million clones from the untreated THP-1 cell library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (1991) Nucleic Acids Res. 19:1954; and Bonaldo et al. (1996) Genome Research 6:791. THP-1 (ATCC TIB 202) is a human promonocyte line derived from peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia.
48	BRSTNOT19	Library was constructed using RNA isolated from breast tissue removed from a 67-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated residual invasive lobular carcinoma. The focus of residual invasive carcinoma was positive for both estrogen and progesterone. Patient history included depressive disorder and benign large bowel neoplasm. Family history included cerebrovascular disease, benign hypertension, congestive heart failure, and lung cancer.
49	ENDPNOT02	Library was constructed using RNA isolated from pulmonary artery endothelial cells removed from a 10-year-old Caucasian male. The cells were treated with TNF alpha and IL-1 beta 10ng/ml each for 20 hours.
50	UTRSNOT05	Library was constructed using RNA isolated from the uterine tissue of a 45-year-old Caucasian female during a total abdominal hysterectomy and total colectomy. Pathology for the associated tumor tissue indicated multiple leiomyomas of the myometrium and a grade 2 colonic adenocarcinoma of the cecum. Patient history included multiple sclerosis and mitral valve disorder. Family history included type I diabetes, cerebrovascular disease, atherosclerotic coronary artery disease, malignant skin neoplasm, hypertension, and malignant neoplasm of the colon.

Table 4 (cont.)

Polynucleotide SEQ ID NO:	Library	Library Comment
51	LIVRTUT11	Library was constructed using 1.1 micrograms of polyA RNA isolated from a treated C3A hepatocyte cell line which is a derivative of Hep G2, a cell line derived from a hepatoblastoma removed from a 15-year-old Caucasian male. The cells were treated with phenobarbital (PB), 1mM for 48 hours. cDNA synthesis was initiated using a NotI-anchored oligo(dT) primer. Double-stranded cDNA was blunted, ligated to EcoRI adaptors, digested with NotI, size-selected, and cloned into the NotI and EcoRI sites of the pINCY vector (Incyte).
52	BRAFNOT02	The library was constructed using RNA isolated from superior frontal cortex tissue removed from a 35-year-old Caucasian male who died from cardiac failure. Pathology indicated moderate leptomeningeal fibrosis and multiple microinfarctions of the cerebral neocortex. Patient history included dilated cardiomyopathy, congestive heart failure, cardiomegaly, and an enlarged spleen and liver.
53	BRAFDIT02	The library was constructed using RNA isolated from diseased right frontal lobe tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident. Patient history included Huntington's disease and emphysema.
54	FIBAUNT01	Library was constructed using RNA isolated from untreated aortic adventitial fibroblasts obtained from a 48-year-old Caucasian male.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	PE Biosystems, Foster City, CA.	
ABI/PARACEL.FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	PE Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	PE Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score $\geq$ GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, and SEQ ID  
10 NO:27,

b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27,

c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27, and

d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27.

30 2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, and SEQ  
35 ID NO:27.

3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, and SEQ ID NO:54.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
9. A method for producing a polypeptide of claim 1, the method comprising:
- a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
- b) recovering the polypeptide so expressed.
10. An isolated antibody which specifically binds to a polypeptide of claim 1.
11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:
- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, and SEQ ID NO:54,
- b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a

polynucleotide sequence selected from the group consisting of SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, and SEQ ID NO:54,

- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

30

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4,



SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, and SEQ ID NO:27.

5

18. A method for treating a disease or condition associated with decreased expression of functional PPIM, comprising administering to a patient in need of such treatment the composition of claim 16.

10

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

15

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

20

21. A method for treating a disease or condition associated with decreased expression of functional PPIM, comprising administering to a patient in need of such treatment a composition of claim 20.

25

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

30

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

35

24. A method for treating a disease or condition associated with overexpression of functional PPIM, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable

conditions, and

b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

5           26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,

b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound,  
10 and

c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

15

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

a) exposing a sample comprising the target polynucleotide to a compound, and  
20 b) detecting altered expression of the target polynucleotide.

28. A method for assessing toxicity of a test compound, said method comprising:

a) treating a biological sample containing nucleic acids with the test compound;

b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at  
25 least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;

c) quantifying the amount of hybridization complex; and

30 d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

## SEQUENCE LISTING

&lt;110&gt; INCYTE GENOMICS, INC.

YUE, Henry  
 TANG, Y. Tom  
 BANDMAN, Olga  
 LAL, Preeti  
 BAUGHN, Mariah R.  
 AZIMZAI, Yalda  
 LU, Dyung Aina M.  
 YANG, Junming

&lt;120&gt; PROTEASES AND PROTEASE INHIBITORS

&lt;130&gt; PF-0727 PCT

&lt;140&gt; To Be Assigned

&lt;141&gt; Herewith

&lt;150&gt; 60/147,986; 60/160,807

&lt;151&gt; 1999-08-09; 1999-10-21

&lt;160&gt; 54

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 444

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 088718CD1

&lt;400&gt; 1

Met	Lys	Val	Val	Pro	Ser	Leu	Leu	Leu	Ser	Val	Leu	Leu	Ala	Gln
1				5					10					15
Val	Trp	Leu	Val	Pro	Gly	Leu	Ala	Pro	Ser	Pro	Gln	Ser	Pro	Glu
				20					25					30
Thr	Pro	Ala	Pro	Gln	Asn	Gln	Thr	Ser	Arg	Val	Val	Gln	Ala	Pro
				35					40					45
Arg	Glu	Glu	Glu	Glu	Asp	Glu	Gln	Glu	Ala	Ser	Glu	Glu	Lys	Ala
				50					55					60
Gly	Glu	Glu	Glu	Lys	Ala	Trp	Leu	Met	Ala	Ser	Arg	Gln	Gln	Leu
				65					70					75
Ala	Lys	Glu	Thr	Ser	Asn	Phe	Gly	Phe	Ser	Leu	Leu	Arg	Lys	Ile
				80					85					90
Ser	Met	Arg	His	Asp	Gly	Asn	Met	Val	Phe	Ser	Pro	Phe	Gly	Met
				95					100					105
Ser	Leu	Ala	Met	Thr	Gly	Leu	Met	Leu	Gly	Ala	Thr	Gly	Pro	Thr
				110					115					120
Glu	Thr	Gln	Ile	Lys	Arg	Gly	Leu	His	Leu	Gln	Ala	Leu	Lys	Pro
				125					130					135
Thr	Lys	Pro	Gly	Leu	Leu	Pro	Ser	Leu	Phe	Lys	Gly	Leu	Arg	Glu
				140					145					150
Thr	Leu	Ser	Arg	Asn	Leu	Glu	Leu	Gly	Leu	Ser	Gln	Gly	Ser	Phe
				155					160					165
Ala	Phe	Ile	His	Lys	Asp	Phe	Asp	Val	Lys	Glu	Thr	Phe	Phe	Asn
				170					175					180
Leu	Ser	Lys	Arg	Tyr	Phe	Asp	Thr	Glu	Cys	Val	Pro	Met	Asn	Phe
				185					190					195
Arg	Asn	Ala	Ser	Gln	Ala	Lys	Arg	Leu	Met	Asn	His	Tyr	Ile	Asn
				200					205					210
Lys	Glu	Thr	Arg	Gly	Lys	Ile	Pro	Lys	Leu	Phe	Asp	Glu	Ile	Asn
				215					220					225
Pro	Glu	Thr	Lys	Leu	Ile	Leu	Val	Asp	Tyr	Ile	Leu	Phe	Lys	Gly

Lys	Trp	Leu	Thr	Pro	Phe	Asp	Pro	Val	Phe	Thr	Glu	Val	Asp	Thr	230	235	240
															245	250	255
Phe	His	Leu	Asp	Lys	Tyr	Lys	Thr	Ile	Lys	Val	Pro	Met	Met	Tyr	260	265	270
Gly	Ala	Gly	Lys	Phe	Ala	Ser	Thr	Phe	Asp	Lys	Asn	Phe	Arg	Cys	275	280	285
His	Val	Leu	Lys	Leu	Pro	Tyr	Gln	Gly	Asn	Ala	Thr	Met	Leu	Val	290	295	300
Val	Leu	Met	Glu	Lys	Met	Gly	Asp	His	Leu	Ala	Leu	Glu	Asp	Tyr	305	310	315
Leu	Thr	Thr	Asp	Leu	Val	Glu	Thr	Trp	Leu	Arg	Asn	Met	Lys	Thr	320	325	330
Arg	Asn	Met	Glu	Val	Phe	Phe	Pro	Lys	Phe	Lys	Leu	Asp	Gln	Lys	335	340	345
Tyr	Glu	Met	His	Glu	Leu	Leu	Arg	Gln	Met	Gly	Ile	Arg	Arg	Ile	350	355	360
Phe	Ser	Pro	Phe	Ala	Asp	Leu	Ser	Glu	Leu	Ser	Ala	Thr	Gly	Arg	365	370	375
Asn	Leu	Gln	Val	Ser	Arg	Val	Leu	Gln	Arg	Thr	Val	Ile	Glu	Val	380	385	390
Asp	Glu	Arg	Gly	Thr	Glu	Ala	Val	Ala	Gly	Ile	Leu	Ser	Glu	Ile	395	400	405
Thr	Ala	Tyr	Ser	Met	Pro	Pro	Val	Ile	Lys	Val	Asp	Arg	Pro	Phe	410	415	420
His	Phe	Met	Ile	Tyr	Glu	Glu	Thr	Ser	Gly	Met	Leu	Leu	Phe	Leu	425	430	435
Gly	Arg	Val	Val	Asn	Pro	Thr	Leu	Leu							440		

&lt;210&gt; 2

&lt;211&gt; 565

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 114551CD1

&lt;400&gt; 2

Met	Ser	Gly	Arg	Ser	Lys	Arg	Glu	Ser	Arg	Gly	Ser	Thr	Arg	Gly	1	5	10	15
Lys	Arg	Glu	Ser	Glu	Ser	Arg	Gly	Ser	Ser	Gly	Arg	Val	Lys	Arg	20	25	30	35
Glu	Arg	Asp	Arg	Glu	Arg	Glu	Pro	Glu	Ala	Ala	Ser	Ser	Arg	Gly	35	40	45	50
Ser	Pro	Val	Arg	Val	Lys	Arg	Glu	Phe	Glu	Pro	Ala	Ser	Ala	Arg	50	55	60	65
Glu	Ala	Pro	Ala	Ser	Val	Val	Pro	Phe	Val	Arg	Val	Lys	Arg	Glu	65	70	75	80
Arg	Glu	Val	Asp	Glu	Asp	Ser	Glu	Pro	Glu	Arg	Glu	Val	Arg	Ala	80	85	90	95
Lys	Asn	Gly	Arg	Val	Asp	Ser	Glu	Asp	Arg	Arg	Ser	Arg	His	Cys	95	100	105	110
Pro	Tyr	Leu	Asp	Thr	Ile	Asn	Arg	Ser	Val	Leu	Asp	Phe	Asp	Phe	110	115	120	125
Glu	Lys	Leu	Cys	Ser	Ile	Ser	Leu	Ser	His	Ile	Asn	Ala	Tyr	Ala	125	130	135	140
Cys	Leu	Val	Cys	Gly	Lys	Tyr	Phe	Gln	Gly	Arg	Gly	Leu	Lys	Ser	140	145	150	155
His	Ala	Tyr	Ile	His	Ser	Val	Gln	Phe	Ser	His	His	Val	Phe	Leu	155	160	165	170
Asn	Leu	His	Thr	Leu	Lys	Phe	Tyr	Cys	Leu	Pro	Asp	Asn	Tyr	Glu	170	175	180	185
Ile	Ile	Asp	Ser	Ser	Leu	Glu	Asp	Ile	Thr	Tyr	Val	Leu	Lys	Pro	185	190	195	200
Thr	Phe	Thr	Lys	Gln	Gln	Ile	Ala	Asn	Leu	Asp	Lys	Gln	Ala	Lys	200	205	210	

```

Leu Ser Arg Ala Tyr Asp Gly Thr Thr Tyr Leu Pro Gly Ile Val
215 220 225
Gly Leu Asn Asn Ile Lys Ala Asn Asp Tyr Ala Asn Ala Val Leu
230 235 240
Gln Ala Leu Ser Asn Val Pro Pro Leu Arg Asn Tyr Phe Leu Glu
245 250 255
Glu Asp Asn Tyr Lys Asn Ile Lys Arg Pro Pro Gly Asp Ile Met
260 265 270
Phe Leu Leu Val Gln Arg Phe Gly Glu Leu Met Arg Lys Leu Trp
275 280 285
Asn Pro Arg Asn Phe Lys Ala His Val Ser Pro His Glu Met Leu
290 295 300
Gln Ala Val Val Leu Cys Ser Lys Lys Thr Phe Gln Ile Thr Lys
305 310 315
Gln Gly Asp Gly Val Asp Phe Leu Ser Trp Phe Leu Asn Ala Leu
320 325 330
His Ser Ala Leu Gly Gly Thr Lys Lys Lys Lys Lys Thr Ile Val
335 340 345
Thr Asp Val Phe Gln Gly Ser Met Arg Ile Phe Thr Lys Lys Leu
350 355 360
Pro His Pro Asp Leu Pro Ala Glu Glu Lys Glu Gln Leu Leu His
365 370 375
Asn Asp Glu Tyr Gln Glu Thr Met Val Glu Ser Thr Phe Met Tyr
380 385 390
Leu Thr Leu Asp Leu Pro Thr Ala Pro Leu Tyr Lys Asp Glu Lys
395 400 405
Glu Gln Leu Ile Ile Pro Gln Val Pro Leu Phe Asn Ile Leu Ala
410 415 420
Lys Phe Asn Gly Ile Thr Glu Lys Glu Tyr Lys Thr Tyr Lys Glu
425 430 435
Asn Phe Leu Lys Arg Phe Gln Leu Thr Lys Leu Pro Pro Tyr Leu
440 445 450
Ile Phe Cys Ile Lys Arg Phe Thr Lys Asn Asn Phe Phe Val Glu
455 460 465
Lys Asn Pro Thr Ile Val Asn Phe Pro Ile Thr Asn Val Asp Leu
470 475 480
Arg Glu Tyr Leu Ser Glu Glu Val Gln Ala Val His Lys Asn Thr
485 490 495
Thr Tyr Asp Leu Ile Ala Asn Ile Val His Asp Gly Lys Pro Ser
500 505 510
Glu Gly Ser Tyr Arg Ile His Val Leu His His Gly Thr Gly Lys
515 520 525
Trp Tyr Glu Leu Gln Asp Leu Gln Val Thr Asp Ile Leu Pro Gln
530 535 540
Met Ile Thr Leu Ser Glu Ala Tyr Ile Gln Ile Trp Lys Arg Arg
545 550 555
Asp Asn Asp Glu Thr Asn Gln Gln Gly Ala
560 565

```

&lt;210&gt; 3

&lt;211&gt; 589

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1261376CD1

&lt;400&gt; 3

```

Met Ala Glu Ser Gly Glu Ser Gly Gly Pro Pro Gly Ser Gln Asp
1 5 10 15
Ser Ala Ala Gly Ala Glu Gly Ala Gly Ala Pro Ala Ala Ala Ala
20 25 30
Ser Ala Asp Ala Lys Ile Met Lys Val Thr Val Lys Thr Pro Lys
35 40 45
Glu Lys Glu Glu Phe Ala Val Pro Glu Asn Ser Ser Val Gln Gln
50 55 60
Phe Lys Glu Glu Ile Ser Lys Arg Phe Lys Ser His Thr Asp Gln

```

	65		70		75
Leu Val Leu Ile	Phe	Ala Gly Lys Ile	Leu Lys Asp Gln Asp	Thr	
	80		85		90
Leu Ser Gln His	Gly	Ile His Asp Gly	Leu Thr Val His Leu	Val	
	95		100		105
Ile Lys Thr Gln	Asn	Arg Pro Gln Asp	His Ser Ala Gln Gln	Thr	
	110		115		120
Asn Thr Ala Gly	Ser	Asn Val Thr Thr	Ser Ser Thr Pro Asn	Ser	
	125		130		135
Asn Ser Thr Ser	Gly	Ser Ala Thr Ser	Asn Pro Phe Gly Leu	Gly	
	140		145		150
Gly Leu Gly Gly	Leu	Ala Gly Leu Ser	Ser Leu Gly Leu Asn	Thr	
	155		160		165
Thr Asn Phe Ser	Glu	Leu Gln Ser Gln	Met Gln Arg Gln Leu	Leu	
	170		175		180
Ser Asn Pro Glu	Met	Met Val Gln Ile	Met Glu Asn Pro Phe	Val	
	185		190		195
Gln Ser Met Leu	Ser	Asn Pro Asp Leu	Met Arg Gln Leu Ile	Met	
	200		205		210
Ala Asn Pro Gln	Met	Gln Gln Leu Ile	Gln Arg Asn Pro Glu	Ile	
	215		220		225
Ser His Met Leu	Asn	Asn Pro Asp Ile	Met Arg Gln Thr Leu	Glu	
	230		235		240
Leu Ala Arg Asn	Pro	Ala Met Met Gln	Glu Met Met Arg Asn	Gln	
	245		250		255
Asp Arg Ala Leu	Ser	Asn Leu Glu Ser	Ile Pro Gly Gly Tyr	Asn	
	260		265		270
Ala Leu Arg Arg	Met	Tyr Thr Asp Ile	Gln Glu Pro Met Leu	Ser	
	275		280		285
Ala Ala Gln Glu	Gln	Phe Gly Gly Asn	Pro Phe Ala Ser Leu	Val	
	290		295		300
Ser Asn Thr Ser	Ser	Gly Glu Gly Ser	Gln Pro Ser Arg Thr	Glu	
	305		310		315
Asn Arg Asp Pro	Leu	Pro Asn Pro Trp	Ala Pro Gln Thr Ser	Gln	
	320		325		330
Ser Ser Ser Ala	Ser	Ser Gly Thr Ala	Ser Thr Val Gly Gly	Thr	
	335		340		345
Thr Gly Ser Thr	Ala	Ser Gly Thr Ser	Gly Gln Ser Thr Thr	Ala	
	350		355		360
Pro Asn Leu Val	Pro	Gly Val Gly Ala	Ser Met Phe Asn Thr	Pro	
	365		370		375
Gly Met Gln Ser	Leu	Leu Gln Gln Ile	Thr Glu Asn Pro Gln	Leu	
	380		385		390
Met Gln Asn Met	Leu	Ser Ala Pro Tyr	Met Arg Ser Met Met	Gln	
	395		400		405
Ser Leu Ser Gln	Asn	Pro Asp Leu Ala	Ala Gln Met Met Leu	Asn	
	410		415		420
Asn Pro Leu Phe	Ala	Gly Asn Pro Gln	Leu Gln Glu Gln Met	Arg	
	425		430		435
Gln Gln Leu Pro	Thr	Phe Leu Gln Gln	Met Gln Asn Pro Asp	Thr	
	440		445		450
Leu Ser Ala Met	Ser	Asn Pro Arg Ala	Met Gln Ala Leu Leu	Gln	
	455		460		465
Ile Gln Gln Gly	Leu	Gln Thr Leu Ala	Thr Glu Ala Pro Gly	Leu	
	470		475		480
Ile Pro Gly Phe	Thr	Pro Gly Leu Gly	Ala Leu Gly Ser Thr	Gly	
	485		490		495
Gly Ser Ser Gly	Thr	Asn Gly Ser Asn	Ala Thr Pro Ser Glu	Asn	
	500		505		510
Thr Ser Pro Thr	Ala	Gly Thr Thr Glu	Pro Gly His Gln Gln	Phe	
	515		520		525
Ile Gln Gln Met	Leu	Gln Ala Leu Ala	Gly Val Asn Pro Gln	Leu	
	530		535		540
Gln Asn Pro Glu	Val	Arg Phe Gln Gln	Gln Leu Glu Gln Leu	Ser	
	545		550		555
Ala Met Gly Phe	Leu	Asn Arg Glu Ala	Asn Leu Gln Ala Leu	Ile	
	560		565		570

Ala Thr Gly Gly Asp Ile Asn Ala Ala Ile Glu Arg Leu Leu Gly  
 575 580 585

Ser Gln Pro Ser

<210> 4  
 <211> 775  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1299481CD1

<400> 4

Met	Thr	Ile	Val	Asp	Lys	Ala	Ser	Glu	Ser	Ser	Asp	Pro	Ser	Ala	
1				5					10					15	
Tyr	Gln	Asn	Gln	Pro	Gly	Ser	Ser	Glu	Ala	Val	Ser	Pro	Gly	Asp	
				20					25					30	
Met	Asp	Ala	Gly	Ser	Ala	Ser	Trp	Gly	Ala	Val	Ser	Ser	Leu	Asn	
				35					40					45	
Asp	Val	Ser	Asn	His	Thr	Leu	Ser	Leu	Gly	Pro	Val	Pro	Gly	Ala	
				50					55					60	
Val	Val	Tyr	Ser	Ser	Ser	Ser	Val	Pro	Asp	Lys	Ser	Lys	Pro	Ser	
				65					70					75	
Pro	Gln	Lys	Asp	Gln	Ala	Leu	Gly	Asp	Gly	Ile	Ala	Pro	Pro	Gln	
				80					85					90	
Lys	Val	Leu	Phe	Pro	Ser	Glu	Lys	Ile	Cys	Leu	Lys	Trp	Gln	Gln	
				95					100					105	
Thr	His	Arg	Val	Gly	Ala	Gly	Leu	Gln	Asn	Leu	Gly	Asn	Thr	Cys	
				110					115					120	
Phe	Ala	Asn	Ala	Ala	Leu	Gln	Cys	Leu	Thr	Tyr	Thr	Pro	Pro	Leu	
				125					130					135	
Ala	Asn	Tyr	Met	Leu	Ser	His	Glu	His	Ser	Lys	Thr	Cys	His	Ala	
				140					145					150	
Glu	Gly	Phe	Cys	Met	Met	Cys	Thr	Met	Gln	Ala	His	Ile	Thr	Gln	
				155					160					165	
Ala	Leu	Ser	Asn	Pro	Gly	Asp	Val	Ile	Lys	Pro	Met	Phe	Val	Ile	
				170					175					180	
Asn	Glu	Met	Arg	Arg	Ile	Ala	Arg	His	Leu	Arg	Phe	Gly	Asn	Gln	
				185					190					195	
Glu	Asp	Ala	His	Glu	Phe	Leu	Gln	Tyr	Thr	Val	Asp	Ala	Met	Gln	
				200					205					210	
Lys	Ala	Cys	Leu	Asn	Gly	Ser	Asn	Lys	Leu	Asp	Arg	His	Thr	Gln	
				215					220					225	
Ala	Thr	Thr	Leu	Val	Cys	Gln	Ile	Phe	Gly	Gly	Tyr	Leu	Arg	Ser	
				230					235					240	
Arg	Val	Lys	Cys	Leu	Asn	Cys	Lys	Gly	Val	Ser	Asp	Thr	Phe	Asp	
				245					250					255	
Pro	Tyr	Leu	Asp	Ile	Thr	Leu	Glu	Ile	Lys	Ala	Ala	Gln	Ser	Val	
				260					265					270	
Asn	Lys	Ala	Leu	Glu	Gln	Phe	Val	Lys	Pro	Glu	Gln	Leu	Asp	Gly	
				275					280					285	
Glu	Asn	Ser	Tyr	Lys	Cys	Ser	Lys	Cys	Lys	Lys	Met	Val	Pro	Ala	
				290					295					300	
Ser	Lys	Arg	Phe	Thr	Ile	His	Arg	Ser	Ser	Asn	Val	Leu	Thr	Leu	
				305					310					315	
Ser	Leu	Lys	Arg	Phe	Ala	Asn	Phe	Thr	Gly	Gly	Lys	Ile	Ala	Lys	
				320					325					330	
Asp	Val	Lys	Tyr	Pro	Glu	Tyr	Leu	Asp	Ile	Arg	Pro	Tyr	Met	Ser	
				335					340					345	
Gln	Pro	Asn	Gly	Glu	Pro	Ile	Val	Tyr	Val	Leu	Tyr	Ala	Val	Leu	
				350					355					360	
Val	His	Thr	Gly	Phe	Asn	Cys	His	Ala	Gly	His	Tyr	Phe	Cys	Tyr	
				365					370					375	
Ile	Lys	Ala	Ser	Asn	Gly	Leu	Trp	Tyr	Gln	Met	Asn	Asp	Ser	Ile	
				380					385					390	
Val	Ser	Thr	Ser	Asp	Ile	Arg	Ser	Val	Leu	Ser	Gln	Gln	Ala	Tyr	

	395		400		405
Val Leu Phe Tyr	Ile Arg Ser His Asp	Val Lys Asn Gly Gly	Glu		
	410		415		420
Leu Thr His Pro	Thr His Ser Pro Gly	Gln Ser Ser Pro Arg	Pro		
	425		430		435
Val Ile Ser Gln	Arg Val Val Thr Asn	Lys Gln Ala Ala Pro	Gly		
	440		445		450
Phe Ile Gly Pro	Gln Leu Pro Ser His	Met Ile Lys Asn Pro	Pro		
	455		460		465
His Leu Asn Gly	Thr Gly Pro Leu Lys	Asp Thr Pro Ser Ser	Ser		
	470		475		480
Met Ser Ser Pro	Asn Gly Asn Ser Ser	Val Asn Arg Ala Ser	Pro		
	485		490		495
Val Asn Ala Ser	Ala Ser Val Gln Asn	Trp Ser Val Asn Arg	Ser		
	500		505		510
Ser Val Ile Pro	Glu His Pro Lys Lys	Gln Lys Ile Thr Ile	Ser		
	515		520		525
Ile His Asn Lys	Leu Pro Val Arg Gln	Cys Gln Ser Gln Pro	Asn		
	530		535		540
Leu His Ser Asn	Ser Leu Glu Asn Pro	Thr Lys Pro Val Pro	Ser		
	545		550		555
Ser Thr Ile Thr	Asn Ser Ala Val Gln	Ser Thr Ser Asn Ala	Ser		
	560		565		570
Thr Met Ser Val	Ser Ser Lys Val Thr	Lys Pro Ile Pro Arg	Ser		
	575		580		585
Glu Ser Cys Ser	Gln Pro Val Met Asn	Gly Lys Ser Lys Leu	Asn		
	590		595		600
Ser Ser Val Leu	Val Pro Tyr Gly Ala	Glu Ser Ser Glu Asp	Ser		
	605		610		615
Asp Glu Glu Ser	Lys Gly Leu Gly Lys	Glu Asn Gly Ile Gly	Thr		
	620		625		630
Ile Val Ser Ser	His Ser Pro Gly Gln	Asp Ala Glu Asp Glu	Glu		
	635		640		645
Ala Thr Pro His	Glu Leu Gln Glu Pro	Met Thr Leu Asn Gly	Ala		
	650		655		660
Asn Ser Ala Asp	Ser Asp Ser Asp Pro	Lys Glu Asn Gly Leu	Ala		
	665		670		675
Pro Asp Gly Ala	Ser Cys Gln Gly Gln	Pro Ala Leu His Ser	Glu		
	680		685		690
Asn Pro Phe Ala	Lys Ala Asn Gly Leu	Pro Gly Lys Leu Met	Pro		
	695		700		705
Ala Pro Leu Leu	Ser Leu Pro Glu Asp	Lys Ile Leu Glu Thr	Phe		
	710		715		720
Arg Leu Ser Asn	Lys Leu Lys Gly Ser	Thr Asp Glu Met Ser	Ala		
	725		730		735
Pro Gly Ala Glu	Arg Gly Pro Pro Glu	Asp Arg Asp Ala Glu	Pro		
	740		745		750
Gln Pro Gly Ser	Pro Ala Ala Glu Ser	Leu Glu Glu Pro Asp	Ala		
	755		760		765
Ala Ala Ser Leu	Phe Pro Phe Ser Glu	Gly			
	770		775		

&lt;210&gt; 5

&lt;211&gt; 351

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1873139CD1

&lt;400&gt; 5

Met Asn Ala Ile Leu	Gln Ser Leu Ser	Asn Ile Glu Gln Phe	Cys
1	5	10	15
Cys Tyr Phe Lys Glu	Leu Pro Ala Val	Glu Leu Arg Asn Gly	Lys
20	25	30	
Thr Ala Gly Arg Arg	Thr Tyr His Thr	Arg Ser Gln Gly Asp	Asn
35	40	45	



```

Asn Val Ser Leu Val Glu Glu Phe Arg Lys Thr Leu Cys Ala Leu
      50      55
Trp Gln Gly Ser Gln Thr Ala Phe Ser Pro Glu Ser Leu Phe Tyr
      65      70
Val Val Trp Lys Ile Met Pro Asn Phe Arg Gly Tyr Gln Gln Gln
      80      85
Asp Ala His Glu Phe Met Arg Tyr Leu Leu Asp His Leu His Leu
      95     100
Glu Leu Gln Gly Gly Phe Asn Gly Val Ser Arg Ser Ala Ile Leu
     110     115
Gln Glu Asn Ser Thr Leu Ser Ala Ser Asn Lys Cys Cys Ile Asn
     125     130
Gly Ala Ser Thr Val Val Thr Ala Ile Phe Gly Gly Ile Leu Gln
     140     145
Asn Glu Val Asn Cys Leu Ile Cys Gly Thr Glu Ser Arg Lys Phe
     155     160
Asp Pro Phe Leu Asp Leu Ser Leu Asp Ile Pro Ser Gln Phe Arg
     170     175
Ser Lys Arg Ser Lys Asn Gln Glu Asn Gly Pro Val Cys Ser Leu
     185     190
Arg Asp Cys Leu Arg Ser Phe Thr Asp Leu Glu Glu Leu Asp Glu
     200     205
Thr Glu Leu Tyr Met Cys His Lys Cys Lys Lys Lys Gln Lys Ser
     215     220
Thr Lys Lys Phe Trp Ile Gln Lys Leu Pro Lys Val Leu Cys Leu
     230     235
His Leu Lys Arg Phe His Trp Thr Ala Tyr Leu Arg Asn Lys Val
     245     250
Asp Thr Tyr Val Glu Phe Pro Leu Arg Gly Leu Asp Met Lys Cys
     260     265
Tyr Leu Leu Glu Pro Glu Asn Ser Gly Pro Glu Ser Cys Leu Tyr
     275     280
Asp Leu Ala Ala Val Val Val His His Gly Ser Gly Val Gly Ser
     290     295
Gly His Tyr Thr Ala Tyr Ala Thr His Glu Gly Arg Trp Phe His
     305     310
Phe Asn Asp Ser Thr Val Thr Leu Thr Asp Glu Glu Thr Val Val
     320     325
Lys Ala Lys Ala Tyr Ile Leu Phe Tyr Val Glu His Gln Ala Lys
     335     340
Ala Gly Ser Asp Lys Leu
     350

```

<210> 6  
 <211> 136  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1903112CD1

```

<400> 6
Met Ala Leu Met Gln Arg Ser Asp Ile Phe Arg Val Ala Ile Ala
  1      5      10
Gly Ala Pro Val Thr Leu Trp Ile Phe Tyr Asp Thr Gly Tyr Thr
      20      25
Glu Arg Tyr Met Gly His Pro Asp Gln Asn Glu Gln Gly Tyr Tyr
      35      40
Leu Gly Ser Val Ala Met Gln Ala Glu Lys Phe Pro Ser Glu Pro
      50      55
Asn Arg Leu Leu Leu Leu His Gly Phe Leu Asp Glu Asn Val His
      65      70
Phe Ala His Thr Ser Ile Leu Leu Ser Phe Leu Val Arg Ala Gly
      80      85
Lys Pro Tyr Asp Leu Gln Ile Tyr Pro Gln Glu Arg His Ser Ile
      95     100
Arg Val Pro Glu Ser Gly Glu His Tyr Glu Leu His Leu Leu His

```

				110						115					120
Tyr	Leu	Gln	Glu	Asn	Leu	Gly	Ser	Arg	Ile	Ala	Ala	Leu	Lys	Val	
				125					130					135	
Ile															

<210> 7  
 <211> 396  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1993044CD1

<400> 7

Met	Ser	Leu	Gly	Trp	Leu	Glu	Arg	Pro	Pro	Ala	Leu	Ser	Arg	Ala	
1				5					10					15	
Ala	Gly	Asp	Gly	Ala	Arg	Arg	Leu	Ser	Gly	Ser	Met	Arg	Gly	Asp	
				20					25					30	
Val	Trp	Leu	Thr	Ser	Ser	Ala	Ala	Gly	Leu	Leu	Arg	Ser	Val	Ala	
				35					40					45	
Gly	Gly	Ser	Trp	Cys	Gly	Gly	Gln	Leu	Arg	Ala	Arg	Gly	Gly	Ser	
				50					55					60	
Gly	Arg	Cys	Val	Ala	Arg	Ala	Met	Thr	Gly	Asn	Ala	Gly	Glu	Trp	
				65					70					75	
Cys	Leu	Met	Glu	Ser	Asp	Pro	Gly	Val	Phe	Thr	Glu	Leu	Ile	Lys	
				80					85					90	
Gly	Phe	Gly	Cys	Arg	Gly	Ala	Gln	Val	Glu	Glu	Ile	Trp	Ser	Leu	
				95					100					105	
Glu	Pro	Glu	Asn	Phe	Glu	Lys	Leu	Lys	Pro	Val	His	Gly	Leu	Ile	
				110					115					120	
Phe	Leu	Phe	Lys	Trp	Gln	Pro	Gly	Glu	Glu	Pro	Ala	Gly	Ser	Val	
				125					130					135	
Val	Gln	Asp	Ser	Arg	Leu	Asp	Thr	Ile	Phe	Phe	Ala	Lys	Gln	Val	
				140					145					150	
Ile	Asn	Asn	Ala	Cys	Ala	Thr	Gln	Ala	Ile	Val	Ser	Val	Leu	Leu	
				155					160					165	
Asn	Cys	Thr	His	Gln	Asp	Val	His	Leu	Gly	Glu	Thr	Leu	Ser	Glu	
				170					175					180	
Phe	Lys	Glu	Phe	Ser	Gln	Ser	Phe	Asp	Ala	Ala	Met	Lys	Gly	Leu	
				185					190					195	
Ala	Leu	Ser	Asn	Ser	Asp	Val	Ile	Arg	Gln	Val	His	Asn	Ser	Phe	
				200					205					210	
Ala	Arg	Gln	Gln	Met	Phe	Glu	Phe	Asp	Thr	Lys	Thr	Ser	Ala	Lys	
				215					220					225	
Glu	Glu	Asp	Ala	Phe	His	Phe	Val	Ser	Tyr	Val	Pro	Val	Asn	Gly	
				230					235					240	
Arg	Leu	Tyr	Glu	Leu	Asp	Gly	Leu	Arg	Glu	Gly	Pro	Ile	Asp	Leu	
				245					250					255	
Gly	Ala	Cys	Asn	Gln	Asp	Asp	Trp	Phe	Ser	Ala	Val	Arg	Pro	Val	
				260					265					270	
Ile	Glu	Lys	Arg	Ile	Gln	Lys	Tyr	Ser	Glu	Gly	Glu	Ile	Arg	Phe	
				275					280					285	
Asn	Leu	Met	Ala	Ile	Val	Ser	Asp	Arg	Lys	Met	Ile	Tyr	Glu	Gln	
				290					295					300	
Lys	Ile	Ala	Glu	Leu	Gln	Arg	Gln	Leu	Ala	Glu	Glu	Glu	Pro	Met	
				305					310					315	
Asp	Thr	Asp	Gln	Gly	Asn	Ser	Met	Leu	Ser	Ala	Ile	Gln	Ser	Glu	
				320					325					330	
Val	Ala	Lys	Asn	Gln	Met	Leu	Ile	Glu	Glu	Glu	Val	Gln	Lys	Leu	
				335					340					345	
Lys	Arg	Tyr	Lys	Ile	Glu	Asn	Ile	Arg	Arg	Lys	His	Asn	Tyr	Leu	
				350					355					360	
Pro	Phe	Ile	Met	Glu	Leu	Leu	Lys	Thr	Leu	Ala	Glu	His	Gln	Gln	
				365					370					375	
Leu	Ile	Pro	Leu	Val	Glu	Lys	Ala	Lys	Glu	Lys	Gln	Asn	Ala	Lys	
				380					385					390	

Lys Ala Gln Glu Thr Lys  
395

<210> 8  
<211> 246  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2292182CD1

<400> 8  
Met Ala Gly Ala Pro Asp Glu Arg Arg Arg Gly Pro Ala Ala Gly  
1 5 10 15  
Glu Gln Leu Gln Gln Gln His Val Ser Cys Gln Val Phe Pro Glu  
20 25 30  
Arg Leu Ala Gln Gly Asn Pro Gln Gln Gly Phe Phe Ser Ser Phe  
35 40 45  
Phe Thr Ser Asn Gln Lys Cys Gln Leu Arg Leu Leu Lys Thr Leu  
50 55 60  
Glu Thr Asn Pro Tyr Val Lys Leu Leu Leu Asp Ala Met Lys His  
65 70 75  
Ser Gly Cys Ala Val Asn Lys Asp Arg His Phe Ser Cys Glu Asp  
80 85 90  
Cys Asn Gly Asn Val Ser Gly Gly Phe Asp Ala Ser Thr Ser Gln  
95 100 105  
Ile Val Leu Cys Gln Asn Asn Ile His Asn Gln Ala His Met Asn  
110 115 120  
Arg Val Val Thr His Glu Leu Ile His Ala Phe Asp His Cys Arg  
125 130 135  
Ala His Val Asp Trp Phe Thr Asn Ile Arg His Leu Ala Cys Ser  
140 145 150  
Glu Val Arg Ala Ala Asn Leu Ser Gly Asp Cys Ser Leu Val Asn  
155 160 165  
Glu Ile Phe Arg Leu His Phe Gly Leu Lys Gln His His Gln Thr  
170 175 180  
Cys Val Arg Asp Arg Ala Thr Leu Ser Ile Leu Ala Val Arg Asn  
185 190 195  
Ile Ser Lys Glu Val Ala Lys Lys Ala Val Asp Glu Val Phe Glu  
200 205 210  
Ser Cys Phe Asn Asp His Glu Pro Phe Gly Arg Ile Pro His Asn  
215 220 225  
Lys Thr Tyr Ala Arg Tyr Ala His Arg Asp Phe Glu Asn Arg Asp  
230 235 240  
Arg Tyr Tyr Ser Asn Ile  
245

<210> 9  
<211> 262  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2331301CD1

<400> 9  
Met Glu Val Tyr Ile Arg His Leu Glu Lys Val Leu Arg Arg Tyr  
1 5 10 15  
Val Gln Arg Leu Gln Trp Leu Leu Ser Gly Ser Arg Arg Leu Phe  
20 25 30  
Gly Thr Val Leu Glu Ser Lys Val Cys Ile Leu Leu Asp Thr Ser  
35 40 45  
Gly Ser Met Gly Pro Tyr Leu Gln Gln Val Lys Thr Glu Leu Val  
50 55 60  
Leu Leu Ile Trp Glu Gln Leu Arg Lys Cys Cys Asp Ser Phe Asn  
65 70 75  
Leu Leu Ser Phe Ala Glu Ser Leu Gln Ser Trp Gln Asp Thr Leu

Val	Glu	Thr	Thr	Asp	Ala	Ala	Cys	His	Glu	Ala	Met	Gln	Trp	Val	80	85	90
				95					100					105			
Thr	His	Leu	Gln	Ala	Gln	Gly	Ser	Thr	Ser	Ile	Leu	Gln	Ala	Leu	110	115	120
Leu	Lys	Ala	Phe	Ser	Phe	His	Asp	Leu	Glu	Gly	Leu	Tyr	Leu	Leu	125	130	135
Thr	Asp	Gly	Lys	Pro	Asp	Thr	Ser	Cys	Ser	Leu	Val	Leu	Asn	Glu	140	145	150
Val	Gln	Lys	Leu	Arg	Glu	Lys	Arg	Asp	Val	Lys	Val	His	Thr	Ile	155	160	165
Ser	Leu	Asn	Cys	Ser	Asp	Arg	Ala	Ala	Val	Glu	Phe	Leu	Arg	Lys	170	175	180
Leu	Ala	Ser	Phe	Thr	Gly	Gly	Arg	Tyr	His	Cys	Pro	Val	Gly	Glu	185	190	195
Asp	Thr	Leu	Ser	Lys	Ile	His	Ser	Leu	Leu	Thr	Lys	Gly	Phe	Ile	200	205	210
Asn	Glu	Lys	Asp	Arg	Thr	Leu	Pro	Pro	Phe	Glu	Gly	Asp	Asp	Leu	215	220	225
Arg	Ile	Leu	Ala	Gln	Glu	Ile	Thr	Lys	Ala	Arg	Ser	Phe	Leu	Trp	230	235	240
Gln	Ala	Gln	Ser	Phe	Arg	Ser	Gln	Leu	Gln	Lys	Lys	Asn	Asp	Ala	245	250	255
Glu	Pro	Lys	Val	Thr	Leu	Ser									260		

&lt;210&gt; 10

&lt;211&gt; 406

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2517512CD1

&lt;400&gt; 10

Met	Ala	Ala	Ala	Val	Arg	Gln	Asp	Leu	Ala	Gln	Leu	Met	Asn	Ser	1	5	10	15
Ser	Gly	Ser	His	Lys	Asp	Leu	Ala	Gly	Lys	Tyr	Arg	Gln	Ile	Leu	20	25	30	35
Glu	Lys	Ala	Ile	Gln	Leu	Ser	Gly	Ala	Glu	Gln	Leu	Glu	Ala	Leu	35	40	45	50
Lys	Ala	Phe	Val	Glu	Ala	Met	Val	Asn	Glu	Asn	Val	Ser	Leu	Val	50	55	60	65
Ile	Ser	Arg	Gln	Leu	Leu	Thr	Asp	Phe	Cys	Thr	His	Leu	Pro	Asn	65	70	75	80
Leu	Pro	Asp	Ser	Thr	Ala	Lys	Glu	Ile	Tyr	His	Phe	Thr	Leu	Glu	80	85	90	95
Lys	Ile	Gln	Pro	Arg	Val	Ile	Ser	Phe	Glu	Glu	Gln	Val	Ala	Ser	95	100	105	110
Ile	Arg	Gln	His	Leu	Ala	Ser	Ile	Tyr	Glu	Lys	Glu	Glu	Asp	Trp	110	115	120	125
Arg	Asn	Ala	Ala	Gln	Val	Leu	Val	Gly	Ile	Pro	Leu	Glu	Thr	Gly	125	130	135	140
Gln	Lys	Gln	Tyr	Asn	Val	Asp	Tyr	Lys	Leu	Glu	Thr	Tyr	Leu	Lys	140	145	150	155
Ile	Ala	Arg	Leu	Tyr	Leu	Glu	Asp	Asp	Asp	Pro	Val	Gln	Ala	Glu	155	160	165	170
Ala	Tyr	Ile	Asn	Arg	Ala	Ser	Leu	Leu	Gln	Asn	Glu	Ser	Thr	Asn	170	175	180	185
Glu	Gln	Leu	Gln	Ile	His	Tyr	Lys	Val	Cys	Tyr	Ala	Arg	Val	Leu	185	190	195	200
Asp	Tyr	Arg	Arg	Lys	Phe	Ile	Glu	Ala	Ala	Gln	Arg	Tyr	Asn	Glu	200	205	210	215
Leu	Ser	Tyr	Lys	Thr	Ile	Val	His	Glu	Ser	Glu	Arg	Leu	Glu	Ala	215	220	225	230
Leu	Lys	His	Ala	Leu	His	Cys	Thr	Ile	Leu	Ala	Ser	Ala	Gly	Gln	230	235	240	

Gln	Arg	Ser	Arg	Met	Leu	Ala	Thr	Leu	Phe	Lys	Asp	Glu	Arg	Cys	
				245					250					255	
Gln	Gln	Leu	Ala	Ala	Tyr	Gly	Ile	Leu	Glu	Lys	Met	Tyr	Leu	Asp	
				260					265					270	
Arg	Ile	Ile	Arg	Gly	Asn	Gln	Leu	Gln	Glu	Phe	Ala	Ala	Met	Leu	
				275					280					285	
Met	Pro	His	Gln	Lys	Ala	Thr	Thr	Ala	Asp	Gly	Ser	Ser	Ile	Leu	
				290					295					300	
Asp	Arg	Ala	Val	Ile	Glu	His	Asn	Leu	Leu	Ser	Ala	Ser	Lys	Leu	
				305					310					315	
Tyr	Asn	Asn	Ile	Thr	Phe	Glu	Glu	Leu	Gly	Ala	Leu	Leu	Glu	Ile	
				320					325					330	
Pro	Ala	Ala	Lys	Ala	Glu	Lys	Ile	Ala	Ser	Gln	Met	Ile	Thr	Glu	
				335					340					345	
Gly	Arg	Met	Asn	Gly	Phe	Ile	Asp	Gln	Ile	Asp	Gly	Ile	Val	His	
				350					355					360	
Phe	Glu	Thr	Arg	Glu	Ala	Leu	Pro	Thr	Trp	Asp	Lys	Gln	Ile	Gln	
				365					370					375	
Ser	Leu	Cys	Phe	Gln	Val	Asn	Asn	Leu	Leu	Glu	Lys	Ile	Ser	Gln	
				380					385					390	
Thr	Ala	Pro	Glu	Trp	Thr	Ala	Gln	Ala	Met	Glu	Ala	Gln	Met	Ala	
				395					400					405	

Gln

&lt;210&gt; 11

&lt;211&gt; 172

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3489039CD1

&lt;400&gt; 11

Met	Leu	Leu	Pro	Asn	Ile	Leu	Leu	Thr	Gly	Thr	Pro	Gly	Val	Gly	
1				5					10					15	
Lys	Thr	Thr	Leu	Gly	Lys	Glu	Leu	Ala	Ser	Lys	Ser	Gly	Leu	Lys	
				20					25					30	
Tyr	Ile	Asn	Val	Gly	Asp	Leu	Ala	Arg	Glu	Glu	Gln	Leu	Tyr	Asp	
				35					40					45	
Gly	Tyr	Asp	Glu	Glu	Tyr	Asp	Cys	Pro	Ile	Leu	Asp	Glu	Asp	Arg	
				50					55					60	
Val	Val	Asp	Glu	Leu	Asp	Asn	Gln	Met	Arg	Glu	Gly	Gly	Val	Ile	
				65					70					75	
Val	Asp	Tyr	His	Gly	Cys	Asp	Phe	Phe	Pro	Glu	Arg	Trp	Phe	His	
				80					85					90	
Ile	Val	Phe	Val	Leu	Arg	Thr	Asp	Thr	Asn	Val	Leu	Tyr	Glu	Arg	
				95					100					105	
Leu	Glu	Thr	Arg	Gly	Tyr	Asn	Glu	Lys	Lys	Leu	Thr	Asp	Asn	Ile	
				110					115					120	
Gln	Cys	Glu	Ile	Phe	Gln	Val	Leu	Tyr	Glu	Glu	Ala	Thr	Ala	Ser	
				125					130					135	
Tyr	Lys	Glu	Glu	Ile	Val	His	Gln	Leu	Pro	Ser	Asn	Lys	Pro	Glu	
				140					145					150	
Glu	Leu	Glu	Asn	Asn	Val	Asp	Gln	Ile	Leu	Lys	Trp	Ile	Glu	Gln	
				155					160					165	
Trp	Ile	Lys	Asp	His	Asn	Ser									
				170											

&lt;210&gt; 12

&lt;211&gt; 517

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5432879CD1

&lt;400&gt; 12

Met	Leu	Ser	Ser	Arg	Ala	Glu	Ala	Ala	Met	Thr	Ala	Ala	Asp	Arg
1				5					10					15
Ala	Ile	Gln	Arg	Phe	Leu	Arg	Thr	Gly	Ala	Ala	Val	Arg	Tyr	Lys
				20					25					30
Val	Met	Lys	Asn	Trp	Gly	Val	Ile	Gly	Gly	Ile	Ala	Ala	Ala	Leu
				35					40					45
Ala	Ala	Gly	Ile	Tyr	Val	Ile	Trp	Gly	Pro	Ile	Thr	Glu	Arg	Lys
				50					55					60
Lys	Arg	Arg	Lys	Gly	Leu	Val	Pro	Gly	Leu	Val	Asn	Leu	Gly	Asn
				65					70					75
Thr	Cys	Phe	Met	Asn	Ser	Leu	Leu	Gln	Gly	Leu	Ser	Ala	Cys	Pro
				80					85					90
Ala	Phe	Ile	Arg	Trp	Leu	Glu	Glu	Phe	Thr	Ser	Gln	Tyr	Ser	Arg
				95					100					105
Asp	Gln	Lys	Glu	Pro	Pro	Ser	His	Gln	Tyr	Leu	Ser	Leu	Thr	Leu
				110					115					120
Leu	His	Leu	Leu	Lys	Ala	Leu	Ser	Cys	Gln	Glu	Val	Thr	Asp	Asp
				125					130					135
Glu	Val	Leu	Asp	Ala	Ser	Cys	Leu	Leu	Asp	Val	Leu	Arg	Met	Tyr
				140					145					150
Arg	Trp	Gln	Ile	Ser	Ser	Phe	Glu	Glu	Gln	Asp	Ala	His	Glu	Leu
				155					160					165
Phe	His	Val	Ile	Thr	Ser	Ser	Leu	Glu	Asp	Glu	Arg	Asp	Arg	Gln
				170					175					180
Pro	Arg	Val	Thr	His	Leu	Phe	Asp	Val	His	Ser	Leu	Glu	Gln	Gln
				185					190					195
Ser	Glu	Ile	Thr	Pro	Lys	Gln	Ile	Thr	Cys	Arg	Thr	Arg	Gly	Ser
				200					205					210
Pro	His	Pro	Thr	Ser	Asn	His	Trp	Lys	Ser	Gln	His	Pro	Phe	His
				215					220					225
Gly	Arg	Leu	Thr	Ser	Asn	Met	Val	Cys	Lys	His	Cys	Glu	His	Gln
				230					235					240
Ser	Pro	Val	Arg	Phe	Asp	Thr	Phe	Asp	Ser	Leu	Ser	Leu	Ser	Ile
				245					250					255
Pro	Ala	Ala	Thr	Trp	Gly	His	Pro	Leu	Thr	Leu	Asp	His	Cys	Leu
				260					265					270
His	His	Phe	Ile	Ser	Ser	Glu	Ser	Val	Arg	Asp	Val	Val	Cys	Asp
				275					280					285
Asn	Cys	Thr	Lys	Ile	Glu	Ala	Lys	Gly	Thr	Leu	Asn	Gly	Glu	Lys
				290					295					300
Val	Glu	His	Gln	Arg	Thr	Thr	Phe	Val	Lys	Gln	Leu	Lys	Leu	Gly
				305					310					315
Lys	Leu	Pro	Gln	Cys	Leu	Cys	Ile	His	Leu	Gln	Arg	Leu	Ser	Trp
				320					325					330
Ser	Ser	His	Gly	Thr	Pro	Leu	Lys	Arg	His	Glu	His	Val	Gln	Phe
				335					340					345
Asn	Glu	Phe	Leu	Met	Met	Asp	Ile	Tyr	Lys	Tyr	His	Leu	Leu	Gly
				350					355					360
His	Lys	Pro	Ser	Gln	His	Asn	Pro	Lys	Leu	Asn	Lys	Asn	Pro	Gly
				365					370					375
Pro	Thr	Leu	Glu	Leu	Gln	Asp	Gly	Pro	Gly	Ala	Pro	Thr	Pro	Val
				380					385					390
Leu	Asn	Gln	Pro	Gly	Ala	Pro	Lys	Thr	Gln	Ile	Phe	Met	Asn	Gly
				395					400					405
Ala	Cys	Ser	Pro	Ser	Leu	Leu	Pro	Thr	Leu	Ser	Ala	Pro	Met	Pro
				410					415					420
Phe	Pro	Leu	Pro	Val	Val	Pro	Asp	Tyr	Ser	Ser	Ser	Thr	Tyr	Leu
				425					430					435
Phe	Arg	Leu	Met	Ala	Val	Val	Val	His	His	Gly	Asp	Met	His	Ser
				440					445					450
Gly	His	Phe	Val	Thr	Tyr	Arg	Arg	Ser	Pro	Pro	Ser	Ala	Arg	Asn
				455					460					465
Pro	Leu	Ser	Thr	Ser	Asn	Gln	Trp	Leu	Trp	Val	Ser	Asp	Asp	Thr
				470					475					480
Val	Arg	Lys	Ala	Ser	Leu	Gln	Glu	Val	Leu	Ser	Ser	Ser	Ala	Tyr
				485					490					495

Leu Leu Phe Tyr Glu Arg Val Leu Ser Arg Met Gln His Gln Ser  
 500 505 510  
 Gln Glu Cys Lys Ser Glu Glu  
 515

<210> 13  
 <211> 346  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5853753CD1

<400> 13  
 Met Val Glu Lys Glu Glu Ala Gly Gly Gly Ile Ser Glu Glu Glu  
 1 5 10 15  
 Ala Ala Gln Tyr Asp Arg Gln Ile Arg Leu Trp Gly Leu Glu Ala  
 20 25 30  
 Gln Lys Arg Leu Arg Ala Ser Arg Val Leu Leu Val Gly Leu Lys  
 35 40 45  
 Gly Leu Gly Ala Glu Ile Ala Lys Asn Leu Ile Leu Ala Gly Val  
 50 55 60  
 Lys Gly Leu Thr Met Leu Asp His Glu Gln Val Thr Pro Glu Asp  
 65 70 75  
 Pro Gly Ala Gln Phe Leu Ile Arg Thr Gly Ser Val Gly Arg Asn  
 80 85 90  
 Arg Ala Glu Ala Ser Leu Glu Arg Ala Gln Asn Leu Asn Pro Met  
 95 100 105  
 Val Asp Val Lys Val Asp Thr Glu Asp Ile Glu Lys Lys Pro Glu  
 110 115 120  
 Ser Phe Phe Thr Gln Phe Asp Ala Val Cys Leu Thr Cys Cys Ser  
 125 130 135  
 Arg Asp Val Ile Val Lys Val Asp Gln Ile Cys His Lys Asn Ser  
 140 145 150  
 Ile Lys Phe Phe Thr Gly Asp Val Phe Gly Tyr His Gly Tyr Thr  
 155 160 165  
 Phe Ala Asn Leu Gly Glu His Glu Phe Val Glu Glu Lys Thr Lys  
 170 175 180  
 Val Ala Lys Val Ser Gln Gly Val Glu Asp Gly Pro Asp Thr Lys  
 185 190 195  
 Arg Ala Lys Leu Asp Ser Ser Glu Thr Thr Met Val Lys Lys Lys  
 200 205 210  
 Val Val Phe Cys Pro Val Lys Glu Ala Leu Glu Val Asp Trp Ser  
 215 220 225  
 Ser Glu Lys Ala Lys Ala Ala Leu Lys Arg Thr Thr Ser Asp Tyr  
 230 235 240  
 Phe Leu Leu Gln Val Leu Leu Lys Phe Arg Thr Asp Lys Gly Arg  
 245 250 255  
 Asp Pro Ser Ser Asp Thr Tyr Glu Glu Asp Ser Glu Leu Leu Leu  
 260 265 270  
 Gln Ile Arg Asn Asp Val Leu Asp Ser Leu Gly Ile Ser Pro Asp  
 275 280 285  
 Leu Leu Pro Glu Asp Phe Val Arg Tyr Cys Phe Ser Glu Met Ala  
 290 295 300  
 Pro Val Cys Ala Val Val Gly Gly Ile Leu Ala Gln Glu Ile Val  
 305 310 315  
 Lys Ala Leu Ser Gln Arg Asp Pro Pro His Asn Asn Phe Phe Phe  
 320 325 330  
 Phe Asp Gly Met Lys Gly Asn Gly Ile Val Glu Cys Leu Gly Pro  
 335 340 345  
 Lys

<210> 14  
 <211> 151  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 411344CD1

<400> 14  
 Met Ala Ser Met Gln Lys Arg Leu Gln Lys Glu Leu Leu Ala Leu  
 1 5 10 15  
 Gln Asn Asp Pro Pro Gly Met Thr Leu Asn Glu Lys Ser Val  
 20 25 30  
 Gln Asn Ser Ile Thr Gln Trp Ile Val Asp Met Glu Gly Ala Pro  
 35 40 45  
 Gly Thr Leu Tyr Glu Gly Glu Lys Phe Gln Leu Leu Phe Lys Phe  
 50 55 60  
 Ser Ser Arg Tyr Pro Phe Asp Ser Pro Gln Val Met Phe Thr Gly  
 65 70 75  
 Glu Asn Ile Pro Val His Pro His Val Tyr Ser Asn Gly His Ile  
 80 85 90  
 Cys Leu Ser Ile Leu Thr Glu Asp Trp Ser Pro Ala Leu Ser Val  
 95 100 105  
 Gln Ser Val Cys Leu Ser Ile Ile Ser Met Leu Ser Ser Cys Lys  
 110 115 120  
 Glu Lys Arg Arg Pro Pro Asp Asn Ser Phe Tyr Val Arg Thr Cys  
 125 130 135  
 Asn Lys Asn Pro Lys Lys Thr Lys Trp Trp Tyr His Asp Asp Thr  
 140 145 150  
 Cys

<210> 15  
 <211> 362  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1256390CD1

<400> 15  
 Met Leu Val Pro Gly Gly Leu Gly Tyr Asp Arg Ser Leu Ala Gln  
 1 5 10 15  
 His Arg Gln Glu Ile Val Asp Lys Ser Val Ser Pro Trp Ser Leu  
 20 25 30  
 Glu Thr Tyr Ser Tyr Asn Ile Tyr His Pro Met Gly Glu Ile Tyr  
 35 40 45  
 Glu Trp Met Arg Glu Ile Ser Glu Lys Tyr Lys Glu Val Val Thr  
 50 55 60  
 Gln His Phe Leu Gly Val Thr Tyr Glu Thr His Pro Met Tyr Tyr  
 65 70 75  
 Leu Lys Ile Ser Gln Pro Ser Gly Asn Pro Lys Lys Ile Ile Trp  
 80 85 90  
 Met Asp Cys Gly Ile His Ala Arg Glu Trp Ile Ala Pro Ala Phe  
 95 100 105  
 Cys Gln Trp Phe Val Lys Glu Ile Leu Gln Asn His Lys Asp Asn  
 110 115 120  
 Ser Ser Ile Arg Lys Leu Leu Arg Asn Leu Asp Phe Tyr Val Leu  
 125 130 135  
 Pro Val Leu Asn Ile Asp Gly Tyr Ile Tyr Thr Trp Thr Thr Asp  
 140 145 150  
 Arg Leu Trp Arg Lys Ser Arg Ser Pro His Asn Asn Gly Thr Cys  
 155 160 165  
 Phe Gly Thr Asp Leu Asn Arg Asn Phe Asn Ala Ser Trp Cys Ser  
 170 175 180  
 Ile Gly Ala Ser Arg Asn Cys Gln Asp Gln Thr Phe Cys Gly Thr  
 185 190 195  
 Gly Pro Val Ser Glu Pro Glu Thr Lys Ala Val Ala Ser Phe Ile  
 200 205 210  
 Glu Ser Lys Lys Asp Asp Ile Leu Cys Phe Leu Thr Met His Ser  
 215 220 225



Tyr	Gly	Gln	Leu	Ile	Leu	Thr	Pro	Tyr	Gly	Tyr	Thr	Lys	Asn	Lys
				230					235					240
Ser	Ser	Asn	His	Pro	Glu	Met	Ile	Gln	Val	Gly	Gln	Lys	Ala	Ala
				245					250					255
Asn	Ala	Leu	Lys	Ala	Lys	Tyr	Gly	Thr	Asn	Tyr	Arg	Val	Gly	Ser
				260					265					270
Ser	Ala	Asp	Ile	Leu	Tyr	Ala	Ser	Ser	Gly	Ser	Ser	Arg	Asp	Trp
				275					280					285
Ala	Arg	Asp	Ile	Gly	Ile	Pro	Phe	Ser	Tyr	Thr	Phe	Glu	Leu	Arg
				290					295					300
Asp	Ser	Gly	Thr	Tyr	Gly	Phe	Val	Leu	Pro	Glu	Ala	Gln	Ile	Gln
				305					310					315
Pro	Thr	Cys	Glu	Glu	Thr	Met	Glu	Ala	Val	Leu	Ser	Val	Leu	Asp
				320					325					330
Asp	Val	Tyr	Ala	Lys	His	Trp	His	Ser	Asp	Ser	Ala	Gly	Arg	Val
				335					340					345
Thr	Ser	Ala	Thr	Met	Leu	Leu	Gly	Leu	Leu	Val	Ser	Cys	Met	Ser
				350					355					360

Leu Leu

&lt;210&gt; 16

&lt;211&gt; 123

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1786774CD1

&lt;400&gt; 16

Met	Ser	Gly	Glu	Glu	Leu	Ser	Glu	Ser	Thr	Pro	Glu	Pro	Gln	Lys
1				5					10					15
Glu	Ile	Ser	Glu	Ser	Leu	Ser	Val	Thr	Arg	Asp	Gln	Asp	Glu	Asp
				20					25					30
Asp	Lys	Ala	Pro	Glu	Pro	Thr	Trp	Ala	Asp	Asp	Leu	Pro	Ala	Thr
				35					40					45
Thr	Ser	Ser	Glu	Ala	Thr	Thr	Thr	Pro	Arg	Pro	Leu	Leu	Ser	Thr
				50					55					60
Pro	Val	Asp	Gly	Ala	Glu	Asp	Pro	Arg	Cys	Leu	Glu	Ala	Leu	Lys
				65					70					75
Pro	Gly	Asn	Cys	Gly	Glu	Tyr	Val	Val	Arg	Trp	Tyr	Tyr	Asp	Lys
				80					85					90
Gln	Val	Asn	Ser	Cys	Ala	Arg	Phe	Trp	Phe	Ser	Gly	Cys	Asn	Gly
				95					100					105
Ser	Gly	Asn	Arg	Phe	Asn	Ser	Glu	Lys	Glu	Cys	Gln	Glu	Thr	Cys
				110					115					120

Ile Gln Gly

&lt;210&gt; 17

&lt;211&gt; 983

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1911808CD1

&lt;400&gt; 17

Met	Ala	Pro	Arg	Leu	Gln	Leu	Glu	Lys	Ala	Ala	Trp	Arg	Trp	Ala
1				5					10					15
Glu	Thr	Val	Arg	Pro	Glu	Glu	Val	Ser	Gln	Glu	His	Ile	Glu	Thr
				20					25					30
Ala	Tyr	Arg	Ile	Trp	Leu	Glu	Pro	Cys	Ile	Arg	Gly	Val	Cys	Arg
				35					40					45
Arg	Asn	Cys	Lys	Gly	Asn	Pro	Asn	Cys	Leu	Val	Gly	Ile	Gly	Glu
				50					55					60
His	Ile	Trp	Leu	Gly	Glu	Ile	Asp	Glu	Asn	Ser	Phe	His	Asn	Ile

Asp Asp Pro Asn Cys	65	Glu Arg Arg Lys	70	Lys Asn Ser Phe Val	75
Leu Thr Asn Leu Gly	80	Ala Thr Cys Tyr	85	Val Asn Thr Phe Leu	90
Val Trp Phe Leu Asn	95	Leu Glu Leu Arg	100	Gln Ala Leu Tyr Leu	105
Pro Ser Thr Cys Ser	110	Asp Tyr Met Leu	115	Gly Asp Gly Ile Gln	120
Glu Lys Asp Tyr Glu	125	Pro Gln Thr Ile	130	Cys Glu His Leu Gln	135
Leu Phe Ala Leu Leu	140	Gln Asn Ser Asn	145	Arg Arg Tyr Ile Asp	150
Ser Gly Phe Val Lys	155	Ala Leu Gly Leu	160	Asp Thr Gly Gln Gln	165
Asp Ala Gln Glu Phe	170	Ser Lys Leu Phe	175	Met Ser Leu Leu Glu	180
Thr Leu Ser Asn Gln	185	Lys Asn Pro Asp	190	Val Arg Asn Ile Val	195
Gln Gln Phe Cys Gly	200	Glu Tyr Ala Tyr	205	Val Thr Val Cys Asn	210
Cys Gly Arg Glu Ser	215	Lys Leu Leu Ser	220	Phe Tyr Glu Leu Glu	225
Leu Asn Ile Gln Gly	230	His Lys Gln Leu	235	Thr Asp Cys Ile Ser	240
Phe Leu Lys Glu Glu	245	Lys Leu Glu Gly	250	Asp Asn Arg Tyr Phe	255
Glu Asn Cys Gln Ser	260	Lys Gln Asn Ala	265	Thr Arg Lys Ile Arg	270
Leu Ser Leu Pro Cys	275	Thr Leu Asn Leu	280	Gln Leu Met Arg Phe	285
Phe Asp Arg Gln Thr	290	Gly His Lys Lys	295	Lys Leu Asn Thr Tyr	300
Gly Phe Ser Glu Ile	305	Leu Asp Met Glu	310	Pro Tyr Val Glu His	315
Gly Gly Ser Tyr Val	320	Tyr Glu Leu Ser	325	Ala Val Leu Ile His	330
Gly Val Ser Ala Tyr	335	Ser Gly His Tyr	340	Ile Ala His Val Lys	345
Pro Gln Ser Gly Glu	350	Trp Tyr Lys Phe	355	Asn Asp Glu Asp Ile	360
Lys Met Glu Gly Lys	365	Lys Leu Gln Leu	370	Gly Ile Glu Glu Asp	375
Ala Glu Pro Ser Lys	380	Ser Gln Thr Arg	385	Lys Pro Lys Cys Gly	390
Gly Thr His Cys Ser	395	Arg Asn Ala Tyr	400	Met Leu Val Tyr Arg	405
Gln Thr Gln Glu Lys	410	Pro Asn Thr Thr	415	Val Gln Val Pro Ala	420
Leu Gln Glu Leu Val	425	Asp Arg Asp Asn	430	Ser Lys Phe Glu Glu	435
Cys Ile Glu Met Ala	440	Glu Met Arg Lys	445	Gln Ser Val Asp Lys	450
Lys Ala Lys His Glu	455	Glu Glu Val Lys	460	Leu Tyr Gln Arg Leu	465
Ala Gly Ala Glu Pro	470	Tyr Glu Phe Val	475	Ser Leu Glu Trp Leu	480
Lys Trp Leu Asp Glu	485	Ser Thr Pro Thr	490	Lys Pro Ile Asp Asn	495
Ala Cys Leu Cys Ser	500	His Asp Lys Leu	505	His Pro Asp Lys Ile	510
Ile Met Lys Arg Ile	515	Ser Glu Tyr Ala	520	Ala Asp Ile Phe Tyr	525
Arg Tyr Gly Gly Gly	530	Pro Arg Leu Thr	535	Val Lys Ala Leu Cys	540
Glu Cys Val Val Glu	545	Arg Cys Arg Ile	550	Leu Arg Leu Lys Asn	555
	560		565		570

```

Leu Asn Glu Asp Tyr Lys Thr Val Asn Asn Leu Leu Lys Ala Ala
575 580 585
Val Lys Gly Ser Asp Gly Phe Trp Val Gly Lys Ser Ser Leu Arg
590 595 600
Ser Trp Arg Gln Leu Ala Leu Glu Gln Leu Asp Glu Gln Asp Gly
605 610 615
Asp Ala Glu Gln Ser Asn Gly Lys Met Asn Gly Ser Thr Leu Asn
620 625 630
Lys Asp Glu Ser Lys Glu Glu Arg Lys Glu Glu Glu Glu Leu Asn
635 640 645
Phe Asn Glu Asp Ile Leu Cys Pro His Gly Glu Leu Cys Ile Ser
650 655 660
Glu Asn Glu Arg Arg Leu Val Ser Lys Glu Ala Trp Ser Lys Leu
665 670 675
Gln Gln Tyr Phe Pro Lys Ala Pro Glu Phe Pro Ser Tyr Lys Glu
680 685 690
Cys Cys Ser Gln Cys Lys Ile Leu Glu Arg Glu Gly Glu Glu Asn
695 700 705
Glu Ala Leu His Lys Met Ile Ala Asn Glu Gln Lys Thr Ser Leu
710 715 720
Pro Asn Leu Phe Gln Asp Lys Asn Arg Pro Cys Leu Ser Asn Trp
725 730 735
Pro Glu Asp Thr Asp Val Leu Tyr Ile Val Ser Gln Phe Phe Val
740 745 750
Glu Glu Trp Arg Lys Phe Val Arg Lys Pro Thr Arg Cys Ser Pro
755 760 765
Val Ser Ser Val Gly Asn Ser Ala Leu Leu Cys Pro His Gly Gly
770 775 780
Leu Met Phe Thr Phe Ala Ser Met Thr Lys Glu Asp Ser Lys Leu
785 790 795
Ile Ala Leu Ile Trp Pro Ser Glu Trp Gln Met Ile Gln Lys Leu
800 805 810
Phe Val Val Asp His Val Ile Lys Ile Thr Arg Ile Glu Val Gly
815 820 825
Asp Val Asn Pro Ser Glu Thr Gln Tyr Ile Ser Glu Pro Lys Leu
830 835 840
Cys Pro Glu Cys Arg Glu Gly Leu Leu Cys Gln Gln Gln Arg Asp
845 850 855
Leu Arg Glu Tyr Thr Gln Ala Thr Ile Tyr Val His Lys Val Val
860 865 870
Asp Asn Lys Lys Val Met Lys Asp Ser Ala Pro Glu Leu Asn Val
875 880 885
Ser Ser Ser Glu Thr Glu Glu Asp Lys Glu Glu Ala Lys Pro Asp
890 895 900
Gly Glu Lys Asp Pro Asp Phe Asn Gln Ile Met His Ala Phe Ser
905 910 915
Val Ala Pro Phe Asp Gln Asn Leu Ser Ile Asp Gly Lys Ile Leu
920 925 930
Ser Asp Asp Cys Ala Thr Leu Gly Thr Leu Gly Val Ile Pro Glu
935 940 945
Ser Val Ile Leu Leu Lys Ala Asp Glu Pro Ile Ala Asp Tyr Ala
950 955 960
Ala Met Asp Asp Val Met Gln Val Cys Met Pro Glu Glu Gly Phe
965 970 975
Lys Gly Thr Gly Leu Leu Gly His
980

```

&lt;210&gt; 18

&lt;211&gt; 227

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1973875CD1

&lt;400&gt; 18

Met Gly Asn Cys Val Gly Arg Gln Arg Arg Glu Arg Pro Ala Ala

1	5	10	15
Pro Gly His Pro Arg	Lys Arg Ala Gly Arg	Asn Glu Pro Leu Lys	
	20	25	30
Lys Glu Arg Leu Lys	Trp Lys Ser Asp Tyr	Pro Met Thr Asp Gly	
	35	40	45
Gln Leu Arg Ser Lys	Arg Asp Glu Phe Trp	Asp Thr Ala Pro Ala	
	50	55	60
Phe Glu Gly Arg Lys	Glu Ile Trp Asp Ala	Leu Lys Ala Ala Ala	
	65	70	75
Tyr Ala Ala Glu Ala	Asn Asp His Glu Leu	Ala Gln Ala Ile Leu	
	80	85	90
Asp Gly Ala Ser Ile	Thr Leu Pro His Gly	Thr Leu Cys Glu Cys	
	95	100	105
Tyr Asp Glu Leu Gly	Asn Arg Tyr Gln Leu	Pro Ile Tyr Cys Leu	
	110	115	120
Ser Pro Pro Val Asn	Leu Leu Leu Glu His	Thr Glu Glu Glu Ser	
	125	130	135
Leu Glu Pro Pro Glu	Pro Pro Pro Ser Val	Arg Arg Glu Phe Pro	
	140	145	150
Leu Lys Val Arg Leu	Ser Thr Gly Lys Asp	Val Arg Leu Ser Ala	
	155	160	165
Ser Leu Pro Asp Thr	Val Gly Gln Leu Lys	Arg Gln Leu His Ala	
	170	175	180
Gln Glu Gly Ile Glu	Pro Ser Trp Gln Arg	Trp Phe Phe Ser Gly	
	185	190	195
Lys Leu Leu Thr Asp	Arg Thr Arg Leu Gln	Glu Thr Lys Ile Gln	
	200	205	210
Lys Asp Phe Val Ile	Gln Val Ile Ile Asn	Gln Pro Pro Pro Pro	
	215	220	225

Gln Asp

&lt;210&gt; 19

&lt;211&gt; 403

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2323917CD1

&lt;400&gt; 19

Met Glu Lys Ser Gln	Lys Ile Asn Pro Phe	Ile Leu His Ile Leu	
1	5	10	15
Gln Glu Val Asp Glu	Glu Ile Lys Lys Gly	Leu Ala Ala Gly Ile	
	20	25	30
Thr Leu Asn Ile Ala	Gly Asn Asn Arg Leu	Val Pro Val Glu Arg	
	35	40	45
Val Thr Gly Glu Asp	Phe Trp Ile Leu Ser	Lys Ile Leu Lys Asn	
	50	55	60
Cys Leu Tyr Ile Asn	Gly Leu Asp Val Gly	Tyr Asn Leu Leu Cys	
	65	70	75
Asp Val Gly Ala Tyr	Tyr Ala Ala Lys Leu	Leu Gln Lys Gln Leu	
	80	85	90
Asn Leu Ile Tyr Leu	Asn Leu Met Phe Asn	Asp Ile Gly Pro Glu	
	95	100	105
Gly Gly Glu Leu Ile	Ala Lys Val Leu His	Lys Asn Arg Thr Leu	
	110	115	120
Lys Tyr Leu Arg Met	Thr Gly Asn Lys Ile	Glu Asn Lys Gly Gly	
	125	130	135
Met Phe Phe Ala Ala	Met Leu Gln Ile Asn	Ser Ser Leu Glu Lys	
	140	145	150
Leu Asp Leu Gly Asp	Cys Asp Leu Gly Met	Gln Ser Val Ile Ala	
	155	160	165
Phe Ala Thr Val Leu	Thr Gln Asn Gln Ala	Ile Lys Ala Ile Asn	
	170	175	180
Leu Asn Arg Pro Ile	Leu Tyr Gly Glu Gln	Glu Glu Ser Thr Val	
	185	190	195

```

His Val Gly Leu Met Leu Lys Glu Asn His Cys Leu Val Ala Leu
                200                205                210
His Met Cys Lys His Asp Ile Lys Asn Ser Gly Ile Gln Gln Leu
                215                220                225
Cys Asp Ala Leu Tyr Leu Asn Ser Ser Leu Arg Tyr Leu Asp Val
                230                235                240
Ser Cys Asn Lys Ile Thr His Asp Gly Met Val Tyr Leu Ala Asp
                245                250                255
Val Leu Lys Ser Asn Thr Thr Leu Glu Val Ile Asp Leu Ser Phe
                260                265                270
Asn Arg Ile Glu Asn Ala Gly Ala Asn Tyr Leu Ser Glu Thr Leu
                275                280                285
Thr Ser His Asn Arg Ser Leu Lys Ala Leu Ser Val Val Ser Asn
                290                295                300
Asn Ile Glu Gly Glu Gly Leu Val Ala Leu Ser Gln Ser Met Lys
                305                310                315
Thr Asn Leu Thr Phe Ser His Ile Tyr Ile Trp Gly Asn Lys Phe
                320                325                330
Asp Glu Ala Thr Cys Ile Ala Tyr Ser Asp Leu Ile Gln Met Gly
                335                340                345
Cys Leu Lys Pro Asp Asn Thr Asp Val Glu Pro Phe Val Val Asp
                350                355                360
Gly Arg Val Tyr Leu Ala Glu Val Ser Asn Gly Leu Lys Lys His
                365                370                375
Tyr Tyr Trp Thr Ser Thr Tyr Gly Glu Ser Tyr Asp His Ser Ser
                380                385                390
Asn Ala Gly Phe Ala Leu Val Pro Val Gly Gln Gln Pro
                395                400

```

<210> 20  
 <211> 372  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2754960CD1

```

<400> 20
Met Ser Lys Ala Phe Gly Leu Leu Arg Gln Ile Cys Gln Ser Ile
1      5      10
Leu Ala Glu Ser Ser Gln Ser Pro Ala Asp Leu Glu Glu Lys Lys
      20      25      30
Glu Glu Asp Ser Asn Met Lys Arg Glu Gln Pro Arg Glu Arg Pro
      35      40      45
Arg Ala Trp Asp Tyr Pro His Gly Leu Val Gly Leu His Asn Ile
      50      55      60
Gly Gln Thr Cys Cys Leu Asn Ser Leu Ile Gln Val Phe Val Met
      65      70      75
Asn Val Asp Phe Thr Arg Ile Leu Lys Arg Ile Thr Val Pro Arg
      80      85      90
Gly Ala Asp Glu Gln Arg Arg Ser Val Pro Phe Gln Met Leu Leu
      95      100     105
Leu Leu Glu Lys Met Gln Asp Ser Arg Gln Lys Ala Val Arg Pro
      110     115     120
Leu Glu Leu Ala Tyr Cys Leu Gln Lys Cys Asn Val Pro Leu Phe
      125     130     135
Val Gln His Asp Ala Ala Gln Leu Tyr Leu Lys Leu Trp Asn Leu
      140     145     150
Ile Lys Asp Gln Ile Thr Asp Val His Leu Val Glu Arg Leu Gln
      155     160     165
Ala Leu Tyr Thr Ile Arg Val Lys Asp Ser Leu Ile Cys Val Asp
      170     175     180
Cys Ala Met Glu Ser Ser Arg Asn Ser Ser Met Leu Thr Leu Pro
      185     190     195
Leu Ser Leu Phe Asp Val Asp Ser Lys Pro Leu Lys Thr Leu Glu
      200     205     210
Asp Ala Leu His Cys Phe Phe Gln Pro Arg Glu Leu Ser Ser Lys

```

	215		220		225
Ser Lys Cys Phe	Cys Glu Asn Cys Gly	Lys Lys Thr Arg Gly	Lys		
	230		235		240
Gln Val Leu Lys	Leu Thr His Leu Pro	Gln Thr Leu Thr Ile	His		
	245		250		255
Leu Met Arg Phe	Ser Ile Arg Asn Ser	Gln Thr Arg Lys Ile	Cys		
	260		265		270
His Ser Leu Tyr	Phe Pro Gln Ser Leu	Asp Phe Ser Gln Ile	Leu		
	275		280		285
Pro Met Lys Arg	Glu Ser Cys Asp Ala	Glu Glu Gln Ser Gly	Gly		
	290		295		300
Gln Tyr Glu Leu	Phe Ala Val Ile Ala	His Val Gly Met Ala	Asp		
	305		310		315
Ser Gly His Tyr	Cys Val Tyr Ile Arg	Asn Ala Val Asp Gly	Lys		
	320		325		330
Trp Phe Cys Phe	Asn Asp Ser Asn Ile	Cys Leu Val Ser Trp	Glu		
	335		340		345
Asp Ile Gln Cys	Thr Tyr Gly Asn Pro	Asn Tyr His Trp Gln	Glu		
	350		355		360
Thr Ala Tyr Leu	Leu Val Tyr Met Lys	Met Glu Cys			
	365		370		

<210> 21  
 <211> 94  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3092341CD1

<400> 21	
Met Leu Arg Gly Val	Leu Gly Lys Thr Phe Arg Leu Val Gly Tyr
1	5 10 15
Thr Ile Gln Tyr Gly	Cys Ile Ala His Cys Ala Phe Glu Tyr Val
	20 25 30
Gly Gly Val Val Met	Val Pro Met Gly His Val Trp Leu Glu Gly
	35 40 45
Asp Asn Leu Gln Asn	Ser Thr Asp Ser Arg Cys Tyr Gly Pro Ile
	50 55 60
Pro Tyr Gly Leu Ile	Arg Gly Arg Ile Phe Phe Lys Ile Trp Leu
	65 70 75
Leu Ser Asp Phe Gly	Phe Leu Arg Ala Ser Pro Asn Gly His Arg
	80 85 90
Phe Ser Asp Asp	

<210> 22  
 <211> 248  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3658034CD1

<400> 22	
Met Asn Thr Glu Arg	Thr Asn Ile Gln Val Thr Val Thr Gly Pro
1	5 10 15
Ser Ser Pro Ser Pro	Val Lys Phe Leu Ile Asp Thr His Asn Arg
	20 25 30
Leu Leu Leu Gln Thr	Ala Glu Leu Ala Val Val Gln Pro Thr Ala
	35 40 45
Val Asn Ile Ser Ala	Asn Gly Phe Gly Phe Ala Ile Cys Gln Leu
	50 55 60
Asn Val Val Tyr Asn	Val Lys Ala Ser Gly Ser Ser Arg Arg Arg
	65 70 75
Arg Ser Ile Gln Asn	Gln Glu Ala Phe Asp Leu Asp Val Ala Val
	80 85 90

```

Lys Glu Asn Lys Asp Asp Leu Asn His Val Asp Leu Asn Val Cys
    95 100 105
Thr Ser Phe Ser Gly Pro Gly Arg Ser Gly Met Ala Leu Met Glu
    110 115 120
Val Asn Leu Leu Ser Gly Phe Met Val Pro Ser Glu Ala Ile Ser
    125 130 135
Leu Ser Glu Thr Val Lys Lys Val Glu Tyr Asp His Gly Lys Leu
    140 145 150
Asn Leu Tyr Leu Asp Ser Val Asn Glu Thr Gln Phe Cys Val Asn
    155 160 165
Ile Pro Ala Val Arg Asn Phe Lys Val Ser Asn Thr Gln Asp Ala
    170 175 180
Ser Val Ser Ile Val Asp Tyr Tyr Glu Pro Arg Arg Gln Ala Val
    185 190 195
Arg Ser Tyr Asn Ser Glu Val Lys Leu Ser Ser Cys Asp Leu Cys
    200 205 210
Ser Asp Val Gln Gly Cys Arg Pro Cys Glu Asp Gly Ala Ser Gly
    215 220 225
Ser His His His Ser Ser Val Ile Phe Ile Phe Cys Phe Lys Leu
    230 235 240
Leu Tyr Phe Met Glu Leu Trp Leu
    245

```

<210> 23  
 <211> 520  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3883861CD1

```

<400> 23
Met Val Ala Arg Val Gly Leu Leu Leu Arg Ala Leu Gln Leu Leu
    1 5 10 15
Leu Trp Gly His Leu Asp Ala Gln Pro Ala Glu Arg Gly Gly Gln
    20 25 30
Glu Leu Arg Lys Glu Ala Glu Ala Phe Leu Glu Lys Tyr Gly Tyr
    35 40 45
Leu Asn Glu Gln Val Pro Lys Ala Pro Thr Ser Thr Arg Phe Ser
    50 55 60
Asp Ala Ile Arg Ala Phe Gln Trp Val Ser Gln Leu Pro Val Ser
    65 70 75
Gly Val Leu Asp Arg Ala Thr Leu Arg Gln Met Thr Arg Pro Arg
    80 85 90
Cys Gly Val Thr Asp Thr Asn Ser Tyr Ala Ala Trp Ala Glu Arg
    95 100 105
Ile Ser Asp Leu Phe Ala Arg His Arg Thr Lys Met Arg Arg Lys
    110 115 120
Lys Arg Phe Ala Lys Gln Gly Asn Lys Trp Tyr Lys Gln His Leu
    125 130 135
Ser Tyr Arg Leu Val Asn Trp Pro Glu His Leu Pro Glu Pro Ala
    140 145 150
Val Arg Gly Ala Val Arg Ala Ala Phe Gln Leu Trp Ser Asn Val
    155 160 165
Ser Ala Leu Glu Phe Trp Glu Ala Pro Ala Thr Gly Pro Ala Asp
    170 175 180
Ile Arg Leu Thr Phe Phe Gln Gly Asp His Asn Asp Gly Leu Gly
    185 190 195
Asn Ala Phe Asp Gly Pro Gly Gly Ala Leu Ala His Ala Phe Leu
    200 205 210
Pro Arg Arg Gly Glu Ala His Phe Asp Gln Asp Glu Arg Trp Ser
    215 220 225
Leu Ser Arg Arg Arg Gly Arg Asn Leu Phe Val Val Leu Ala His
    230 235 240
Glu Ile Gly His Thr Leu Gly Leu Thr His Ser Pro Ala Pro Arg
    245 250 255
Ala Leu Met Ala Pro Tyr Tyr Lys Arg Leu Gly Arg Asp Ala Leu

```

Leu Ser Trp Asp	260	Val Leu Ala Val	265	Gln Ser Leu Tyr Gly	270
Asp	275	Val Ala Val Gln	280	Lys	285
Pro Leu Gly Gly	290	Leu Pro Gly Lys Leu	295	Phe	300
Thr Asp Phe Glu	305	Thr Trp Asp Ser Tyr	310	Arg Arg	315
Pro Glu Thr Gln	320	Pro Lys Tyr Cys	325	His Ser Ser Phe Asp	330
Ile Thr Val Asp	335	Arg Gln Gln Gln Leu	340	Tyr Ile Phe Lys Gly	345
His Phe Trp Glu	350	Val Ala Ala Asp Gly	355	Asn Val Ser Glu Pro	360
Pro Leu Gln Glu	365	Arg Trp Val Gly Leu	370	Pro Pro Asn Ile Glu	375
Ala Ala Val Ser	380	Leu Asn Asp Gly Asp	385	Phe Tyr Phe Phe Lys	390
Gly Arg Cys Trp	395	Arg Phe Arg Gly Pro	400	Lys Pro Val Trp Gly	405
Pro Gln Leu Cys	410	Arg Ala Gly Gly Leu	415	Pro Arg His Pro Asp	420
Ala Leu Phe Phe	425	Pro Leu Arg Arg	430	Leu Ile Leu Phe Lys	435
Ala Arg Tyr Tyr	440	Val Leu Ala Arg Gly	445	Gly Leu Gln Val Glu	450
Tyr Tyr Pro Arg	455	Leu Gln Asp Trp	460	Gly Gly Ile Pro Glu	465
Val Ser Gly Ala	470	Leu Pro Arg Pro Asp	475	Gly Ser Ile Ile Phe	480
Arg Asp Asp Arg	485	Tyr Trp Arg Leu Asp	490	Gln Ala Lys Leu Gln	495
Thr Thr Ser Gly	500	Arg Trp Ala Thr Glu	505	Leu Pro Trp Met Gly	510
Trp His Ala Asn	515	Ser Gly Ser Ala Leu	520	Phe	

&lt;210&gt; 24

&lt;211&gt; 422

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4993873CD1

&lt;400&gt; 24

Met Gly Pro Ala Trp	1	Leu Trp Leu Leu Gly	10	Thr Gly Ile Leu Ala	15
Ser Val His Cys Gln	20	Pro Leu Leu Ala His	25	Gly Asp Lys Ser Leu	30
Gln Gly Pro Gln Pro	35	Pro Arg His Gln Leu	40	Ser Glu Pro Ala Pro	45
Ala Tyr His Arg Ile	50	Thr Pro Thr Ile Thr	55	Asn Phe Ala Leu Arg	60
Leu Tyr Lys Glu Leu	65	Ala Ala Asp Ala Pro	70	Gly Asn Ile Phe Phe	75
Ser Pro Val Ser Ile	80	Ser Thr Thr Leu Ala	85	Leu Leu Ser Leu Gly	90
Ala Gln Ala Asn Thr	95	Ser Ala Leu Ile Leu	100	Glu Gly Leu Gly Phe	105
Asn Leu Thr Glu Thr	110	Pro Glu Ala Asp Ile	115	His Gln Gly Phe Arg	120
Ser Leu Leu His Thr	125	Leu Ala Leu Pro Ser	130	Pro Lys Leu Glu Leu	135
Lys Val Gly Asn Ser	140	Leu Phe Leu Asp Lys	145	Arg Leu Lys Pro Arg	150
Gln His Tyr Leu Asp	155	Ser Ile Lys Glu Leu	160	Tyr Gly Ala Phe Ala	165



Phe	Ser	Ala	Asn	Phe	Thr	Asp	Ser	Val	Thr	Thr	Gly	Arg	Gln	Ile
			170						175					180
Asn	Asp	Tyr	Leu	Arg	Arg	Gln	Thr	Tyr	Gly	Gln	Val	Val	Asp	Cys
			185						190					195
Leu	Pro	Glu	Phe	Ser	Gln	Asp	Thr	Phe	Met	Val	Leu	Ala	Asn	Tyr
			200						205					210
Ile	Phe	Phe	Lys	Ala	Lys	Trp	Lys	His	Pro	Phe	Ser	Arg	Tyr	Gln
			215						220					225
Thr	Gln	Lys	Gln	Ala	Ser	Phe	Phe	Val	Asp	Glu	Arg	Thr	Ser	Leu
			230						235					240
Gln	Val	Pro	Met	Met	His	Gln	Lys	Glu	Met	His	Arg	Phe	Leu	Tyr
			245						250					255
Asp	Gln	Asp	Leu	Ala	Cys	Thr	Val	Leu	Gln	Ile	Glu	Tyr	Arg	Gly
			260						265					270
Asn	Ala	Leu	Ala	Leu	Leu	Val	Leu	Pro	Asp	Pro	Gly	Lys	Met	Lys
			275						280					285
Gln	Val	Glu	Ala	Ala	Leu	Gln	Pro	Gln	Thr	Leu	Arg	Lys	Trp	Gly
			290						295					300
Gln	Leu	Leu	Leu	Pro	Ser	Leu	Leu	Asp	Leu	His	Leu	Pro	Arg	Phe
			305						310					315
Ser	Ile	Ser	Gly	Thr	Tyr	Asn	Leu	Glu	Asp	Ile	Leu	Pro	Gln	Ile
			320						325					330
Gly	Leu	Thr	Asn	Ile	Leu	Asn	Leu	Glu	Ala	Asp	Phe	Ser	Gly	Val
			335						340					345
Thr	Gly	Gln	Leu	Asn	Lys	Thr	Ile	Ser	Lys	Val	Ser	His	Lys	Ala
			350						355					360
Met	Val	Asp	Met	Ser	Glu	Lys	Gly	Thr	Glu	Ala	Gly	Ala	Ala	Ser
			365						370					375
Gly	Leu	Leu	Ser	Gln	Pro	Pro	Ser	Leu	Asn	Thr	Met	Ser	Asp	Pro
			380						385					390
His	Ala	His	Phe	Asn	Arg	Pro	Phe	Leu	Leu	Leu	Leu	Trp	Glu	Val
			395						400					405
Thr	Thr	Gln	Ser	Leu	Leu	Phe	Leu	Gly	Lys	Val	Val	Asn	Pro	Val
			410						415					420

Ala Gly

<210> 25  
 <211> 114  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5208004CD1

Met	Arg	Trp	Arg	Gln	Arg	Ser	Phe	Leu	Leu	Arg	Leu	Phe	Leu	Gly
1				5					10					15
Ser	Leu	Arg	Gly	Gly	Gln	His	His	Pro	Pro	Leu	Thr	Leu	Pro	Ser
			20						25					30
Ala	Ser	Ser	Leu	Pro	Phe	Ser	Thr	Leu	Ser	Leu	Leu	Leu	Ala	Ser
			35						40					45
Ser	Leu	Ser	Cys	Cys	Leu	Val	Ser	Pro	Cys	Pro	Lys	Thr	Pro	Gly
			50						55					60
Ser	Phe	Val	Leu	Leu	Pro	Trp	Pro	Pro	Pro	Arg	Arg	Arg	Ser	Gln
			65						70					75
Ala	Pro	Ser	Pro	Pro	Arg	Gly	Ile	His	Thr	Thr	Gly	Ser	Cys	Trp
			80						85					90
Gly	Trp	Gly	Ser	Pro	Ala	Gly	Phe	Leu	Met	Pro	Cys	Ala	Gln	Gly
			95						100					105
Ser	Ala	Ala	Val	Ile	Phe	Gly	Leu	Ser						
			110											

<210> 26  
 <211> 742  
 <212> PRT  
 <213> Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5267783CD1

&lt;400&gt; 26

Met	Pro	Ala	Gly	Gly	Lys	Gly	Ser	His	Pro	Ser	Ser	Thr	Pro	Gln
1				5					10					15
Arg	Val	Pro	Asn	Arg	Leu	Ile	His	Glu	Lys	Ser	Pro	Tyr	Leu	Leu
				20					25					30
Gln	His	Ala	Tyr	Asn	Pro	Val	Asp	Trp	Tyr	Pro	Trp	Gly	Glu	Glu
				35					40					45
Ala	Phe	Asp	Lys	Ala	Arg	Lys	Glu	Asn	Lys	Pro	Ile	Phe	Leu	Ser
				50					55					60
Val	Gly	Tyr	Ser	Thr	Cys	His	Trp	Cys	His	Met	Met	Glu	Glu	Glu
				65					70					75
Ser	Phe	Gln	Asn	Glu	Glu	Ile	Gly	Arg	Leu	Leu	Ser	Glu	Asp	Phe
				80					85					90
Val	Ser	Val	Lys	Val	Asp	Arg	Glu	Glu	Arg	Pro	Asp	Val	Asp	Lys
				95					100					105
Val	Tyr	Met	Thr	Phe	Val	Gln	Ala	Thr	Ser	Ser	Gly	Gly	Gly	Trp
				110					115					120
Pro	Met	Asn	Val	Trp	Leu	Thr	Pro	Asn	Leu	Gln	Pro	Phe	Val	Gly
				125					130					135
Gly	Thr	Tyr	Phe	Pro	Pro	Glu	Asp	Gly	Leu	Thr	Arg	Val	Gly	Phe
				140					145					150
Arg	Thr	Val	Leu	Leu	Arg	Ile	Arg	Glu	Gln	Trp	Lys	Gln	Asn	Lys
				155					160					165
Asn	Thr	Leu	Leu	Glu	Asn	Ser	Gln	Arg	Val	Thr	Thr	Ala	Leu	Leu
				170					175					180
Ala	Arg	Ser	Glu	Ile	Ser	Val	Gly	Asp	Arg	Gln	Leu	Pro	Pro	Ser
				185					190					195
Ala	Ala	Thr	Val	Asn	Asn	Arg	Cys	Phe	Gln	Gln	Leu	Asp	Glu	Gly
				200					205					210
Tyr	Asp	Glu	Glu	Tyr	Gly	Gly	Phe	Ala	Glu	Ala	Pro	Lys	Phe	Pro
				215					220					225
Thr	Pro	Val	Ile	Leu	Ser	Phe	Leu	Phe	Ser	Tyr	Trp	Leu	Ser	His
				230					235					240
Arg	Leu	Thr	Gln	Asp	Gly	Ser	Arg	Ala	Gln	Gln	Met	Ala	Leu	His
				245					250					255
Thr	Leu	Lys	Met	Met	Ala	Asn	Gly	Gly	Ile	Arg	Asp	His	Val	Gly
				260					265					270
Gln	Gly	Phe	His	Arg	Tyr	Ser	Thr	Asp	Arg	Gln	Trp	His	Val	Pro
				275					280					285
His	Phe	Glu	Lys	Met	Leu	Tyr	Asp	Gln	Ala	Gln	Leu	Ala	Val	Ala
				290					295					300
Tyr	Ser	Gln	Ala	Phe	Gln	Leu	Ser	Gly	Asp	Glu	Phe	Tyr	Ser	Asp
				305					310					315
Val	Ala	Lys	Gly	Ile	Leu	Gln	Tyr	Val	Ala	Arg	Ser	Leu	Ser	His
				320					325					330
Arg	Ser	Gly	Gly	Phe	Tyr	Ser	Ala	Glu	Asp	Ala	Asp	Ser	Pro	Pro
				335					340					345
Glu	Arg	Gly	Gln	Arg	Pro	Lys	Glu	Gly	Ala	Tyr	Tyr	Val	Trp	Thr
				350					355					360
Val	Lys	Glu	Val	Gln	Gln	Leu	Leu	Pro	Glu	Pro	Val	Leu	Gly	Ala
				365					370					375
Thr	Glu	Pro	Leu	Thr	Ser	Gly	Gln	Leu	Leu	Met	Lys	His	Tyr	Gly
				380					385					390
Leu	Thr	Glu	Ala	Gly	Asn	Ile	Ser	Pro	Ser	Gln	Asp	Pro	Lys	Gly
				395					400					405
Glu	Leu	Gln	Gly	Gln	Asn	Val	Leu	Thr	Val	Arg	Tyr	Ser	Leu	Glu
				410					415					420
Leu	Thr	Ala	Ala	Arg	Phe	Gly	Leu	Asp	Val	Glu	Ala	Val	Arg	Thr
				425					430					435
Leu	Leu	Asn	Ser	Gly	Leu	Glu	Lys	Leu	Phe	Gln	Ala	Arg	Lys	His
				440					445					450
Arg	Pro	Lys	Pro	His	Leu	Asp	Ser	Lys	Met	Leu	Ala	Ala	Trp	Asn
				455					460					465

Gly	Leu	Met	Val	Ser	Gly	Tyr	Ala	Val	Thr	Gly	Ala	Val	Leu	Gly	
				470					475					480	
Gln	Asp	Arg	Leu	Ile	Asn	Tyr	Ala	Thr	Asn	Gly	Ala	Lys	Phe	Leu	
				485					490					495	
Lys	Arg	His	Met	Phe	Asp	Val	Ala	Ser	Gly	Arg	Leu	Met	Arg	Thr	
				500					505					510	
Cys	Tyr	Thr	Gly	Pro	Gly	Gly	Thr	Val	Glu	His	Ser	Asn	Pro	Pro	
				515					520					525	
Cys	Trp	Gly	Phe	Leu	Glu	Asp	Tyr	Ala	Phe	Val	Val	Arg	Gly	Leu	
				530					535					540	
Leu	Asp	Leu	Tyr	Glu	Ala	Ser	Gln	Glu	Ser	Ala	Trp	Leu	Glu	Trp	
				545					550					555	
Ala	Leu	Arg	Leu	Gln	Asp	Thr	Gln	Asp	Arg	Leu	Phe	Trp	Asp	Ser	
				560					565					570	
Gln	Gly	Gly	Gly	Tyr	Phe	Cys	Ser	Glu	Ala	Glu	Leu	Gly	Ala	Gly	
				575					580					585	
Leu	Pro	Leu	Arg	Leu	Lys	Asp	Asp	Gln	Asp	Gly	Ala	Glu	Pro	Ser	
				590					595					600	
Ala	Asn	Ser	Val	Ser	Ala	His	Asn	Leu	Leu	Arg	Leu	His	Gly	Phe	
				605					610					615	
Thr	Gly	His	Lys	Asp	Trp	Met	Asp	Lys	Cys	Val	Cys	Leu	Leu	Thr	
				620					625					630	
Ala	Phe	Ser	Glu	Arg	Met	Arg	Arg	Val	Pro	Val	Ala	Leu	Pro	Glu	
				635					640					645	
Met	Val	Arg	Ala	Leu	Ser	Ala	Gln	Gln	Gln	Thr	Leu	Lys	Gln	Ile	
				650					655					660	
Val	Ile	Cys	Gly	Asp	Arg	Gln	Ala	Lys	Asp	Thr	Lys	Ala	Leu	Val	
				665					670					675	
Gln	Cys	Val	His	Ser	Val	Tyr	Ile	Pro	Asn	Lys	Val	Leu	Ile	Leu	
				680					685					690	
Ala	Asp	Gly	Asp	Pro	Ser	Ser	Phe	Leu	Ser	Arg	Gln	Leu	Pro	Phe	
				695					700					705	
Leu	Ser	Thr	Leu	Arg	Arg	Leu	Glu	Asp	Gln	Ala	Thr	Ala	Tyr	Val	
				710					715					720	
Cys	Glu	Asn	Gln	Ala	Cys	Ser	Val	Pro	Ile	Thr	Asp	Pro	Cys	Glu	
				725					730					735	
Leu	Arg	Lys	Leu	Leu	His	Pro									
				740											

<210> 27  
 <211> 734  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5583922CD1

<400>	27														
Met	Trp	Gly	Leu	Leu	Leu	Ala	Leu	Ala	Ala	Phe	Ala	Pro	Ala	Val	
1				5					10					15	
Gly	Pro	Ala	Leu	Gly	Ala	Pro	Arg	Asn	Ser	Val	Leu	Gly	Leu	Ala	
				20					25					30	
Gln	Pro	Gly	Thr	Thr	Lys	Val	Pro	Gly	Ser	Thr	Pro	Ala	Leu	His	
				35					40					45	
Ser	Ser	Pro	Ala	Gln	Pro	Pro	Ala	Glu	Thr	Ala	Asn	Gly	Thr	Ser	
				50					55					60	
Glu	Gln	His	Val	Arg	Ile	Arg	Val	Ile	Lys	Lys	Lys	Lys	Val	Ile	
				65					70					75	
Met	Lys	Lys	Arg	Lys	Lys	Leu	Thr	Leu	Thr	Arg	Pro	Thr	Pro	Leu	
				80					85					90	
Val	Thr	Ala	Gly	Pro	Leu	Val	Thr	Pro	Thr	Pro	Ala	Gly	Thr	Leu	
				95					100					105	
Asp	Pro	Ala	Glu	Lys	Gln	Glu	Thr	Gly	Cys	Pro	Pro	Leu	Gly	Leu	
				110					115					120	
Glu	Ser	Leu	Arg	Val	Ser	Asp	Ser	Arg	Leu	Glu	Ala	Ser	Ser	Ser	
				125					130					135	
Gln	Ser	Phe	Gly	Leu	Gly	Pro	His	Arg	Gly	Arg	Leu	Asn	Ile	Gln	

	140		145		150
Ser Gly Leu Glu Asp	Gly Asp Leu Tyr Asp	Gly Ala Trp Cys Ala			
	155		160		165
Glu Glu Gln Asp Ala	Asp Pro Trp Phe Gln	Val Asp Ala Gly His			
	170		175		180
Pro Thr Arg Phe Ser	Gly Val Ile Thr Gln	Gly Ser Asn Ser Val			
	185		190		195
Trp Arg Tyr Asp Trp	Val Thr Ser Tyr Lys	Val Gln Phe Ser Asn			
	200		205		210
Asp Ser Arg Thr Trp	Trp Gly Ser Arg Asn	His Ser Ser Gly Met			
	215		220		225
Asp Ala Val Phe Pro	Ala Asn Ser Asp Pro	Glu Thr Pro Val Leu			
	230		235		240
Asn Leu Leu Pro Glu	Pro Gln Val Ala Arg	Phe Ile Arg Leu Leu			
	245		250		255
Pro Gln Thr Trp Leu	Gln Gly Gly Ala Pro	Cys Leu Arg Ala Glu			
	260		265		270
Ile Leu Ala Cys Pro	Val Ser Asp Pro Asn	Asp Leu Phe Leu Glu			
	275		280		285
Ala Pro Ala Ser Gly	Ser Ser Asp Pro Leu	Asp Phe Gln His His			
	290		295		300
Asn Tyr Lys Ala Met	Arg Lys Leu Met Lys	Gln Val Gln Glu Gln			
	305		310		315
Cys Pro Asn Ile Thr	Arg Ile Tyr Ser Ile	Gly Lys Ser Tyr Gln			
	320		325		330
Gly Leu Lys Leu Tyr	Val Met Glu Met Ser	Asp Lys Pro Gly Glu			
	335		340		345
His Glu Leu Gly Glu	Pro Glu Val Arg Tyr	Val Ala Gly Met His			
	350		355		360
Gly Asn Glu Ala Leu	Gly Arg Glu Leu Leu	Leu Leu Leu Met Gln			
	365		370		375
Phe Leu Cys His Glu	Phe Leu Arg Gly Asn	Pro Arg Val Thr Arg			
	380		385		390
Leu Leu Ser Glu Met	Arg Ile His Leu Leu	Pro Ser Met Asn Pro			
	395		400		405
Asp Gly Tyr Glu Ile	Ala Tyr His Arg Gly	Ser Glu Leu Val Gly			
	410		415		420
Trp Ala Glu Gly Arg	Trp Asn Asn Gln Ser	Ile Asp Leu Asn His			
	425		430		435
Asn Phe Ala Asp Leu	Asn Thr Pro Leu Trp	Glu Ala Gln Asp Asp			
	440		445		450
Gly Lys Val Pro His	Ile Val Pro Asn His	His Leu Pro Leu Pro			
	455		460		465
Thr Tyr Tyr Thr Leu	Pro Asn Ala Thr Val	Ala Pro Glu Thr Arg			
	470		475		480
Ala Val Ile Lys Trp	Met Lys Arg Ile Pro	Phe Val Leu Ser Ala			
	485		490		495
Asn Leu His Gly Gly	Glu Leu Val Val Ser	Tyr Pro Phe Asp Met			
	500		505		510
Thr Arg Thr Pro Trp	Ala Ala Arg Glu Leu	Thr Pro Thr Pro Asp			
	515		520		525
Asp Ala Val Phe Arg	Trp Leu Ser Thr Val	Tyr Ala Gly Ser Asn			
	530		535		540
Leu Ala Met Gln Asp	Thr Ser Arg Arg Pro	Cys His Ser Gln Asp			
	545		550		555
Phe Ser Val His Gly	Asn Ile Ile Asn Gly	Ala Asp Trp His Thr			
	560		565		570
Val Pro Gly Ser Met	Asn Asp Phe Ser Tyr	Leu His Thr Asn Cys			
	575		580		585
Phe Glu Val Thr Val	Glu Leu Ser Cys Asp	Lys Phe Pro His Glu			
	590		595		600
Asn Glu Leu Pro Gln	Glu Trp Glu Asn Asn	Lys Asp Ala Leu Leu			
	605		610		615
Thr Tyr Leu Glu Gln	Val Arg Met Gly Ile	Ala Gly Val Val Arg			
	620		625		630
Asp Lys Asp Thr Glu	Leu Gly Ile Ala Asp	Ala Val Ile Ala Val			
	635		640		645

Asp	Gly	Ile	Asn	His	Asp	Val	Thr	Thr	Ala	Trp	Gly	Gly	Asp	Tyr
				650					655					660
Trp	Arg	Leu	Leu	Thr	Pro	Gly	Asp	Tyr	Met	Val	Thr	Ala	Ser	Ala
				665					670					675
Glu	Gly	Tyr	His	Ser	Val	Thr	Arg	Asn	Cys	Arg	Val	Thr	Phe	Glu
				680					685					690
Glu	Gly	Pro	Phe	Pro	Cys	Asn	Phe	Val	Leu	Thr	Lys	Thr	Pro	Lys
				695					700					705
Gln	Arg	Leu	Arg	Glu	Leu	Leu	Ala	Ala	Gly	Ala	Lys	Val	Pro	Pro
				710					715					720
Asp	Leu	Arg	Arg	Arg	Leu	Glu	Arg	Leu	Arg	Gly	Gln	Lys	Asp	
				725					730					

&lt;210&gt; 28

&lt;211&gt; 2080

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 088718CB1

&lt;400&gt; 28

tgaaggactt	ttccaggacc	caaggccaca	cactggaagt	cttgacagctg	aagggaggca	60
ctccttggcc	tccgcagccg	atcacatgaa	gggtggtgcc	agtctcctgc	tctccgtcct	120
cctggcacag	gtgtggctgg	taccggctt	ggccccagct	cctcagtcgc	cagagacccc	180
agccccctag	aaccagacca	gcagggtagt	gcaggctccc	agggagggaag	aggaagatga	240
gcaggaggcc	agcgaggaga	aggccggtga	ggaagagaaa	gcctggctga	tggccagcag	300
gcagcagctt	gccaaggaga	cttcaaactt	cggattcagc	ctgctgcgaa	agatctccat	360
gaggcacgat	ggcaacatgg	tcttctctcc	atttggcatg	tccttggcca	tgacaggctt	420
gatgctgggg	gccacagggc	cgactgaaac	ccagatcaag	agagggctcc	acttgcaggc	480
cctgaagccc	accaagcccc	ggctcctgcc	ttccctcttt	aagggaactca	gagagacctt	540
ctcccgcaac	ctggaactgg	gcctctcaca	ggggagtttt	gccttcaccc	acaaggattt	600
tgatgtcaaa	gagactttct	tcaatttatc	caagaggtat	tttgatacag	agtgcgtgcc	660
tatgaatttt	cgcaatgcct	cacaggccaa	aaggctcatg	aatcattaca	ttaacaaaga	720
gactcggggg	aaaattccca	aactgtttga	tgagattaat	cctgaaacca	aattaattct	780
tgtcgattac	atcttgttca	aagggaatg	gttgaccccc	tttgacctg	ctttcaccca	840
agtcgacact	ttccacctgg	acaagtacaa	gaccattaag	gtgccccatga	tgtacggtgc	900
aggcaagttt	gcctccacct	ttgacaagaa	ttttcgttgt	catgtcccca	aactgcccta	960
ccaaggaaat	gccaccatgc	tggtggtcct	catggagaaa	atgggtgacc	acctcgccct	1020
tgaaggactt	ctgaccacag	acttgggtgga	gacatggctc	agaaacatga	aaaccagaaa	1080
catggaagtt	ttctttccga	agttcaagct	agatcagaag	tatgagatgc	atgagctgct	1140
taggcagatg	ggaatcagaa	gaatcttctc	accctttgct	gaccttagtg	aactctcagc	1200
tactggaaga	aatctccaag	tatccagggt	tttacaagaa	acagtgattg	aagttgatga	1260
aaggggcact	gaggcagtg	caggaatctt	gtcagaaaat	actgcttatt	ccatgcctcc	1320
tgtcatcaaa	gtggaccggc	catttcattt	catgatctat	gaagaaacct	ctggaatgct	1380
tctgtttctg	ggcagggtgg	tgaatccgac	tctctataaa	ttcaggacat	gcataagcaa	1440
cttcgtgctg	tagtagatgc	tgaatctgag	gtatcaaaaca	cacacaggat	accagcaatg	1500
gatggcaggg	gagagtgttc	cttttgttct	taactagttt	aggggtgtct	caaataaata	1560
cagtagtccc	cacttatctg	aggggggatac	atcaaagac	ccccagcaga	tgccctgaaac	1620
ggtggacagt	gctgaacctt	atatatatatt	tttctacac	atacatacct	atgataaagt	1680
ttaattttata	aattaggcac	agtaagagat	taacaataat	aacaacatta	agtaaaatga	1740
gttacttgaa	cgcaagcact	gcaataccat	aacagtcaaa	ctgattatag	agaaggctac	1800
taagtgtactc	atgggcgagg	agcatagaca	gtgtggagac	attgggcaag	gggagaattc	1860
acatcctggg	tgggacagag	caggacgatg	caagattcca	tcccactact	cagaatggca	1920
tgctgcttaa	gacttttaga	ttgtttattt	ctggaatttt	tcatttaagt	tttttgacc	1980
atggttgacc	atggttaact	gagactgcag	aaagcaaaac	catggataag	ggaggactac	2040
tacaaaagca	ttaaattgat	acatatattt	taaaaaaaa			2080

&lt;210&gt; 29

&lt;211&gt; 2225

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 114551CB1

```

<400> 29
gtgcttggcg cctgcgctgg acgactcggc cggtagtgga gatgtccggc cggcttaagc 60
gggagtcctg cgggtccact cgcgggaagc gagagtcctga gtcgcggggc agctccggtc 120
gcgtcaagcg ggagcgagat cgggagcggg agcctgaggg ggcgagctcc cggggcagcc 180
ctgtgcgcgt gaagcgggag ttcgagccgg cgagcgcgcg cgaggccccg gcttctgttg 240
tcccgtttgt gcgggtgaag cgggagcggc aggtcgatga ggactcggag cctgagcggg 300
aggtgcgagc aaagaatggc cgagtggatt ctgaggaccg gaggagccgc cactgcccg 360
acctggacac catcaacagg agtgtgctgg actttgactt tgagaaactg tgttctatct 420
ccctctcaca catcaatgct tatgcctgtc tgggtgtgtg caagtacttt caaggccggg 480
gtttgaagtc tcacgcctac attcacagtg tccagtttag ccaccatgtt tctctcaacc 540
tccacaccct caagttttac tgccttcag acaactatga gatcatcgat tctcattgg 600
aggatatcac gtatgtgttg aagcccactt tcacaaagca gcaaattgca aacttggaca 660
agcaagccaa attgtccggg gcatatgatg tactaccctta cctgcgggtt attgtgggac 720
tgaataacat aaaggccaat gattatgcca acgctgtcct tcaggctcta tctaattgtc 780
ctcctctccg gaactacttt ctggaagaag acaattataa gaacatcaa cgtcctccag 840
gggatatcat gttcttgttg gtccagcgtt ttggagagct gatgagaaag ctctggaacc 900
ctcgaaattt caaggcacat gtgtctcccc atgagatgct tcaggcagtt gtaacttgca 960
tgaagaagac ttttcagatc accaaacaag gagatggcgt tgactttctg tcttggtttc 1020
tgaatgctct gcactcagct ctggggggca caaagaagaa aaagaagact attgtgactg 1080
atgttttcca ggggtccatg aggatcttca ctaaaaagct tccccatcct gatctgccag 1140
cagaagaaaa agagcagttg ctccataatg acgagtacca ggagacaatg gtggagtcca 1200
cttttatgta cctgacgctg gaccttcccta ctgccccct ctacaaggac gagaaggagc 1260
agctcatcat tccccagtg ccactcttca acatcctggc taagttcaat ggcactcactg 1320
agaaggataa taagacttac aaggagaact ttctgaagcg cttccagctt accaagttgc 1380
ctccatatct aatcttttgt atcaagagat tcactaagaa caacttcttt gttgagaaga 1440
atccaactat tgtcaatttc cctattacaa atgtggatct gagagaatac ttgtctgaag 1500
aagtacaagc agtacacaag aataccacct atgacctcat tgccaacatc gtgcatgacg 1560
gcaagccctc cgagggtctc taccggatcc acgtgcttca tcatgggaca ggcaaatggt 1620
atgaattaca agacctccag gtgactgaca tcttcccca gatgatcaca ctgtcagagg 1680
cttacattca gatttggaag aggcgagata atgatgaaac caaccagcag ggggcttgaa 1740
ggaggcgctc agggctttgc tccaaggggc tgtggctgat gatggtaaat aagaacacag 1800
aagctgtagc tgaacacagg ctggctgggtg ggcttccctag gccagcccag cttgtatggg 1860
ttctggctac accagagcac caagagccca cttgcctggg atggccccac actgtcactc 1920
agctgttctt tgatcatttt tttctagatt gatgtcctt tctcccatgc attgagctcc 1980
catctagctt cagcagggca gaaccttct ccagatgtgt gtaacttatg tcttgagtat 2040
ctgggagtag ttgaagaaca gataattctt tccaaacatc aagccttggg attcttggag 2100
caagcagaaa gccagtaact tcgctctgtt agaggtggag gattttccca tgggtccccc 2160
catttcttga tttgtatttt tagatggatt aaatagtctc ctgtttttaa accaaaaaaa 2220
aaaaa 2225

```

```

<210> 30
<211> 3287
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1261376CB1

```

```

<400> 30
caggcgctgt acccggcgtg ctccgcgcgg cgccggcgaa gggacgtggg ggaaggggca 60
gggaggagga agcggtggtt gctgcggatg tgggtgtgag cgagcggcgc ctgaacacac 120
ggcggtctgc gagcgctga cccgggcctg cgccagagcc tgcaccgagc tccggggccc 180
cacaccgct acgggtggccc tgcgcccgtt gctactgagg cggcgtgctc tgcattcttc 240
gctgtccagg cctgcgggct ctgggtgtctg ctggctcctc cttgctcgcc tgctccctcc 300
tgcttgctct agtcaccgcc gccgcggcgg ccacagccat ggcgagaggt ggtgaaagcg 360
cggttccctc gggctcccag gatagcgcgc cggagccga aggtgctggc gccccgcgg 420
ccgctgcctc cgcggacgcc aaaatcatga aagtcaccgt gaagaccccg aaggaaaagg 480
aggaattcgc cgtgcccag aatagctcgg tccagcagtt taagggaagaa atctctaaac 540
gttttaaatc acatactgac caacttgtgt tgatatttgc tggaaaaatt ttgaaagatc 600
aagatacctt gagtcagcat ggaattcatg atggacttac tgttcaacct gtcattaaaa 660
cacaaaacag gctcaggat cattcagctc agcaaacaaa tacagctgga agcaatgtta 720
ctacatcatc aactcctaata agtaactcta catctgggtc tgctactagc aaccttttg 780
gttttaggtg ccttggggga cttgcaggct tgagttagct gggtttgaat actaccaact 840
tctctgaact acagagtcag atgcagcgac aacttttgtc taacctgaa atgatggtcc 900
agatcatgga aaatccctt gtccagagca tgctctcaaa tctgacctg atgagacgt 960
taattatggc caatccacaa atgcagcagt tgatacagag aaatccagaa attagtcata 1020

```

tgttgaataa	tccagatata	atgagacaaa	cgttggaaact	tgccaggaat	ccagcaatga	1080
tgcaggagat	gatgaggaac	caggaccgag	ctttgagcaa	cctagaaagc	atcccagggg	1140
gatataatgc	tttaaggcgc	atgtacacag	atattcagga	accaatgctg	agtgtctgcac	1200
aagagcagtt	tggttgtaaat	ccatttgctt	ccttgggtgag	caatacatcc	tctgggtgaag	1260
gtagtcaacc	ttcccgtaca	gaaaatagag	atccactacc	caatccatgg	gctccacaga	1320
cttcccagag	ttcatcagct	tccagcgga	ctgccagcac	tgtgggtggc	actactggta	1380
gtactgccag	tggcacttct	gggcagagta	ctactgcgcc	aaatttgggtg	cctggagtag	1440
gagctagtat	gttcaacaca	ccaggaatgc	agagcttggt	gcaacaaata	actgaaaacc	1500
cacaacttat	gcaaaacatg	ttgtctgccc	cctacatgag	aagcatgatg	cagtcactaa	1560
gccagaatcc	tgaccttgct	gcacagatga	tgctgaataa	tcccctattt	gctggaaatc	1620
ctcagcttca	agaacaaatg	agacaacagc	tcccaacttt	cctccaacaa	atgcagaatc	1680
ctgatacact	atcagcaatg	tcaaacctta	gagcaatgca	ggccttggtta	cagattcagc	1740
aggggtttaca	gacatttagca	acggaagccc	cgggcctcat	cccaggggtt	actcctggct	1800
tgggggcatt	aggaagcact	ggaggctctt	caggaaactaa	tggatctaac	gccacacctt	1860
gtgaaaacac	aagtcccaca	gcaggaacca	ctgaacctgg	acatcagcag	tttattcagc	1920
agatgcttca	ggctcttgct	ggagtaaatic	ctcagctaca	gaatccagaa	gtcagatttc	1980
agcaacaact	ggaacaactc	agtgcgaatg	gatttttgaa	ccgtgaagca	aacttgcaag	2040
ctctaatagc	aacaggaggt	gatataaatg	cagctattga	aagggttact	ggctcccagc	2100
catcatagca	gcatttctgt	atcttgaaaa	aatgtaattt	atttttgata	acggctctta	2160
aacttttaaaa	tacctgcttt	atttcatttt	gactcttgga	attctgtgct	gttataaaca	2220
aacccaatat	gatgcatttt	aagggtggagt	acagtaagat	gtgtgggttt	ttctgtattt	2280
ttcttttctg	gaacagtggg	aattaaggct	actgcatgca	tcacttctgc	atttattgta	2340
attttttaaa	aacatcacct	tttatagttg	ggtgaccaga	ttttgtctcg	catctgtcca	2400
gtttatttgc	tttttaaaaca	ttagcctatg	gtagtaattt	atgtagaata	aaagcattaa	2460
aaagaagcaa	atcatttgca	ctctataatt	tgtggtacag	tattgcttat	tgtgactttg	2520
gcatgcattt	ttgcaaacaa	tgctgtaaga	tttatactac	tgataatttt	gttttatttg	2580
tatacaatat	agagtatgca	catttgggac	tgcatctctg	gaaacatact	gcaataggct	2640
ctctgagcaa	aacacctgta	actaaaaaag	tgaagataag	aaaatactct	taaagctgag	2700
tatttccctaa	ttgtatagaa	tcttacagca	tctttgacaa	acatctccca	gcaaaagtgc	2760
cgggttagtca	ggtttggtga	aaatacagta	gaaaagctga	ttctggttat	ctctttaagg	2820
acaattaatt	gtacagacac	ataatgtaac	attgtctcaa	cattcattca	cagattgcat	2880
gtaaattacc	ttaatctttg	tcagagctga	aggaacactg	tagtataccc	caaagtgcac	2940
ttgcctagga	cttctcagct	tctcccatag	gtagtttaac	aggcattaaa	atttgtaatt	3000
gaaatgttgc	tttcaactgaa	aaagtgtctt	gatgtttcag	ttatttttaa	tcgccataaa	3060
aaaaatagaac	tatctttttg	gttttatctgt	tttctcatgc	acaggcaata	gacaaaattta	3120
aatagagttg	tgagccaatt	gtttctgaag	tgtttttgta	gttctattaa	gaaatagtta	3180
aatattgtgc	ttttcagagc	ctcagagaca	gggggacgcg	gtgccggggt	ggggcagcgg	3240
aatctgtcct	ggatggggcc	agcttaaata	atactggcaa	ccaagat		3287

<210> 31  
 <211> 2412  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1299481CB1

<400> 31						
gggctgtgtg	cgccggcgcc	ggcggcgcc	gaggggggatg	gagcgagcgc	cgagccgggt	60
cagagttgaa	caatgaccat	agttgacaaa	gcttctgaat	cttcagaccc	atcagcctat	120
cagaatcagc	ctggcagctc	cgaggcagtc	tcacctggag	acatggatgc	aggttctgcc	180
agctgggggtg	ctgtgtcttc	attgaatgat	gtgtcaaatc	acacactttc	tttaggacca	240
gtacctgggtg	ctgtagttta	ttcgagttca	tctgtacctg	ataaatcaaa	accatcacca	300
caaaaggatc	aagccctagg	tgatggcatc	gctcctccac	agaaagtct	tttcccatct	360
gagaagattt	gtcttaagtg	gcaacaaact	catagagttg	gagctgggct	ccagaatttg	420
ggcaatacct	gttttgccaa	tcagcactg	cagtgtttta	cctacacacc	acctcttgcc	480
aattacatgc	tatcacatga	acactccaaa	acatgtcatg	cagaaggctt	ttgtatgatg	540
tgtacaatgc	aagcacatat	tacccaggca	ctcagtaatc	ctggggacgt	tattaaacca	600
atgtttgtca	tcaatgagat	gcggcgtata	gctaggcacc	tccgttttgg	aaaccaagaa	660
gatgcccatg	aattccttca	atacactgtt	gatgctatgc	agaaagcatg	cttgaatggc	720
agcaataaat	tagacagaca	caccagggcc	accactcttg	tttgtcagat	tttgaggaga	780
tacctaagat	ctagagtcaa	atgttttaaa	tgcaagggcg	tttcagatac	ttttgatcca	840
tatcttgata	taacattgga	gataaaggct	gctcagagtg	tcaacaaggc	attggagcag	900
tttgtgaagc	cggaacagct	tgatggagaa	aactcgtaca	agtgcagcaa	gtgtaaaaag	960
atggttccag	cttcaaagag	gttcaactatc	catagatcct	ctaattgtct	tacactttct	1020
ctgaaacggt	ttgcaaattt	taccgggtgga	aaaattgcta	aggatgtgaa	ataccctgag	1080

tatcttgata	ttcggccata	tatgtctcaa	cccaacggag	agccaattgt	ctacgtcttg	1140
tatgcagtcg	tggteccacac	tggttttaat	tgccatgctg	gccattactt	ctgctacata	1200
aaagctagca	atggcctctg	gtatcaaagt	aatgactcca	ttgtatctac	cagtgtatatt	1260
agatcgggtac	tcagccaaca	agcctatgtg	ctcttttata	tcaggtccca	tgatgtgaaa	1320
aatggagggtg	aacttactca	tcccacccat	agccccggcc	agtccctctc	ccgccccgtc	1380
atcagtcagc	gggttgtcac	caacaaacag	gctgcgccag	gctttatcgg	accacagctt	1440
ccctctcaca	tgataaagaa	tccacctcac	ttaaatggga	ctggaccatt	gaaagacacg	1500
ccaagcagtt	ccatgtcgag	tcctaacggg	aattccagtg	tcaacagggc	tagtccgtgt	1560
aatgcttcag	cttctgtcca	aaactgggtc	gttaatatgt	cctcagtgat	cccagaacat	1620
cctaagaaac	aaaaaattac	aatcagttat	cacaacaagt	tgccctgttc	ccagtgtcag	1680
tctcaacctc	accttcatag	taattctttg	gagaacccta	ccaagcccgt	tccctcttct	1740
accattacca	attctgcagt	acagtctacc	tccaacgcac	ctacgatgtc	agtttctagt	1800
aaagttaaca	aaccgatccc	ccgcagtgaa	tctgtctccc	agcccgtagt	gaatggcaaa	1860
tccaagtcga	actccagcgt	gctggtgcc	tggtgcgccg	agtcctctga	ggactctgac	1920
gaggagtcaa	aggggctggg	caaggagaat	gggattggta	cgattgtgag	ctcccactct	1980
cccgcccaag	atgccgaaga	tgaggaggcc	actccgcacg	agcttcaaga	acccatgacc	2040
ctaaacggtg	ctaatagtgc	agacagcgac	agtgaccgca	aagaaaacgg	cctagcgctc	2100
gatggtgcc	gctgccaaag	ccagcctgcc	ctgcactcag	aaaatccctt	tgctaaggca	2160
aacggtcttc	ctggaaagt	gatgcctgct	cttttgctgt	ctctcccaga	agacaaaatc	2220
ttagagacct	tcaggcttag	caacaaactg	aaaggctcga	cggatgaaat	gagtgcacct	2280
ggagcagaga	ggggccctcc	cgaggaccgc	gacgccgagc	ctcagcctgg	cagccccgcc	2340
gccgaatccc	tggaggagcc	agatgcggcc	gcaagcttat	tccccttag	tgagggttaa	2400
ttttagcttg	ca					2412

<210> 32  
 <211> 2286  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1873139CB1

<400> 32						
gaaaacccgg	cgctccggaag	cccgtgcttt	ctttgacgca	agggctcgag	acgcagccgc	60
cgctccgga	gcgcccggct	agaagcgaca	ccagacggag	cctccggagt	tccctccgcc	120
ccacctcgcc	gggtcctgga	gccgcagtc	tcccagctgc	cctcctcggt	gccatggagt	180
gtccacacct	gagctccagc	gtctgcattg	ctccggactc	agccaagttc	cccaacggct	240
ccccgtcgct	ctgggtgctgc	agcgttatcg	ctgtgatgat	tttgtgggta	atgacaccaa	300
gctgggactg	gtacagaaaag	tcagagaaca	cttcagaaac	ttggaaaact	cagctttcac	360
agctgacagg	cataagaaaa	gaaaactttt	ggaaaactca	acactaaaca	gcaagttatt	420
aaaagttaaat	ggaagcacca	ctgccatttg	tgccacaggc	cttcggaatt	tggggaacac	480
atgtttcatg	aatgccatcc	ttcagtcact	cagtaacatt	gagcagtttt	gctgtttatt	540
caaagcaactg	cccgcctggg	agttaaggaa	tgggaaaaca	gcaggaaggc	ggacatacca	600
caccaggagc	caaggggata	acaatgtgtc	tttggtagaa	gagtttagaa	agacactctg	660
tgctttatgg	caaggcagcc	agactgcatt	tagcccagag	tccttatatt	atgttgtttg	720
gaagattatg	ccaaacttta	ggggctatca	acagcaggac	gccccatgat	tcctgcctga	780
ccctttggac	cacctacact	tggaacttca	ggcggttttc	aacgggtgtt	cccgctcagc	840
aattctgcag	gagaattcta	ctctgtctgc	aagtaacaag	tgttgcataa	atggagcatc	900
tactgttgtc	acggctatat	tcggaggcat	tctccaaaat	gaggttaact	gcctcatatg	960
tgggacagaa	tctagaaaag	ttgatccatt	cctagacctt	tcattagata	ttccaagtca	1020
gttcagaagt	aagcgtctca	agaatcaaga	aaatggacca	gtttgtctgt	tacgagattg	1080
tcttcgcagt	tttaccgact	tagaagaact	tgatgagaca	gagttatata	tgtgccataa	1140
atgcaaaaag	aaacaaaagt	ccacaaaaaa	gttttggtatt	caaaaactac	ccaaggtgct	1200
atgcttacat	ttgaaaagat	ttcattggac	agcatattta	agaaaataag	ttgatacata	1260
cgtagaattt	ccactgagag	gcctagacat	gaaatgctac	ttactagagc	ctgagaacag	1320
tggcccgagg	agctgcctgt	atgacctcgc	cgctgtgggt	gtgcaccatg	gttccggggg	1380
tggttcttga	cattacacag	catacgcaac	tcacgaaggc	cgctggttcc	acttcaatga	1440
cagtactgta	acactgactg	acgaggagac	tgtggtgaag	gcgaaggcct	acatcctttt	1500
ctacgtggaa	caccaggcca	aagctggatc	ggataaaact	taataccttc	tccaaatcat	1560
cattaccaca	ccataccaga	gaaacatttc	cagtttttcca	caaatacttg	atacaagatt	1620
taatttcatt	atgcactttt	caatttcccta	tttttgattt	agttttgtca	atggtagtga	1680
cttactgaac	atgggcacca	actaattttg	ttgttgttct	accagaaaac	ctcagcagat	1740
gttttgattt	gctgcttttag	ttgtaataat	tcaattttta	taggtagttg	taagaactta	1800
gtcttatattg	actttttttat	tttatgttaa	ggttttccagt	tctcactttt	aggcacattt	1860
acatcaatgc	ttttgttccct	ctcacatgct	gaaagcaaga	tgtgttccct	attgtgaaga	1920
gcgacacaa	tgccctgctgc	ctttccacag	ctataatgga	catcagggtg	actctaaatc	1980



```

aaggatcatg tgtgcacaat acttgtggcc cacaaaatTT cacaatgact gctgaggaat 2040
cattcttttt gcctgtaaaa tataacaaag ggcattcatta agtagaccag gtaattactg 2100
cttgtctctc aaggctgctg tctttatcag cactaactaa ataaatttgt tggttcagtt 2160
gtacttgtcc tgcaaataca agaattactc tcttTgttgg tttttttggT tttggggcat 2220
acttgtttgc ggggaggtaa gatgggagta aagaccaaT acatgtaatg tttaaaaaaa 2280
aaaaaa

```

```

<210> 33
<211> 873
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1903112CB1

```

```

<400> 33
taaaaacatt tatcatcttt gaaagcactc ttacgagact tttattcatc agaatcagtg 60
aaatgtttct ctttttttga cagggTcaaa tagaaattga cgatcaggtg gaaggactcc 120
aatatctagc ttctcgatat gatttcattg acttagatcg tgtgggcac caccgctggT 180
cctatggagg atacctctcc ctgatggcat taatgcagag gtcagatata ttcagggttg 240
ctattgctgg ggcctcagtc actctgtgga tcttctatga tacaggatac accgaacgtT 300
atatgggtca ccctgaccag aatgaacagg gctattactt aggatctgtg gccatgcaag 360
cagaaaagtt cccctctgaa ccaaatcgtt tactgctctt acatggtttc ctggatgaga 420
atgtccattt tgcacatacc agtatattac tgagtttttt agtgagggtc ggaaagccat 480
atgattttaca gatctatcct caggagagac acagcataag agttcctgaa tcgggagaac 540
attatgaact gcatcttttg cactaccttc aagaaaacct tggatcacgt attgctgctc 600
taaaagtgat ataattttga cctgtgtaga actctctggT atacactggc tatTTaacca 660
aatgaggagg tttaatcaat agaaaacaca gaattgatca tcacattttg atacctgcca 720
tgtaacatct actcctgaaa ataaatgtgg tgccatgcag gggTctacgg tttgtggtag 780
taatctaata ccttaacccc acatgctcaa aatcaaatga tacatatTcc tgagagaccc 840
agcaatacca taagaattac taaaaaaaaa aaa

```

```

<210> 34
<211> 1658
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1993044CB1

```

```

<400> 34
ggaagagttt cgatgtctct agggTgggcta gagcgtcctc ccgcgctcag tcgcgctgca 60
ggTgacggcg cccggaggct gtcgggaagt atgcggggTg acgtgtggTt gacgagctcg 120
gcggcgggTt tgctgagatc tgtggccggc ggcagctggT gcgggggggca gctgagagcg 180
agaggTggat cggggcggtg tgtggccagg gccatgacgg gcaatgccgg ggagTggTgc 240
ctcatggaaa gcgaccccg ggtcttcacc gagctcatta aaggattcgg ttgccgagga 300
gcccaggtag aagaaatatg gagtTtagag cctgagaatt ttgaaaaatt aaagccagTt 360
catgggttaa ttttctttt caagtggcag ccaggagaag aaccagcagg ctctgtggTt 420
caggactccc gacttgacac gatatttttt gctaagcagg gctaagcagg taattaataa tgctTgtgct 480
actcaagcca tagtgagtgt gttactgaac tgtaccaccc aggatgtcca tttaggcgag 540
acattatcag agTTtaaaaga attttcacaa agttttgatg cagctatgaa aggctTggca 600
ctgagcaatt cagatgtgat tgcacaagta cacaacagTt tcgccagaca gcaaatgtTt 660
gaatttgata cgaagacatc agcaaaaagaa gaagatgctt ttcactTtgt cagttatgtt 720
cctgttaatg ggagactgta tgaattagat ggattaagag aaggaccgat tgatttaggt 780
gcatgcaate aagatgattg gttcagTgca gtaaggcctg tcatagaaaa aaggatacaa 840
aagtacagtg aaggtgaaat tcgattTaat ttaatggcca ttgtgtctga cagaaaaatg 900
atatatgagc agaagatagc agagttacaa agacaacttg cagaggagga acccatggat 960
acagatcaag gtaatagtat gTTaagtgtc attcagTcag aagTtgccaa aaatcagatg 1020
cttattgaag aagaagtaca gaaattaaaa agatacaaga ttgagaatat cagaaggaag 1080
cataattatc tgcctttcat tatggaaattg ttaaagactt tagcagaaca ccagcagTta 1140
ataccactag tagaaaaggc aaaaagaaaa cagaacgcaa agaaagctca ggaaaccaa 1200
tgaagatgtt ttcagatatg tacacatttc tgcttctgca catattTtca tggaaaccat 1260
tatgtataaa gaacttagag caacatccta attggctcag tgcacgtTtg gcaatagtgc 1320
cagcctgtct tgtctTtaat gcatggattc ataaactTct tccctacctg catcatgtgc 1380
atgtagtgca tattaatga aagtgatatt aagaatgctt tcccaaattc cattattTga 1440

```

```

catttgaggt cctgaccact gtttaagttt tcttgggttg tcttaactaa ccattaatgg 1500
aggcctgaga aaatggcgcc aaacggatta ttgctttaat ctgtaaattt aaaataacgt 1560
aacaattttg cacattggtc agatccttgg attaaaccgg gttggtccta atttgcttcc 1620
tccatccata accgggaaaa aaaattccgg gttttccc 1658

```

```

<210> 35
<211> 1255
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2292182CB1

```

```

<400> 35
gcctaacgtc ttcagccag ccaggctgcc cttcttccct gcggaggagg ggectgggag 60
gtcgcgttgg cgggaggagg gttaccttcc ccagtctcgc tctggccgcc tgagccagga 120
ggaagcagcg gcgaggtctg cgggaggcat ggcgggagct ccggacgagc gccggcgggg 180
ccccgcggca ggggagcagc tgcagcagca acacgtctct tgccagggtc tccccgagcg 240
tctggcccag gggaatcccc agcaagggtt cttctccagc ttcttccca gcaaccagaa 300
gtgccagctt aggtcctctga agacgctgga gacaaatcca tatgtcaaac ttctgcttga 360
tgctatgaaa cactcaggtt gtgctgttaa caaagataga cacttttctt gcgaagactg 420
taatgggaaat gtcagtggag gttttgatgc ttcaacatct cagatagttt tgtgccagaa 480
taatattccat aatcaggccc atatgaacag agtggtcaca cacgagctta ttcattgcatt 540
tgatcattgt cgtgcccctg tgcactgggt caccaacatc agacatttgg cgtgctcaga 600
ggttcgagct gctaaccctta gtggagactg ctcacttgct aatgaaatat tcaggttaca 660
ttttggatta aaacaacacc accagacttg tgtgcgagac agagccactc tttctatcct 720
ggctgttagg aatatcagca aagaagtagc taaaaaggct gttgatgaag tttttgaatc 780
ttgtttcaat gaccatgaac cttttggaag gatcccatc aacaagactt atgcaagata 840
tgctcacaga gactttgaaa accgtgatcg gtattattca aatatatgag cacaatgaca 900
tttttatatt acagagcttc catcatcatg aagaaaaaaa aatttgggtc tcaagtgaac 960
aaacatataa aatgtaaaca atccctacaa tccagataaa acagagaaga ctgtgattct 1020
agcatattat cagaaaaagt aactatggca aaaaatgaaa ctgctttaa ctccaaggca 1080
gaaattgtat ctttatagct aaccatttag taaataaata cattacattt ttgtatttta 1140
gtcatttggt tttattacat gtgattattt taaaatttga ttatgaaacc tgtgaagtgt 1200
gtcttcccat gatgttgata ggtgaccact gttggtattt acattaaatg taaag 1255

```

```

<210> 36
<211> 1139
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2331301CB1

```

```

<400> 36
tgcttttaaa aaaaaaaaaa gcttctaate ctctggggaa agatctgctt ccttccccat 60
ttgatgcaat gaaataaccgt gtgactcttc cccagcgttg agatccatgg agtgggtgaga 120
cacatccagt ggacgcccag ggagatggag gtgtacatca ggcacttgga gaaggtgtta 180
aggcgtctatg tccagaggct gcagtggctg ctgtccggga gccgccgact gtttggcacc 240
gttttgagga gcaaagtatg catattgctg gacacgtcag ggtccatggg cccctacctg 300
cagcaggtga agacagagct ggttttgctg atttgggaac agctgcggaa gtgctgtgac 360
agttttaacc tgctcagctt tgcagagagc cttcagtcac gccaggacac gctggtggag 420
accacagatg cagcgtgtca tgaggctatg caatgggtga cccacctgca agctcagggc 480
agcacctcca tcttgcaagc attgctgaaa gctttcagtt tccatgatct ggaaggattg 540
tacctcctga ccgacggaaa gccagacaca agctgcagcc ttgtcctaaa tgaagtccaa 600
aaactcaggg agaaaagaga tgtgaaagtg cacaccattt ccttgaactg ctcagacaga 660
gcggcggttg agttcctgag aaagctggct tccctcaccg gcggacgcta tcaactgcct 720
gtgggtgagg acacactctc caaaattcac agcctgctga ccaaaggctt catcaatgaa 780
aaggatcgca cattgccacc atttgaagga gatgatttaa ggatcctggc ccaggaaatc 840
accaaggcca gaagcttctc ctggcaggcc caatccttca gatcccaact ccagaagaaa 900
aatgatgcag aaccaaggtt cactcttctc tagagaagtg ttctcagaaa agtcactctg 960
caaaggacca ctactgagc aaatctcagc cccgagggca ggatgggatg aaatgctgtg 1020
acctgcagga aacatgattc ctggtaccag gactctctgg aagctgagga aggaaagact 1080
ttgtcttttg tgtgaggggc cattcccggg tcttaatcta actctccagc cctgtccgg 1139

```

<210> 37  
 <211> 1753  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2517512CB1

<400> 37  
 gtggcggact agcccagget cccacacccg acgctctggc ccacacagac gctactctgt 60  
 agcatctcag gttccctctg gctgcactct ggaggaccac actcggtttc tttttggctg 120  
 ccagaggccc ccgcatccac cgtcgagctg ggagaaagat ggcggcagcc gtgcgacagg 180  
 atttggcccc gctcatgaat tccgagcggt ctcataaaga tctggctggc aagtatcgtc 240  
 agatcctgga aaaagccatt cagttatctg gagcagaaca actagaagct ttgaaagctt 300  
 ttgtggaagc aatggtaaat gagaatgtca gtctcgtgat ctgcggcgag ttgctgactg 360  
 atttttgcac acatcttctc aacttgccct atagcacagc caaagaaatc tatcacttca 420  
 ccttggaaaa gatccagcct agagtcattt catttgagga gcaggttgtt tccataagac 480  
 agcatcttgc atctatatat gagaaagaag aagattggag aaatgcagcc caagtgttgg 540  
 tgggaattcc tttggaacaa ggacaaaaac agtacaatgt agattataaa ctggagactt 600  
 acttgaagat tgctaggcta tatctggagg atgatgatcc agtccaggca gaggttaca 660  
 taaatcgagc atcgttgctt cagaaatgaat caaccaatga acaattacag atacattata 720  
 aggtatgcta tgcacgtggt cttgattata gaagaaaatt cattgaagct gcacaaaggt 780  
 acaatgagct ctcttacaag acaatagttc acgaaagtga aagactagag gccttaaaac 840  
 atgctttgca ctgtacgata ttagcatcag cagggcagca gcgttctcgg atgctagcta 900  
 ctctttttaa ggatgaaagg tgccagcaac ttgctgccta tgggaccta gagaaaatgt 960  
 atctagatag gatcatcaga ggaaatcaac ttcaagaatt tgctgccatg ctgatgcctc 1020  
 accaaaaagc aactacagct gatggttcca gcactcttga cagagctggt attgaacaca 1080  
 atttgttgtc tgcaagcaaa ttatataata atattacctt cgaagaactt ggagctcttt 1140  
 tagagatccc tgcagctaag gcggaagaaga tagcatctca aatgataacc gaaggacgta 1200  
 tgaatggatt tattgaccag attgatggaa tagttcattt tgaaacacga gaagccctgc 1260  
 caacgtggga taagcagata caatcacttt gtttccaagt agcccatgga agcccagatg gctcagttaa 1320  
 ttagtcaaac agcaccagaa tggacagcac aagccatgga agcccagatg cagtttccact 1380  
 tctttgcaga acttctgtgc acatgacatc tttttccatg ttgtgcagat cgtggttccg 1440  
 atctccaaag catttgcatc atgaccttat acatttcaat cctttttatg ctggattccg 1500  
 tttaaagaag acattattag agcaggaagt acaagcattt aaaatatgta gtgcccatat 1560  
 atttcagggt ctctgtgtat taagctaact cagatgtttt gaaagctttt tcttttaaca 1620  
 gaggtgaaat atctgtggct aaaaagtgtg agatttgtga taactttgta gtcattgtaa 1680  
 acttaagtgc ttcattgcctc tccaaatgtg gttattctaa taaatggaga aatgagccaa 1740  
 aaaaaaaaaaaa aaa 1753

<210> 38  
 <211> 580  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3489039CB1

<400> 38  
 cgaggcggccc gaagccgctc cggcggggac catgttgctt ccgaacatcc tgetcaccgg 60  
 tacaccaggg gttggaaaaa ccacactagg caaagaactt gcgtcaaaat caggactgaa 120  
 atacattaat gtgggtgatt tagctcgaga agagcaattg tatgatggct atgatgaaga 180  
 gtatgactgt cccatttttag atgaagacag agtagttgat gagttagata accaaatgag 240  
 agaaggtgga gttattgttg attaccatgg ttgtgatctt ttcctgaac gctggtttca 300  
 tatagttttt gtgctgagaa cagataccaa tgtattgtac gaaagacttg aaacaagggg 360  
 ttataatgag aagaaactaa cagacaatat tcagtgtgag atttttcaag ttctttatga 420  
 agaagccaca gcatcctaca aggaagaaat cgtgcacatg ctgcccagta ataaaccaga 480  
 agagctagaa aataatgtag atcagatctt gaaatggatt gagcagtgga tcaaagatca 540  
 taactcttga cttataaggc tagctactta ataatacact 580

<210> 39  
 <211> 2186  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5432879CB1

<400> 39  
 gtccggcgccg ggcggcgccg gtagcggagg agacgggttc aggcctccgg tgcggctgca 60  
 atgctgagct cccggggccga ggcggcgatg accggggccg acagggccat ccagcgcttc 120  
 ctgcggaaccg gggcgggcgt cagatataaa gtcatgaaga actggggagt tataggtgga 180  
 attgctgctg ctcttcgagc aggaatatat gttatttggg gtccattac agaaagaaag 240  
 aagcgtagaa aagggttgt gcttggtcct gttatttag ggaacacctg cttcatgaac 300  
 tccctgctac aaggcctgtc tgctgtcct gctttcatca ggtggctgga agagtccacc 360  
 tccagtagct ccagggatca gaaggagccc cctcacacc agtatttatc cttaacactc 420  
 ttgcaccttc tgaagcctt gtcctgccaa gaagtactg atgatgaggt cttagatgca 480  
 agctgcttgg tggatgtctt aagaatgtac agatggcaga tctcatcatt tgaagaacag 540  
 gatgctcacg aattattcca tgtcattacc tcgtcattgg aagatgagcg agaccgccag 600  
 cctcggtgca cacatttgtt tgatgtgcat tccctggagc agcagtcaga aataactccc 660  
 aaacaaatta cctgccgcac aagaggggtca cctcacccc catccaatca ctggaagtct 720  
 caacatcctt ttcattggaag actcactagt aatatggctt gcaaacactg tgaacaccag 780  
 agtctgttcc gatttgatag ctttgatagc ctttcactaa gtattccagc cgccacatgg 840  
 ggtcacccat tgaccctgga ccaactgcct caccacttca tctcatcaga atcagtgcgg 900  
 gatgttgtgt gtgacaactg tacaaagatt gaagccaagg gaacgttgaa cggggaaaag 960  
 gtggaacacc agaggaccac ttttggttaa cagttaaaac tagggaagct cctcagtggt 1020  
 ctctgcatcc acctacagcg gctgagctgg tccagccacg gcacgcctct gaagcggcat 1080  
 gagcacgtgc agttcaatga gttcctgatg atggacattt acaagtacca cctccttgga 1140  
 cataaaccta gtcaacacaa ccttaaactg aacaagaacc cagggccctac actggagctg 1200  
 caggatgggc cgggagcccc ctgaatcagc cagggggccc caaaacacag 1260  
 atttttatga atggcgctg ctccccatct ttattgccc aaagctgtcagc gccgatgccc 1320  
 tccctctccc cagttgttcc cgactacagc tctccacat acctcttccg gctgatggca 1380  
 gtgtgtgccc accatggaga catgcaactct ggacactttg tcaettaccg acggtcccca 1440  
 ccttctgcca ggaacctct ctcaactagc aatcagtggt tgtgggtctc cgatgacact 1500  
 gtccgcaagg ccagcctgca ggagggtcct tctccagcg cctacctgct gttctacgag 1560  
 cgctgctctt ccaggatgca gatggcactg tctgcaactgt ccaggaaaaa agtaaaactg 1620  
 cctcctgcaa ggctagagct gctccactag agccttcag ccttctggtg tgttctaaga 1740  
 tactgttgcg tgtgcaagcg gccccactag agccccagtt cagagctcct gaacagtcct 1800  
 gcaggctcca cctgggagcc ctggtgtgtt taattaaaag aactcagatt cttgaagcca cgttttcat 1920  
 gttctgtgt ctgatacagg taggtgttct cagaggggag gtaccttgt ctaatcaacg tttccactta 1980  
 attgtaatgt ttttaataag caggcccata aaaattgttg acaagaatta atgaaattat 2040  
 taaaggcaac aatttagaag aaaaagtgcc tttcacttcc gattgctttt gtacacgtc 2100  
 cattgtgaaa tattecttcc aggtacttca aaggatagca agagaacagg taaatgatgc 2160  
 ctaaagaaca ccttctttt tctatg 2186

<210> 40  
 <211> 2522  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5853753CB1

<400> 40  
 ctccggggcgt gctgcccggc gcggtaggtg gcgcgcgggt ccggcgggcg gttggcttga 60  
 gcgggaccgg agctgaggca ggaagagccg gcgccatggt ggagaaggag gaggctggcg 120  
 gcggcattag cgaggaggag gcggcacagt atgaccggca gatccgctg tggggactgg 180  
 agggcccagaa acggtgctgg gctctctcgg tgcctcttgt cggttgaaa ggacttgggg 240  
 ctgaaattgc caagaatctc atcttggcag gagtgaagg actgaccatg ctggatcacg 300  
 aacaggtaac tccagaagat cccggagctc agttcttgat tcgtactggg tctgttggcc 360  
 gaaatagggc tgaagcctct ttggagcgag ctccagaatct caaccccctg gtggatgtga 420  
 aggtggacac tgaggatata gagaagaaac cagagtcatt tttcactcaa tctgatgctg 480  
 tgtgtctgac ttgctgtccc agggatgtca tagttaaagt tgaccagatc tgtcacaaaa 540  
 atagcatcaa gttctttaca ggagatgttt ttggctacca tggatacaca tttgccaatc 600  
 taggagagca tgagtttgta gaggagaaaa ctaaagttagc caaagttagc caaggagtag 660  
 aagatggggc cgacaccaag agagcaaaa cttgattcttc tgagacaacg atggctaaaa 720  
 agaaggtggc cttctgcctt gttaaagaag ccttgagggt ggactggagc agtgagaaag 780  
 caaaggctgc tctgaagcgc acgacctccg actactttct ccttcaagtg ctcttaaaag 840

tccgtacaga	taaaggaaga	gatcccagtt	ctgatacata	tgaggaagat	tctgagttgt	900
tgctccagat	acgaaatgat	gtgcttgact	cactgggtat	tagtcctgac	ctgcttctctg	960
aggactttgt	caggtactgc	ttctccgaga	tggccccagt	gtgtgcggtg	gttggaggga	1020
ttttggcaca	ggaaattgtg	aaggccctgt	ctcagcgagg	ccctccctac	aacaacttct	1080
tcttcttcga	tgcatgaag	gggaatggga	ttgtggagtg	ccttgcccc	aagtgaactc	1140
aagatttggc	agccccagag	atgccaaactg	cagcatgccc	acctgtatcc	cctgtccct	1200
tccttcgatga	aggcatctcc	aggcaaggaa	aactgaagtc	attggcccga	tacaaaacat	1260
ttcttgcaac	gaaggaggtg	gtgcgcagct	gctgcttccc	atcaccagca	gctgctcgac	1320
aagggggcgca	gggtggctgt	ctttgttcca	gcactgttca	ggctgcctgt	catcccgggc	1380
ctgccagctc	ccctgagtga	tgagcacttc	caagcaccoc	tctgcccttt	ctctgtcctt	1440
atgctgtccc	ggcctcgcca	gccctctggg	gcattgtggg	agatgcctgc	caggaatgag	1500
caagetctgt	tgctcgggag	ccctctgtca	ccctcttgga	cttatccccc	acctgatacc	1560
ttatagagaa	aagtgtgaat	tcagggtggag	agtaggcccc	ggccccatga	ggcaccagt	1620
gaagcacagc	tccaagttca	gacagggtgcc	cttagagagg	aaaaccatga	caggcaaatg	1680
catttctctt	ggagtttgag	accctgacaa	acaacagggtg	gcatctgggtg	tgctgttctt	1740
gagttttcgt	ttaggattag	ttgagttcca	gctgggtttt	gggagaaagg	agatgctacc	1800
aagtcttgga	tgttagggcg	agaccctgca	agttgagtat	tagagagctt	gtctttcaag	1860
gcagggttct	ggggcttcag	ggctaggagg	gaggagcctg	cccttttaac	agaacccag	1920
tcacatgcgg	ctcaagtcac	tcagaggctg	ttgcatttca	gggctatggt	ggctccttgt	1980
ttacctctca	aaccacagct	gtttgtgttt	cacatatgtt	gtgaattttc	cttgggtctt	2040
tttaaaggaa	tgataataaa	gttacttgct	ttaggatttg	cttgtttttc	ttccacttca	2100
gaagcttctg	agaggggaatg	ggatgatcct	accagttgcc	ttttcagacc	tgaggctcta	2160
actcaagaga	ttctctctct	ccctcaccat	tcctgccacc	attttttctg	ggtgatgcag	2220
caagagttaa	attgttcaca	ttctagaatg	tgtagaagct	tctggctcca	gttgtctaat	2280
ggataattca	gctaaatagt	ggctattttc	agtggcaaga	attataataa	taaagggaag	2340
tcaaaagtga	tgcttcatat	gccccaaagt	ccacagccgc	tacgggtcac	agctttttatt	2400
ggaagaccag	gtttgccttg	atgcagccaa	agtgcgttcc	agttcacccc	actcttgctt	2460
gtgtttcttc	tttatgtaaa	ttgtgtgatg	tttcaataaa	tcagtctctt	tcctaaaaaa	2520
aa						2522

<210> 41  
 <211> 2167  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 411344CB1

<400> 41						
gcgcgggtctt	gggggggatgg	ttccatcatg	gcgtcaatgc	agaaacgact	acagaaagaa	60
ctgtttggctt	tgcaaaatga	cccacctcct	ggaatgacct	taaatgagaa	gagtgttcaa	120
aattcaatta	cacagtggat	tgtagacatg	gaagggtgcac	caggtagcctt	atatgaagg	180
gaaaaatttc	aacttctatt	taaatttagt	agtcgatatc	cttttgactc	tcctcaggtc	240
atgtttactg	gtgaaaatat	tcctgttcat	cctcatgttt	atagcaatgg	tcatatctgt	300
ttatccattc	taacagaaga	ctgggtcccca	gcgctctcag	tccaatcagt	ttgtcttagc	360
attattagca	tgctttccag	ctgcgaaggaa	aagagacgac	caccggataa	ttctttttat	420
gtgcgaacat	gtaacaagaa	tccaaagaaa	acaaaatggg	ggtatcatga	tgatacttgt	480
tgatgccact	gttatcatcc	tcctagcaga	agatagtcc	actgagaaaa	tgagcacttt	540
gatcattcag	tcctttgaact	ttaacctttg	actggaagtg	acctataggc	aatgaagact	600
acttctcttt	actgcatttt	tactcgtgtg	cattctgggc	gcatgttgat	cgctggttca	660
gtccaggcaa	ctgacatgct	tttattagtc	atacagtatt	aatgcagggtg	tcaggaaatg	720
tcaaatataa	ttccattttt	tatttttatt	tttttaagct	tttggaagg	ctccagggtcc	780
tcattgtattg	tgcaataaca	atgacttcct	tggcggtttt	ggtacgttca	ttgccggcaa	840
tggcggttgt	aacaggaaaa	gttttcatta	actcctgcc	ttcaatgatt	aatgcattgat	900
agggcctatg	aaatgaactt	actggttata	gtgggaatat	aaataaagt	agggatccaa	960
cattacttta	aaagtcaccc	caactgttta	tatttggatt	ctatgcactg	tgatcctaag	1020
gttaacagca	tgaattaaca	tgcgtcttta	aaggactgta	atgaaagatc	attgcatatt	1080
tattgaattg	tttatatcta	ctgtcaagtt	gttttgacat	ggaagatttt	caagtaacat	1140
tggcagagag	gtacagtatg	ttatccctat	gggtgaaaata	aattaatttg	ttgtatatag	1200
ttcctcaatc	tctgaagtaa	aggtatgagt	aatatagggt	atgaatgggt	taatcaaggc	1260
tttatttttg	aagtaagaaa	aatggcagtg	atgattaaat	tgctgcagtc	cataatttgg	1320
gcttgtttatt	tgtacattaa	agattttttc	caagtaagtt	acactctgtt	aacttctctg	1380
tagccatcag	catgagccct	actgcctaaa	cactatttca	tttatttatg	tttggaacc	1440
ccgtaaacat	ttttgtttgc	aatcttggtt	cttttggtat	aagtcaagtt	tgaatgttac	1500
aatactttta	ttgaaacttt	tgtaagtttt	tttcttgtaa	attttcttta	cttgtgagta	1560
tcattcttgc	ctttaatcct	gtaccctaaa	ataagaaata	catttttgac	agaggcttaa	1620

tgttttaaca	aaagagtgtg	gacattttta	ttttaaaatt	taggcaaaag	tcactatcaa	1680
atgggtgtct	atttgtctca	cacagccata	tagtttttcc	tggagggttt	tgttttgttg	1740
ttgttgaaaa	gactttgctt	acagctagat	gaaactttct	atagaaaaaa	aaaattgttg	1800
aaaggtccag	ttctcagtac	catgtgagtt	aatgatacta	caactaagtt	ctttttaaaa	1860
agtgattaat	gtattttata	aattaccctt	tcacatatgc	aaaatctgtt	tctactacaa	1920
tgttattttt	actaatgcct	tattgtggca	ctctttttga	aacatcctgc	agtgaatata	1980
tgaatcaatt	tgggctttaa	actgaaagcc	agttgggctg	aaaggtttga	aatacgtacc	2040
ccagtaaaac	cattcaatcc	ataattggta	aataatattt	taaaaatggg	tttaatccgt	2100
atagatgaca	ttttgtagct	ttggacatgg	tggtaatata	ggggcattaa	ttttacactc	2160
caaggta						2167

<210> 42  
 <211> 1826  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1256390CB1

<220>  
 <221> unsure  
 <222> 1755  
 <223> a, t, c, g, or other

<400> 42						
ccccgcctt	tctatagcat	agttctcttt	agggtggaatg	attcctataa	gattttctcat	60
tattaaatca	tgcatttttc	aagatggaat	caatctttga	tttaattctaa	gctgatattc	120
tcatttgta	gaagaacaac	ctacatgcta	gagagagagg	aggaaatata	cccacgacca	180
cacagccagt	tagtatccag	ttgggtgctg	actccagcca	ggtgtcctgc	ctcatggtag	240
ttaaatgata	tatagaaaag	gtaaaatttt	aaagaaatat	ttattaatat	attcctataa	300
aacattttta	aggtaaccac	ataaaaaatg	ttatattttc	cattccaaag	taaatgctaa	360
gcatgtttat	taatgaagca	gtacttctga	ttagtatatg	acattctgaa	gttaattaaa	420
ctcattgcac	taaatgtgtc	ttccttggtg	tagtggagga	tttgaggatt	ggaatataga	480
gtagagtgtc	tgtttaagcc	cttgcctgaa	accctttatc	ttttggggat	gctggttcct	540
ggagggtcgt	gatatgatag	atccttagcc	caacacagac	aagagattgt	ggacaagtca	600
gtgagtccat	ggagcctgga	gacgtattcc	tataacatat	accaccccat	gggagagatc	660
tatgagtggc	tgagagagat	cagtgaagaag	tacaaggaag	tggtgacaca	gcatttccta	720
ggagtgtac	atgagaccca	ccccatgtat	tatctgaaga	tcagccaacc	atctggtaat	780
cccaagaaaa	tcatttggat	ggactgtgga	attcagccca	gagaatggat	tgtctctgct	840
ttttgccaat	ggttcgtcaa	agaaattcta	caaaaccata	aagacaactc	aagtatacgc	900
aagctcctta	ggaacctgga	cttctatgtc	cttccagttc	ttaacataga	tggttatatc	960
tacatttggg	caactgatcg	tctttggagg	aaatcccgtt	caccccataa	taatggcaca	1020
tgttttggga	cggatctcaa	tcgaaatttc	aatgcattct	ggtgtagtat	tggtgcctct	1080
agaaactgcc	aagatcaaac	attctgtggg	acagggccag	tgtctgaacc	agagactaaa	1140
gctgttgcca	gcttcataga	gagcaagaag	gatgatattt	tgtgcttctc	gaccatgcac	1200
tcttatgggc	agttaattct	cacaccttac	ggctacacca	aaaataaaatc	aagtaaccac	1260
ccagaaatga	ttcaagtttg	acagaaggca	gcaaatgcat	tgaaagcaaa	gtatgggaacc	1320
aattatagag	ttggatcgag	tgcagatatt	ttatatgcct	catcagggtc	ttcaagagat	1380
tgggcccggg	acattgggat	tcccttctca	tatacgtttg	agctgaggga	cagtgggaaca	1440
tatgggtttg	ttctgccaga	agctcagatc	cagcccacct	gtgaggagac	catggaggct	1500
gtgctgtcag	tccctggatga	tgtgtatgcg	aaacactggc	actcggacag	tgtggaagg	1560
gtgacatctg	ccactatgct	gctgggctg	ctgggtgtct	gcatgtctct	tctctaagtg	1620
cattctgccc	aggttttttt	ctgccagaaa	ttccccttta	tttttccctc	tagtaaagtc	1680
tttctttgat	actaattctg	acattacctg	tgggaattttt	gctgacttcc	acagattttt	1740
ttttctctcc	ttctntaact	gtttgtacac	actttaatgt	aataatttac	agacttgtgt	1800
agtgagatag	gatattacat	gccctg				1826

<210> 43  
 <211> 1371  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1786774CB1

&lt;400&gt; 43

```

gcaaaaaattg tttcaaaaaa tttgtgagga ttttgattcc tatctcgttc aaatTTTTTgg 60
ttcatcgtca cctcaacctg gatttgggat gtcaggggaa gaactcagtg aatctactcc 120
agagcctcaa aaagaaattt ctgagtcatt gagtgccacc agagaccagg atgaagatga 180
taaggtccca gagccaacgt gggtgatga tctgcctgcc actacctcat ctgagggcac 240
caccaccccc aggccactgc tcagcacccc tgtggatggg gcagaggatc ctgatgtttt 300
ggaagccttg aagcctggaa actgtggtga atatgtggtt cgatggtatt atgacaaaca 360
ggteaactct tgtgccgat tttggttcag tggctgtaat ggctcaggaa atagattcaa 420
cagtgaaaaag gaatgtcaag aaacctgcat tcaaggatga gcaagtaaat tggcctgtct 480
ctatcaaaaag catagaactc cctaatttcc acatattcac ccaatacaaa tacagcacta 540
tatttgagtg tatactgagt atttacaact tatacatgta attgaattct cactacagcc 600
ctaggatgta catattatta accacttata taggtaagaa agctgaggct ctgagaagtt 660
tagtaacttg tcaactgtca cccaactaaa aagtttcaga gctgaggatt tagacttaga 720
gctgtgtaac ttcaatacac agactctatc tacttcacaa cctgcaatgt gattctgatt 780
ccttttaattc ctgttgatg tactatgtca gctcaaaacc ctacctgtt cctgccccat 840
acctccaccc actcacctcc ctaacctcct tatgtccctc acagtagcaa gatgtaggtg 900
ataggaagga cttcgggtgtg agaattagaa atgatgtaaa tgtttacgca ggagtgtctg 960
gataggagtc gggatggtga gggtagttag atttttgcct cacttgccct gaaagtggta 1020
atagggagaa accaatctga ttaacaatta ctacagactg tcactttgta 1080
ttcctccaac atgtttggta acaagtgcct aatgtatgtt aaaataaaga aggtttttat 1140
acctttccat taaaatatgt cagtgggccc ttccatttta tggagtggaa tgggaaggcc 1200
cttgacagcc aggaaccact tgaagctggc atccactctt gaacagagag tattaaagac 1260
aggattcaca ctgaaaagtg agccacccaa ttgaaatgcg agtaatggag agtaacggag 1320
aatcgttga cctgggaag cagaggttgc agagagccgg gattgcccc 1371

```

&lt;210&gt; 44

&lt;211&gt; 3634

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1911808CB1

&lt;400&gt; 44

```

gcggcggcag tggcgcgcac aggtgattga ctggccagct gcctgaagga gcgccaggtc 60
ctccttgctg gcagggtggc aagcccatg gggcgccggt gcagaccgcg gcggcgctg 120
cggcgggtctg gctcgggagg cgttcctggg gccaaaggcca tggccccgcg gctgcagctg 180
gagaaggcgg cctggcgctg ggccggagacg gtgcggcccg aggaggtgtc gcaggagcat 240
atcgagaccg cttaccgcct ctggctggag ccttgcattc gcggcgtgtg cacagcaaac 300
tgcaaaaggaa atccgaattg cttgggtggg attggtgagc atatttgggt agggaaaata 360
gatgaaaata gttttcataa catcgatgat cccaactgtg agaggagaaa aaagaactca 420
tttgtgggccc tgactaacct tggagccact tgttatgtca acacatttct tcaagtgtgg 480
ttttctcaact tggagcttcg gcaggcactc tacttatgtc caagcacttg tagtgactac 540
atgctgggag acggcatcca agaagaaaaa gattatgagc ctcaaacaaat ttgtgagcat 600
ctccagtaact tgtttgcctt gttgcaaaac agtaataggc gatacattga tccatcagga 660
tttgttaaag ccttgggcct ggacactgga caacagcagg atgctcaaga attttcaaag 720
ctcttttatgt ctctattgga agatactttg tctaaccaaa agaattcaga tgtgcgcaat 780
attgttcaac agcagttctg tggagaatat gcectatgtaa ctgtttgcaa ccagtgtggc 840
agagaagtcta agcttttgtc aaaattttat gagctggagt taaatatcca agggcacaaa 900
cagttaacag attgtatctc ggaatttttg aaggaagaaa aattagaagg agacaatcgc 960
tattttttgcg agaactgtca aagcaaacag aatgcaacaa gaaagattcg acttcttagc 1020
cttctttgca ctctgaactt gcagctaattg cgttttgtct ttgacaggca aactggacat 1080
aagaaaaagc tgaataccta cattggcttc tcagaaaattt tggatatgga gccttatgtg 1140
gaacataaag gtgggtccta cgtgtatgaa ctccagcgag tccatcatac cagaggagtg 1200
agtgtcttatt ctggccacta catcgcccac gtgaaagatc cacagtctgg tgaatggtat 1260
aagtttaatg atgaagacat agaaaagatg gaggggaaga aattacaact agggattgag 1320
gaagatctag cagaaccttc taagtctcag acacgtaaac ccaagtgtgg caaaggaaact 1380
cattgtcttc gaaatgcata tatgttgggt tatagactgc aaactcaaga aaagcccaac 1440
actactgttc aagttccagc ctttcttcaa gagctggtag atcgggataa ttccaaattt 1500
gaggagtggg gtattgaaat ggtgagatg cgtaagcaaa gtgtggataa agggaaaagca 1560
aaacacgaag aggttaagga gctgtaccaa aggttacctg ctggagctga gcccttatgag 1620
tttgtctctc tggaaatggc gcaaaaagtg ttggatgaat caacacctac caacctattt 1680
gataatcacg cttgcctgtg ttcccatgac aagcttcacc cggataaaat atcaattatg 1740
aagaggatat ctgaatatgc agctgacatt ttctatagta gatatggagg aggtccaaga 1800
ctaactgtga aagccctgtg taaggaatgt gtagtagaac gttgtcgcct attgcgtctg 1860
aagaaccaac taaatgaaga ttataaaact gttaataatc tgctgaaagc agcagtaaaag 1920

```

```
ggcagcgaatg gatttttgggt ggggaagtc tctttgcgga gttggcgcca gctagctctt 1980
gaacagctgg atgagcaaga tggatgatga gaacaaagca acggaagat gaacggtagc 2040
accttaataa aagatgaatc aaaggaagaa agaaaagaag aggaggaatt aaattttaat 2100
gaagatatcc tgtgtccaca tggtagtata tgcataatctg aaaaatgaaag aaggcttggt 2160
tctaaagagg cttggagcaa actgcagcag tactttccaa aggtctctga gtttccaagt 2220
tacaagaggt gctgttcaca gtgcaagatt ttagaaagag aaggggaaga aaatgaagcc 2280
ttacataaga tgattgcaaa cgagcaaaaag acttctctcc caaatttgtt ccaggataaa 2340
aacagaccgt gtctcagtaa ctggccagag gatacggatg tctctacat cgtgtctcag 2400
ttctttgtag aagagtggcg gaaatttggt agaaagccta caagatgcag cctgtgttca 2460
tcagttggga acagtgtctt tttgtgtccc cacggggggc tcatgtttac atttgcttcc 2520
atgaccaaaag aagattctaa acttatagct ctcatatggc ccagtgaagt gcaaatgata 2580
caaaagctct ttgtttgtga tcatgtaatt aaaatcacga gaattgaagt gggagatgta 2640
aacccttcag aaacacagta tatttctgag cccaaactct gtccagaatg cagagaaggc 2700
ttattgtgtc agcagcagag ggacctgctg gaatacactc aagccaccat ctatgtccat 2760
aaagtgtgtg ataataaaaa ggtgatgaag gattcggctc cggaactgaa tgtgagttagt 2820
tctgaaacag aggaggacaa ggaagaagct aaaccagatg gagaaaaaga tccagatttt 2880
aatcaaatca tgcattgcat ttcagttgct ccttttgacc agaatttgct aattgatgga 2940
aagattttta gtgatgactg tgccacccta ggcacccttg gcgtcattcc tgaatctgtc 3000
attttattga aggtgatga accaattgca gattatgctg caatggatga tgtcatgcaa 3060
gtttgtatgc cagaagaagg gtttaaagggt actggtcttc ttggacatta atctttgaa 3120
acttgctgac tgctaagaaa tgaccagagg ggaagaggag tttgacatgt tagggcata 3180
aagcaaaagt ttcaaacgtg actgtcccaa atgccttatg tcaataaag cagattgcac 3240
cagacttgaa ggaaatgttt ccaattttat atttaagggg ggtggtgggt gggagggggc 3300
aagtaagac ggaacaagtt tagtagcagt aatagtaaat catgtttaca tatgagatt 3360
atagtcgtgg gaggggaata aagttctgtt atatttcctt gctcgagttc cataccagat 3420
gcgttggtcc ataaaggatt gtatcaagta gatgggacaa cattctgttc tgaacgaaaa 3480
gtaattttag agacataacc tgcttaccac tgctgtctt tgattcatat tctactttca 3540
ataaagcatg aaagtgaaga acttgtaaaa aaaa 3600
3634
```

<210> 45  
<211> 1661  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 1973875CB1

```
<400> 45
ctcgagccga ttcggtctga ggcggactgg tctccggccg ggcaccgtcg cggggccccc 60
tgcccccggc acctgggacc gtgctgggga gtctgccact tccctctctc cctggcccgc 120
caaagttttg gcggaccatc cgtggtgggt gagcgcgcgc ccggggggag atcggggagc 180
gcccagtcgc gggcgccggg agccattgac ccgggacgcc gccgtccgct gagcagccga 240
ccaccctgcc gctccgggtg catggggact ggctgaggag ccagcatggg caactgctgt 300
gggagacagc gccgggagag gccggcagcc ccgggacacc ccgcgaagcg agcaggacgc 360
aatgagcccc tgaagaaaga gcggtctaaag tggaaagagcg actaccccat gactgacggg 420
cagctgcgga gcaaacggga tgagtcttgg gacacagcgc ctgccttcga gggccgcaag 480
gagatctggg atgcccctaa ggctgccgcc tatgtgtctg aagccaacga ccacgagctg 540
gcccaggcca tcttgatgg agccagcatc accctgcctc atggcaccct ctgtgaatgc 600
tacgatgagc tgggcaatcg ctaccagctg cccatctact gcctgtcacc gccggtgaac 660
ctgctgctgg agcacacgga ggaggagagc ctggagcccc ccgagcctcc accagcgtg 720
cgccgtgagt tcccgtgaa ggtgcgcctg tccacgggca aggacgtgag gctcagcgcc 780
agcctgcccc acacagtggg gcagctcaag aggcagctgc acgcccagga gggcatcgag 840
ccatcgtggc agcgggtggtt cttctccggg aagctgtctc cagaccgcac acgggtccag 900
gagaccaaga tccagaaaga ttttgtcatc cagggtcatc tcaaccagcc cccaccacc 960
caggatgatg gggcccacgg acccctggga agaggcccc cctggagcac taggccccca 1020
cctgtctgct gccttccagt gctgtcattt tcttcagggg ccttccccctc ggtgtggctg 1080
gtgggtgagc cgtgaaggga cctgtccttt cagggcacta cgcgccacca gttcccggta 1140
cccaggagc aggcagccac acacgggctt tgcaaccttg tcagagaaaa ggccaacagg 1200
gcccacccc tgctgtctc ccgaagcac ttogagccac aaggggcccc caggagccc 1260
ctggagccca gatctgtcat ctggtgctgc cagctgtgct cactctggtt ttctgtctag 1320
ggtctgaagc agctgctgtc tccctctctt gccccatcc cctggctctc cctggggcac 1380
agtgccactc ccttggaagg gagggaaacca cccgtgagcc ccagggttg ggaagcctga 1440
ggcgggctc tgctctctcc tgccccagc acaattggga gagatgagc ggggtgtgga 1500
cgctgggct gtctggcag ggtctgcaca ggccatgtc ctggctgtaa ccaggcag 1560
gggaggtctg cagcctggct atggcctcca cagcaggtcc ctgtggacag acatatctgc 1620
```



atatttatca ataaagcctt ttgctccttc aaaaaaaaaa a

1661

&lt;210&gt; 46

&lt;211&gt; 1910

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2323917CB1

&lt;400&gt; 46

gtactccgcg	cgcagccctg	gcggtccagc	gcgaatctcc	agaatctggt	ctccagaaac	60
attattctaa	tctgtgtatg	gaaaaatccc	agaaaattaa	tccttttata	ttgcataatac	120
ttcaagaagt	ggatgaagaa	attaaaaagg	ggctagcagc	aggaatcaca	ttaaacattg	180
ctggtaacaa	tcgcttagtg	ccagtagaaa	gagttacagg	tgaagatttt	tggattcttt	240
ccaaaatttt	aaagaattgt	ctgtatatta	atgggtttgga	tgttggtat	aaccttctat	300
gtgatgttgg	tgcatactat	gctgcgaaac	tgcttcagaa	acaacttaac	ctcattttact	360
taaacctcat	gtttaatgat	attgggcccg	aagggtggaga	attgattgct	aaagtgtac	420
ataagaatcg	gactctgaaa	tacctaagaa	tgactggaaa	caaaattgaa	aataaagggtg	480
gaatgttttt	tgtgcaatg	ctgcaaatta	attcatcatt	agagaaatta	gatctgggtg	540
actgtgatct	gggaatgcaa	agtgtgatag	catttgctac	agtactaact	caaaaccaag	600
caattaaggc	aataaaccta	aaccgacct	tactgtacgg	tgaacaggaa	gagctacag	660
tcctatgtagg	cctcatgttg	aaagaaaatc	actgtcttgt	tgactacac	atgtgtaagc	720
atgatataaa	aaacagtggg	atacaacagt	tatgtgatgc	actgtatctg	aacagttagcc	780
tgcgctacct	tgatgtcagc	tgcaacaaaa	taactcatga	tggaatgggtg	tatttggctg	840
atgtactgaa	aagcaacact	accctggaag	taatagatct	ttcctttaac	agaatagaaa	900
atgcaggcgc	aaactatctc	agtgaactc	ttacttcaca	caacaggagt	cttaaagcgt	960
tgtcagtagt	cagcaacaac	atagagggag	aaggacttgt	tgcaacttca	caatcaatga	1020
aaacaaatct	cactttctct	catatctaca	tttggggaaa	caaatttgat	gaggctacgt	1080
gtatagcata	ttcagactta	attcaaatgg	gttgtctaaa	accagacaat	acagatgtgg	1140
agccatttgt	ggtagatgga	cgtgtatatc	ttgcagaagt	ctccaatggc	cttaaaaagc	1200
attattattg	gacatcaact	tatggagaat	cttatgacca	ctcatctaata	gcagggtttg	1260
ctcttgttcc	agtaggtcaa	cagccatgaa	aaagtagctg	taaaataatt	ttcatacact	1320
tgtcttattg	tttcacagag	attaatatct	tatagcctat	tttcttttat	aaatataaac	1380
aagttaacttt	tttcaaatgt	gtaaaagata	atgactgtat	taaactttgt	ataactgtca	1440
aaaaaaaaaa	aaagggggcg	gccccgatta	gtgagctcgt	ccccgcggtg	atttactccc	1500
ggggccccgt	cccttggggg	gggttaacca	gtctttccgc	caaggggtgg	gtcctgtata	1560
tacaggcgct	ttggaaagag	aagtcgtgct	gtccctgggc	tcttcttctg	gtcggaggggc	1620
atcttactcg	tctgtccccg	ctggtgtaac	atagtaccgc	cacgcgtgtg	gtcctgtgga	1680
tgttggaaat	aagtgccttg	tggagtctcg	ctcgggtgtg	gacgtaggta	gatcctcgca	1740
gacgtttctca	tatcggtctg	gcgttggctg	tggccccgct	tcccgtggtt	gtgttgcga	1800
ggtagtggtg	ttgcacctgt	atcttatattg	gtccctccgc	cgcgcgggtg	ggaggcggtg	1860
ttggtttagt	agctcgtggc	ggttgttgtg	ttcctcaacc	cacacacaat		1910

&lt;210&gt; 47

&lt;211&gt; 2162

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2754960CB1

&lt;400&gt; 47

gacagcgcaa	ccgcactcca	gcctggacga	cagaatgaga	ctccgtctca	aaaaaaaaaa	60
aaaaaaaaaa	aagcgagggc	tggaagtctg	gagcaggtgc	gcggtctcaa	cggcagccgc	120
gggaagctcg	ggccggcagg	gtttcccgcc	acgctggcgc	ccagctcccc	gcgcggaggc	180
cgtgtgaagt	ttcgttttcc	attcagtggg	aaacgaaagc	tgggcggggg	gccacgagcg	240
cggggccaga	ccaaggcggg	cccgagcgcg	aacttcgggc	ccagctcggt	ccccggtcca	300
gtccccgacg	ggaactcagc	agcggaggct	ggacgcttgc	atggcgcttg	agagattcca	360
tctgtccttg	ctcacataag	cgttctctgg	aagtgaagtc	gtgctgtcct	gaacgcgggc	420
caggcagctg	cggcctgggg	gttttgaggt	gatcacgaat	gagcaaggcg	tttgggctcc	480
tgaggcaaat	ctgtcagtc	atcctggctg	agtcctcgca	gtccccggca	gatcttgaag	540
aaaagaagga	agaagacagc	aacatgaaga	gagagcagcc	cagagagcgt	cccagggcct	600
gggactaccc	tcatggcctg	gttggtttac	acaacattgg	acagacctgc	tgcttcaact	660
ccttgattca	ggtgttcgta	atgaatgtgg	acttcaccag	gatattgaag	aggatcacgg	720

```

tgcccagggg agctgacgag cagaggagaa gcgtcccttt ccagatgctt ctgctgctgg 780
agaagatgca ggacagccgg cagaaagcag tgcggccctt ggagctggcc tactgcctgc 840
agaagtgcaa cgtgcccttg tttgtccaac atgatgctgc ccaactgtac ctcaaactct 900
ggaacctgat taaggaccag atcactgatg tgcacttggt ggagagactg caggccctgt 960
atacgatccg ggtgaaggac tccttgattt gcgttgactg tgccatggag agtagcagaa 1020
acagcagcat gctcaccttc ccactttctc tttttgatgt ggactcaaag cccctgaaga 1080
cactggagga cgccctgcac tgctttcttc agccagggga gttatcaagc aaaagcaagt 1140
gcttctgtga gaactgtggg aagaagacc gtgggaaaca ggtcttgaag ctgacccatt 1200
tgccccagac cctgacaatc cacctcatgc gattctccat caggaattca cagacgagaa 1260
agatctgcca ctccctgtac ttccccaga gcttggattt cagccagatc ctccaatga 1320
agcgagagtc ttgtgatgct gaggagcagt ctggagggca gtatgagctt tttgctgtga 1380
ttgcgcacgt gggaatggca gactccggtc attactgtgt ctacatccgg aatgctgtgg 1440
atggaaaatg gttctgtctc aatgactcca atatttgctt ggtgtcctgg gaagacatcc 1500
agtgtacata cggaaatcct aactaccact ggcaggaaac tgcatactct ctggtttaca 1560
tgaagatgga gtgctaattg aaatgcccaa aaccttcaga gattgacacg ctgtcatttt 1620
ccatttccgt tcttgatctc acggagtctt ctaagagatt ttgcaatgag gagaagcatt 1680
gttttcaaac tatataactg agccttattt ataattaggg atattatcaa aatatgtaac 1740
catgagggcc ctccaggtct gatcagtcag atggatgct ttcaccagca gaccggcca 1800
tgtggctgct cgttcctggg tgctcgtgc tggccaagac attagccctt tagttatgag 1860
cctgtgggaa cttcaggggt tcccagtggg gagagcagtg gcagtgggag gcatctgggg 1920
gccaaaggtc agtggcaggg ggtatttcag tattatacaa ctgctgtgac cagacttgta 1980
tactggctga atatcagtgc tgtttgtaat ttttcacttt gagaaccaac attaatcca 2040
tatgaaatcaa gtgttttgta actgctattc atttattcag caaatattta ttgactcatc 2100
cttctccata agatagtgtg ataaacacag tcatgaataa agttattttc cacaaaaaaa 2160
aa 2162

```

&lt;210&gt; 48

&lt;211&gt; 578

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3092341CB1

&lt;400&gt; 48

```

gaaaggccca cctgtgtcct ggttgagggt ctccagggtt ctttggggcc cgaggaattt 60
cagacctgaa caatcaggtg gacaacttct tgtgactttg ggtattctga aattactcat 120
cttgactcta ctgtgccttg atgtagctgt gtgttttggt tgtatcctgt tcatggccaa 180
tgggtggcaga gtctacatag aactatgctt cgtggtgttc tggggaaaac ctttcgactt 240
gttggctata ctattcaata tggctgtata gctcattgtg cttttgaata cgttgggtgt 300
gttgtcatgg tgccaatggg tcatgttttg ttagaagggt acaatctaca gaattctaca 360
gattccaggt gctatggacc tattccatat ggactaataa gaggacgaat cttctttaag 420
atttggcttc tgagtgattt tggattttta cgtgccagcc ctaatggcca cagattttct 480
gatgattagt aagcatttat tcttttgact tgattattgt ctctttttca tgtgaattta 540
ttactccctg tgaaccctgt tacttaccaa taaactat 578

```

&lt;210&gt; 49

&lt;211&gt; 1300

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3658034CB1

&lt;400&gt; 49

```

agggaaatccc aattatgagg cggctaagca ggcaaagaaa tagcttgggt ggttttgcatt 60
ctactcagga taccactgtg gctttaaagg ctgtctgaat ttgcagccct aatgaatata 120
gaaaggacaa atatccaagt gaccgtgacg gggcctagct caccaagtc tgtaaagttt 180
cgtattgaca cacacaaccg cttactcctt cagacagcag agcttgcctg ggtacagcca 240
acggcagtta atatttccgc aaatggtttt ggatttgcta tttgtcagct caatgttgta 300
tataatgtga aggtctctgg gtcttctaga agacgaagat ctatccaaaa tcaagaagcc 360
tttgatttag atgttgctgt aaaagaaaat aaagatgatc tcaatcatgt ggatttgaat 420
gtgtgtacaa gcttttcggg cccgggtagg agtggcatgg ctcttatgga agttaacct 480
ttaagtggct ttatgggtgc ttcagaagca atttctctga gcgagacagt gaagaaagt 540
gaatatgatc atggaaaact caacctctat ttagattctg taaatgaaac ccagttttgt 600

```

```

gttaatatcc ctgctgtgag aaacttttaa gtttcaaata cccaagatgc ttcagtgtcc 660
atagtggatt actatgagcc aaggagacag gcggtgagaa gttacaactc tgaagtgaag 720
ctgtcctcct gtgacctttg cagtgatgtc cagggctgcc gtccttgtga ggatggagct 780
tcaggetccc atcatcactc ttcagtcatt tttatTTTTt gtttcaagct tctgtacttt 840
atggaacttt ggctgtgatt tttttttaa ggactctgtg taacactaac atttccagta 900
gtcacatgtg attgttttgt tttcgtagaa gaatactgct tctatTTTTg aaaaagagtt 960
ttttttcttt ctatggggtt gcagggatgg tgtacaacag gtcctagcat gtatagctgc 1020
atagatttct tcacctgatc tttgtgtgga agatcagaat gaatgcagtt gtgtgtctat 1080
attttccctt ctcaaaatct tttagaattt ttttggaggt gtttgttttc tccagaataa 1140
aggattactt ttagaatagg tattctcctc attttgtgaa agaaatgaac ctagattctt 1200
aagcattatt acacatccat gtttgcttaa agatggattt ccctgggaat gggagaaaaa 1260
agccagcagg aggagcttca tctgttccct tcccacctcc 1300

```

<210> 50  
 <211> 2241  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3883861CB1

```

<400> 50
gctccccgag ccgggctgca ccggaggcgg cgagatggtc gcgcgcgtcg gcctcctgct 60
gcgcgccttg cagctgctac tgtggggcca cctggacgcc cagcccgagg agcgcgagg 120
ccaggagctg cgcaaggagg cggaggcatt cctagagaag tacggatacc tcaatgaaca 180
gggtcccaaaa gctcccacct ccactcgatt cagcgatgcc atcagagcgt ttcagtgggt 240
gtcccgctca cctgtcagcg gcgtgttggg ccgcgccacc ctgcgccaga tgactcgtcc 300
ccgctgcggg gttacagata ccaacagtta tgcggcctgg gctgagagga tcagtactt 360
gtttgtctaga caccggacca aaatgaggcg taagaaacgc tttgcaaagc aaggtaacaa 420
atggtacaag cagcacctct cctaccgcct gggtgaactg cctgagcacc tgccggagcc 480
ggcagttcgg ggcgcgtg gcgcgccttg ccagttgtgg agcaacgtct cagcgtgga 540
gtttctgggag gccccagcca caggccccgc tgacatccgg ctacacctct tccaagggga 600
ccacaacgat gggctgggca atgcctttga tggcccaggg ggccgctgg cgcacgctt 660
cctgccccgc cgcggcgaag cgcacttcga ccaagatgag cgttggtccc tgagccgcc 720
ccgcgggcgc aacctgttcg tgggtgctgg gcacgagatc ggtcacacc ttggcctcac 780
ccactgcgcc gcgcgcgcgc cgtcatggc gccctactac aagaggctgg gccgcgacgc 840
gctgctcagc tgggacgacg tgcctggcgt gcagagcctg tatgggaagc ccctaggggg 900
ctcagtgccc gtccagctcc caggaaagct gttcaactgac tttgagacct gggactccta 960
cagcccccaa ggaaggcgcc ctgaaacgca tactgccact ctctcttcca 1020
tgccatcact gatggcaacg tctcagagcc ccgtccactg caggaaagat atttctggga 1080
gggtggcagct gatggcaacg attgaggctg cggcagtgct attgaatgat ggagatttct gggtcgggct 1140
gccccccaac tgcctggaggt tccggggccc gccatcctga cgcgcctctc ttcttccctc ctctgcgccg 1260
ccgggcaggg ggccctgccc cccgctacta cgtgctggcc cgagggggac tgcaagtga 1380
cctcactctc ccccgaaagt tgcaggactg gggaggcatc cctgaggagg tcagcggcgc 1440
gcccacttac cccgatggct ccatcatctt ctcccgagat gaccgctact ggcgcctcga 1500
cctgcccagg ccaggccaaa ctgcaggcaa ccacctcggg ccctggggcc accgagctgc cctggatggg 1560
ctgctggcat gccaaactcg ggagcgcctt gttctgaagg cacctcctca cctcagaaac 1620
tggttggtgct ctacgggcaa aatcatgttc cccacccccg gggcagaacc cctcttagaa 1680
gcctctgagt ccctctgcag aagaccgggc agcaaagcct ccactctggg gtctgtctgc 1740
ctttgttctt tgaagaatgc agcattgtct ttgtctgtcc ccaccaatg gaggtggggg 1800
tgggtcaaat cttaggaaaa gcaaaaaagg gtcccagatc ccttggccct tctctccag 1860
gactctctat ctccccaggc ctttgtttct teggttaaag gtacagttcc tttcaagagg 1920
taacagcact gggatccaag caggggggatg aaaaactcag cagagaaatt cgagaccatt 1980
ttgcaagact gtgcctttct cctcaggacc ccttgctca gttcttgaaa aacggtgtca 2040
tatttagtca gaggccccac cccaggaag catggatggg gatgaaggca caggcgtctc 2100
caacctcaga ggccctttgt ggggtcagga cacagagtgg gagggagact gatgcaggcc 2160
taccagtccc tggctttttg tctggggctg gaataaagag gtgccttcag ctggtggggc 2220
gagaggcagg aaaaaaaaaa a 2241

```

<210> 51  
 <211> 1860  
 <212> DNA  
 <213> Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4993873CB1

&lt;400&gt; 51

```
agaattgacc tgacaaatat tgaccaggtt gctgtgatct tcaaacacca ttttctgtg 60
ggtagagggtg atgcagtgtt gaagacctgg gcccttgcct agtgccctttg ctctagaatg 120
ggtccagctt ggcttttggct actgggaaca gggatcctgg cctctgtcca ctgtcagccc 180
cttcttgccc atggagataa aagtctgcag gggcctcaac ccccaggca tcagctctca 240
gagccagccc ccgcctacca cagaatcaca cccaccatta ccaattttgc tttgcgtttg 300
tataaagagc tggcagcaga cgcctccgga aacatcttct tctcgccagt gagcatctcc 360
accacctgtg cctgtctctc tcttggggcc caagctaaca cctcagctct gatcctggag 420
ggcctgggat tcaacctcac agaaacctct gaagccgaca tccaccaggg ctccgggagc 480
ctcctccaca ccttggccct gccagcccc aaactcgaa taaaagtagg aaactccctg 540
ttcctagaca agcgactaaa gcctcggcag cactatttgg acagcatcaa ggagctttat 600
ggagcttttg ctttttctgc caacttcaca gattctgtta caactgggag gcagattaat 660
gactatttga gaaggcaaac atacgggcaa gtctgtggact gcctcccgga gttcagccag 720
gacacgttca tgggtcttgc caattacatc ttcttcaaag ccaagtggaa gcacctttc 780
agtcgtctac agacccagaa gcaggcaagt ttctttgtgg atgagaggac ttctctccag 840
gtccccatga tgcacaaaaa ggaaatgcac agattcctct atgaccagga tttggcttgc 900
actgtcctcc agatagaata cagaggaaat gccttggcgc tgctggctct cctgacccg 960
gggaaaaatga agcagggtgga ggctgctctg cagccacaga ccttgagaaa atggggccaa 1020
ttgctcctgc ccagtctgtt ggatttgcac ttgccaaagt tttcaatttc tggacatat 1080
aacctggaag acatacttcc ccaaattggt ctcaccaaca tactcaactt agaagctgac 1140
ttctcaggag tcaactgggca gctcaacaaa accatctcca aggtgtcaca caaggcgatg 1200
gtggacatga gtgagaaggg gaccgagccc gggcctgctt caggcctctc ctcccagccc 1260
ccatctctga acaccatgtc agaccacatc gcccacttca acaggccttt cctcttgcctc 1320
ctttgggagg tcaccaccca gagcttactc ttctgggaa aagttgtcaa cccagttgca 1380
gggtaaccat ggtgggaggc caggagtatt cttatctcat cctggacca acagataggc 1440
cagaaccagc ctgcctctgt gggctgtctt gtggttcagt taatcagtg gccaagattc 1500
taataaagt gaccttgggt tctgtgaaaa aaaaaaaaag aacaaccaac gcataaagtt 1560
gcggcgtaaa ggatgctgaa agcaaatacc ccggtttgcc cgtccaaatt taacataagg 1620
ggggggccgg cagtacagaa ggtccctcag agtggtgttg atggcgtcg actgattggc 1680
ggggggggcg agagagttt tacgcattct tgcacgtgtt agcaccggg gggggggggc 1740
ccataggacc cccgagctg gaggttgggt ttgcgcgggc actcttgggg ggtgtgttat 1800
gaccgacttt tagtggcaca aactgtgggg cttctcatat gtgtgcgcgc gcttctgttt 1860
```

&lt;210&gt; 52

&lt;211&gt; 550

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5208004CB1

&lt;400&gt; 52

```
ggcgggtggc tggaggacac ggtgtaggcc ttgccgtgt ctgggttctt cgccactgca 60
ggagcaggac tgtttctgga aacactgggc tgctggtgac ccgtcaaact tctcccacaa 120
attctgaatc cgagaaagtg aaggaaggat ggtggggaag tgaggaggca ggagcagagg 180
ccacagggac cgaccagaga tgcggtggag acagaggagc ttctttctca ggctgtttct 240
gggaagcctg agaggcggcc agcaccacc ctcgctcact ctccctcag cctctctgct 300
tcccttctca accctttctc tctctctgc ctcttccct tcttgcgtgc ttgttagtcc 360
ctgtcccaaa actcctggct cctttgttct gctgcctgg ccccccacca ggaggaggtc 420
tcaggcacca tccccccac gagggatcca cacaacaggt tcatgctggg gctggggggg 480
ccccgctggg ttcttgatgc cttgtgcaca ggaagtgtc gcagtcattt ttggactctc 540
ctgaatgtgt
```

&lt;210&gt; 53

&lt;211&gt; 2597

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5267783CB1

```

<400> 53
gacctggttgg gccgcgtcct tetgctgccc cgcgcggtg caggcctcgc cgcgagccgc 60
agggtgtcctg gagtctggcc caggacctgg cccacagga gtcccagcag gggtagctcc 120
tcccgggaca aggaccgaag tgcgacggtc agtagttcag tgcccagcgc tgctggaggg 180
aaaggaagcc atccttcac taccacccag aggggtcccca accgcctgat ccacgagaag 240
tcaccatacc tcctacaaca tgccataaat cctgtggact ggtaccctg gggagaggaa 300
gccttcgaca aggccaggaa ggaaaacaag ccgattttcc tctcagtcgg gtactccacc 360
tgccactggg gccacatgat ggaagaggag tcttccaga atgaggagat tggccgcctg 420
ctcagtgagg actttgtgag tgtgaaggta gaccgtgagg agcggcctga cgtggacaag 480
gtgtacatga cgttcgtgca ggccaccagc agcggcgggg gctggcccat gaatgtgtgg 540
ctgactccca acctccagcc ctttgtcggg ggcacctatt tccctcctga ggtatggctg 600
acctcgagtc gcttcgcac agtgttgtg agaatacgag aacagtggaa acagaacaag 660
aacacctgc tagaaaatag ccagcgtgtc accactgccc tgctggcccc atcagagatc 720
agcgtgggtg accgccagct gccgcctct gccgccaccg tgaacaatcg ctgcttccag 780
cagctggatg agggctatga tgaggaaata ggtggcttcg ctgaggcccc caagtttccc 840
acgcgggtga tcctgagctt cctgttctcc tactggctca gccatcgact gactcaggat 900
ggctctcggg cccagcagat ggcccttgc atacctgaaa tgatggctaa cgggggcac 960
cgggaccatg tggggcaggg ctttccacc tactccacag accgccagt gacgtccct 1020
cactttgaga agatgctcta tgaccaggca cagctcgtc tggcctatt gcaggcctc 1080
cagctctctg gtgatgaatt ctactctgac gtggccaaag gcacctgca gtactgggt 1140
cggagcctga gccaccggtc cggaggcttc tatagcgag aagatgcaga ctgccccca 1200
gagcggggcc agcggcccaa agagggcgcc tactatgtgt ggacgggcaa agaggttcag 1260
cagctcctcc cggagcctgt gttgggtgcc accgagccgc tgacctcagg ccagctccc 1320
atgaagcact acggcctcac agaggctggt aacatcagcc ccagtcagg cccaagggg 1380
gagctgcagg gctgaccgtc gctgaccgtc cgggtactgc tggagctgac tgctgcccc 1440
tttggcttgg atgtggaggc cgtgcggacc ttgctcaatt cagggctgga gaagctctc 1500
caggccccga agcatcggcc caagccgcac ctggacagca agatgctggc tgcctggaat 1560
ggcttgatgg tgtcaggcta tgctgtgact ggggctgtcc tgggccaaga cagctgac 1620
aactatgcca ccaatggtgc caagtctctg aagcggcaca tgtttgatg ggccagtggc 1680
cgctgatgc ggacctgcta caccggccct ggggggactg tggagcacag caaccaccc 1740
tgctggggct tcctggagga ctacgccttc gtggtgcggg gctctgcggc tgcaggacac acaggacag 1800
gctcaccagg agagtgcgtg gtcgagtggt tggcggctac ttctgcagt aggctgagc cagcgccaa 1860
ctcttttggg actcccagg gtctgaagga cgaccaggat ggagcagagc ccagcgccaa 1920
ctgccccctg gccacaacc tgctccggct gcatggcttc acggggccaca aggactggat ggacaagtgt 2040
gcccacaacc tgctccggct tgaccgcctt tcccgagcgc atgcgtcgtg tcccgggtggc gttgcccag 2100
atggtccgcg cccctcagc ccagcagcag accctcaagc agatcgatg ctgtggagac 2160
cgtcaggcca aggaaccaa ggccctggtg cagtgcgtcc actctgtcta cattcctaac 2220
aaggtgctga ttctggctga tggggacccc tegagcttcc tgtcccgcca gctgccttc 2280
ctgagtaccc tccgacggtt ggaagaccag gccactgcat atgtgtgtga gaatcaagcc 2340
tgctcagtcg ccatcactga tccctgcgaa ttacgaaaac tactacatcc atgactgc 2400
caacccccct ggggtggggc agaagggtga gcatcccaac tgactagaga ctcaggccct 2460
gcagggccct atagaacctg tggccatccc tgagcaccct gccaccagg gacctcgcc 2520
atactactg ccccccttgg gcacccact accctagaat aaacttaaca gtgtcccg 2580
gtaaaaaaaa aaaaaaa 2597

```

```

<210> 54
<211> 2385
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 5583922CB1

```

```

<400> 54
gaccgacccg ccacccgcgc tagcccgccg gcccctggca ctcaatcccc gccatgtggg 60
ggctcctgct cgccttgccc gccttcgcgc cggcgtcgg cccggtctg ggggcccga 120
ggaactcggg gctgggccc cgcagcccg ggaccaccaa ggtcccagg tcgaccccg 180
ccctgcatag cagcccgcca cagcccgccg cgagacagc taacgggac tcagaacagc 240
atgtccggat tcgagtcac aagaagaaaa aggtcattat gaagaagcgg aagaagctaa 300
ctctaactcg cccaccccca ctggtgactg cggggccctt tgtgacccc actccagcag 360
ggacctcga ccccgctgag aaacaagaaa caggctgtcc tcccttgggt ctggagccc 420
tgcgagtctc agatagccgg cttgaggcat ccagcagcca gtccttgggt ctggagccc 480
accgaggacg gctcaacatt cagtcaggcc tggaggacgg cgatctatat gatggagcct 540
ggtgtgctga ggagcaggac gccgatccat ggtttcagg ggacgctggg caccacccc 600
gcttctcggg tgttatcaca cagggcagta actctgtctg gaggtatgac tgggtcacat 660

```

catacaaggt	ccagttcagc	aatgacagtc	ggacctggtg	gggaagtagg	aaccacagca	720
gtgggatgga	cgcagtattt	cctgccaatt	cagacccaga	aactccagtg	ctgaacctcc	780
tgccggagcc	ccaggtggcc	cgcttcattc	gcctgctgcc	ccagacctgg	ctccaggagg	840
gcgcgccttg	cctccgggca	gagatcctgg	cctgcccagt	ctcagacccc	aatgacctat	900
tccttgaggc	ccctgcgtcg	ggatcctctg	accctctaga	ctttcagcat	cacaattaca	960
aggccatgag	gaagctgatg	aagcaggtaac	aagagcaatg	ccccaacatc	acccgcattc	1020
acagcattgg	gaagagctac	cagggcctga	agctgtatgt	gatggaaatg	tcggacaagc	1080
ctggggagca	tgagctgggg	gagcctgagg	tgcgctacgt	ggctggcatg	catgggaacg	1140
aggccctggg	gcgggagttg	cttctgctcc	tgatgcagtt	cctgtgccat	gagttcctgc	1200
gagggaaacc	acgggtgacc	cgctgctctc	ctgagatgcy	cattcacctg	ctgccctcca	1260
tgaacctga	tggtctatgag	atcgctacc	accggggttc	agagctgggtg	ggctgggccc	1320
agggccgctg	gaacaaccag	agcatcgatc	ttaaccataa	ttttgctgac	ctcaacacac	1380
cactgtggga	agcacaggac	gatgggaagg	tgccccacat	cgcccccaac	catcacctgc	1440
cattgcccac	ttactacacc	ctgcccgaatg	ccaccgtggc	tcctgaaacg	cgggcagtaa	1500
tcaagtggat	gaagcggatc	ccctttgtgc	taagtgccaa	cctccacggg	ggtgagctcg	1560
tggtgtccta	cccattcgac	ctgtactcgca	ccccgtgggc	tgcccgcgag	ctcacgcca	1620
caccagatga	tgctgtgttt	cgctgggtca	gcactgtcta	tgctggcagt	aatctggcca	1680
tgaggacac	cagccgccga	ccctgccaca	gccaggactt	ctccgtgcac	ggcaacatca	1740
tcaacggggc	tgactggcac	acgggtcccc	ggagcatgaa	tgacttcagc	tacctacaca	1800
ccaactgctt	tgaggctcact	gtggagctgt	cctgtgacaa	gttccctcac	gagaatgaat	1860
tgcccagga	gtgggagaa	aacaaagacg	ccctcctcac	ctacctggag	caggtgcgca	1920
tgggcattgc	aggagtgggtg	agggacaagg	acacggagct	tgggattgct	gacgtgtca	1980
ttgccgtgga	tgggattaac	catgacgtga	ccacggcgtg	gggcggggat	tattggcgtc	2040
tgctgacccc	aggggactac	atggtgactg	ccagtgccga	gggctaccat	tcagtgcac	2100
ggaactgtcg	ggtcaccttt	gaagaggggc	ccttcccctg	caatttcgtg	ctcaccaaga	2160
ctcccaaaca	gaggctgcgc	gagctgctgg	cagctggggc	caaggtgccc	ccggaccttc	2220
gcaggcgctt	ggagcggcta	aggggacaga	aggattgata	cctgcggttt	aagagcccta	2280
gggcaggctg	gacctgtcaa	gacgggaagg	ggaagagtag	agagggaggg	acaaagtgag	2340
gaaaaggctg	tcattaaagc	taccgggcac	cttaaaaaaa	aaaaa		2385

## SEQUENCE LISTING

&lt;110&gt; INCYTE GENOMICS, INC.

YUE, Henry  
 TANG, V. Tom  
 BANDMAN, Olga  
 LAL, Preeti  
 BAUGHN, Mariah R.  
 AZIMZAI, Valda  
 LU, Dyang Aina M.  
 YANG, Junming

&lt;120&gt; PROTEASES AND PROTEASE INHIBITORS

&lt;130&gt; PF-0727 PCT

&lt;140&gt; To Be Assigned

&lt;141&gt; Herewith

&lt;150&gt; 60/147,986; 60/160,807

&lt;151&gt; 1999-08-09; 1999-10-21

&lt;160&gt; 54

&lt;170&gt; PERL Program

&lt;210&gt; 1

&lt;211&gt; 444

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 088718CD1

&lt;400&gt; 1

```

Met Lys Val Val Pro Ser Leu Leu Leu Ser Val Leu Leu Ala Gln
  1      5      10      15
Val Trp Leu Val Pro Gly Leu Ala Pro Ser Pro Gln Ser Pro Glu
  20      25      30
Thr Pro Ala Pro Gln Asn Gln Thr Ser Arg Val Val Gln Ala Pro
  35      40      45
Arg Glu Glu Glu Glu Asp Glu Gln Glu Ala Ser Glu Glu Lys Ala
  50      55      60
Gly Glu Glu Glu Lys Ala Trp Leu Met Ala Ser Arg Gln Gln Leu
  65      70      75
Ala Lys Glu Thr Ser Asn Phe Gly Phe Ser Leu Leu Arg Lys Ile
  80      85      90
Ser Met Arg His Asp Gly Asn Met Val Phe Ser Pro Phe Gly Met
  95     100     105
Ser Leu Ala Met Thr Gly Leu Met Leu Gly Ala Thr Gly Pro Thr
  110     115     120
Glu Thr Gln Ile Lys Arg Gly Leu His Leu Gln Ala Leu Lys Pro
  125     130     135
Thr Lys Pro Gly Leu Leu Pro Ser Leu Phe Lys Gly Leu Arg Glu
  140     145     150
Thr Leu Ser Arg Asn Leu Glu Leu Gly Leu Ser Gln Gly Ser Phe
  155     160     165
Ala Phe Ile His Lys Asp Phe Asp Val Lys Glu Thr Phe Phe Asn
  170     175     180
Leu Ser Lys Arg Tyr Phe Asp Thr Glu Cys Val Pro Met Asn Phe
  185     190     195
Arg Asn Ala Ser Gln Ala Lys Arg Leu Met Asn His Tyr Ile Asn
  200     205     210
Lys Glu Thr Arg Gly Lys Ile Pro Lys Leu Phe Asp Glu Ile Asn
  215     220     225
Pro Glu Thr Lys Leu Ile Leu Val Asp Tyr Ile Leu Phe Lys Gly

```

Lys	Trp	Leu	Thr	Pro	Phe	Asp	Pro	Val	Phe	Thr	Glu	Val	Asp	Thr	230	235	240
															245	250	255
Phe	His	Leu	Asp	Lys	Tyr	Lys	Thr	Ile	Lys	Val	Pro	Met	Met	Tyr	260	265	270
Gly	Ala	Gly	Lys	Phe	Ala	Ser	Thr	Phe	Asp	Lys	Asn	Phe	Arg	Cys	275	280	285
His	Val	Leu	Lys	Leu	Pro	Tyr	Gln	Gly	Asn	Ala	Thr	Met	Leu	Val	290	295	300
Val	Leu	Met	Glu	Lys	Met	Gly	Asp	His	Leu	Ala	Leu	Glu	Asp	Tyr	305	310	315
Leu	Thr	Thr	Asp	Leu	Val	Glu	Thr	Trp	Leu	Arg	Asn	Met	Lys	Thr	320	325	330
Arg	Asn	Met	Glu	Val	Phe	Phe	Pro	Lys	Phe	Lys	Leu	Asp	Gln	Lys	335	340	345
Tyr	Glu	Met	His	Glu	Leu	Leu	Arg	Gln	Met	Gly	Ile	Arg	Arg	Ile	350	355	360
Phe	Ser	Pro	Phe	Ala	Asp	Leu	Ser	Glu	Leu	Ser	Ala	Thr	Gly	Arg	365	370	375
Asn	Leu	Gln	Val	Ser	Arg	Val	Leu	Gln	Arg	Thr	Val	Ile	Glu	Val	380	385	390
Asp	Glu	Arg	Gly	Thr	Glu	Ala	Val	Ala	Gly	Ile	Leu	Ser	Glu	Ile	395	400	405
Thr	Ala	Tyr	Ser	Met	Pro	Pro	Val	Ile	Lys	Val	Asp	Arg	Pro	Phe	410	415	420
His	Phe	Met	Ile	Tyr	Glu	Glu	Thr	Ser	Gly	Met	Leu	Leu	Phe	Leu	425	430	435
Gly	Arg	Val	Val	Asn	Pro	Thr	Leu	Leu							440		

&lt;210&gt; 2

&lt;211&gt; 565

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 114551CD1

&lt;400&gt; 2

Met	Ser	Gly	Arg	Ser	Lys	Arg	Glu	Ser	Arg	Gly	Ser	Thr	Arg	Gly	1	5	10	15
Lys	Arg	Glu	Ser	Glu	Ser	Arg	Gly	Ser	Ser	Gly	Arg	Val	Lys	Arg	20	25	30	35
Glu	Arg	Asp	Arg	Glu	Arg	Glu	Pro	Glu	Ala	Ala	Ser	Ser	Arg	Gly	35	40	45	50
Ser	Pro	Val	Arg	Val	Lys	Arg	Glu	Phe	Glu	Pro	Ala	Ser	Ala	Arg	50	55	60	65
Glu	Ala	Pro	Ala	Ser	Val	Val	Pro	Phe	Val	Arg	Val	Lys	Arg	Glu	65	70	75	80
Arg	Glu	Val	Asp	Glu	Asp	Ser	Glu	Pro	Glu	Arg	Glu	Val	Arg	Ala	80	85	90	95
Lys	Asn	Gly	Arg	Val	Asp	Ser	Glu	Asp	Arg	Arg	Ser	Arg	His	Cys	95	100	105	110
Pro	Tyr	Leu	Asp	Thr	Ile	Asn	Arg	Ser	Val	Leu	Asp	Phe	Asp	Phe	110	115	120	125
Glu	Lys	Leu	Cys	Ser	Ile	Ser	Leu	Ser	His	Ile	Asn	Ala	Tyr	Ala	125	130	135	140
Cys	Leu	Val	Cys	Gly	Lys	Tyr	Phe	Gln	Gly	Arg	Gly	Leu	Lys	Ser	140	145	150	155
His	Ala	Tyr	Ile	His	Ser	Val	Gln	Phe	Ser	His	His	Val	Phe	Leu	155	160	165	170
Asn	Leu	His	Thr	Lys	Phe	Tyr	Cys	Leu	Pro	Asp	Asn	Tyr	Glu		170	175	180	185
Ile	Ile	Asp	Ser	Ser	Leu	Glu	Asp	Ile	Thr	Tyr	Val	Leu	Lys	Pro	185	190	195	200
Thr	Phe	Thr	Lys	Gln	Gln	Ile	Ala	Asn	Leu	Asp	Lys	Gln	Ala	Lys	200	205	210	



```

Leu Ser Arg Ala Tyr Asp Gly Thr Thr Tyr Leu Pro Gly Ile Val
215 220
Gly Leu Asn Asn Ile Lys Ala Asn Asp Tyr Ala Asn Ala Val Leu
230 235
Gln Ala Leu Ser Asn Val Pro Pro Leu Arg Asn Tyr Phe Leu Glu
245 250
Glu Asp Asn Tyr Lys Asn Ile Lys Arg Pro Pro Gly Asp Ile Met
260 265
Phe Leu Leu Val Gln Arg Phe Gly Glu Leu Met Arg Lys Leu Trp
275 280
Asn Pro Arg Asn Phe Lys Ala His Val Ser Pro His Glu Met Leu
290 295
Gln Ala Val Val Leu Cys Ser Lys Lys Thr Phe Gln Ile Thr Lys
305 310
Gln Gly Asp Gly Val Asp Phe Leu Ser Trp Phe Leu Asn Ala Leu
320 325
His Ser Ala Leu Gly Gly Thr Lys Lys Lys Lys Lys Thr Ile Val
335 340
Thr Asp Val Phe Gln Gly Ser Met Arg Ile Phe Thr Lys Lys Leu
350 355
Pro His Pro Asp Leu Pro Ala Glu Glu Lys Glu Gln Leu Leu His
365 370
Asn Asp Glu Tyr Gln Glu Thr Met Val Glu Ser Thr Phe Met Tyr
380 385
Leu Thr Leu Asp Leu Pro Thr Ala Pro Leu Tyr Lys Asp Glu Lys
395 400
Glu Gln Leu Ile Ile Pro Gln Val Pro Leu Phe Asn Ile Leu Ala
410 415
Lys Phe Asn Gly Ile Thr Glu Lys Glu Tyr Lys Thr Tyr Lys Glu
425 430
Asn Phe Leu Lys Arg Phe Gln Leu Thr Lys Leu Pro Pro Tyr Leu
440 445
Ile Phe Cys Ile Lys Arg Phe Thr Lys Asn Asn Phe Phe Val Glu
455 460
Lys Asn Pro Thr Ile Val Asn Phe Pro Ile Thr Asn Val Asp Leu
470 475
Arg Glu Tyr Leu Ser Glu Glu Val Gln Ala Val His Lys Asn Thr
485 490
Thr Tyr Asp Leu Ile Ala Asn Ile Val His Asp Gly Lys Pro Ser
500 505
Glu Gly Ser Tyr Arg Ile His Val Leu His His Gly Thr Gly Lys
515 520
Trp Tyr Glu Leu Gln Asp Leu Gln Val Thr Asp Ile Leu Pro Gln
530 535
Met Ile Thr Leu Ser Glu Ala Tyr Ile Gln Ile Trp Lys Arg Arg
545 550
Asp Asn Asp Glu Thr Asn Gln Gln Gly Ala
560 565

```

&lt;210&gt; 3

&lt;211&gt; 589

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1261376CD1

&lt;400&gt; 3

```

Met Ala Glu Ser Gly Glu Ser Gly Gly Pro Pro Gly Ser Gln Asp
1 5 10 15
Ser Ala Ala Gly Ala Glu Gly Ala Gly Ala Pro Ala Ala Ala Ala
20 25 30
Ser Ala Asp Ala Lys Ile Met Lys Val Thr Val Lys Thr Pro Lys
35 40 45
Glu Lys Glu Glu Phe Ala Val Pro Glu Asn Ser Ser Val Gln Gln
50 55 60
Phe Lys Glu Glu Ile Ser Lys Arg Phe Lys Ser His Thr Asp Gln

```

				65					70					75
Leu	Val	Leu	Ile	Phe	Ala	Gly	Lys	Ile	Leu	Lys	Asp	Gln	Asp	Thr
				80					85					90
Leu	Ser	Gln	His	Gly	Ile	His	Asp	Gly	Leu	Thr	Val	His	Leu	Val
				95					100					105
Ile	Lys	Thr	Gln	Asn	Arg	Pro	Gln	Asp	His	Ser	Ala	Gln	Gln	Thr
				110					115					120
Asn	Thr	Ala	Gly	Ser	Asn	Val	Thr	Thr	Ser	Ser	Thr	Pro	Asn	Ser
				125					130					135
Asn	Ser	Thr	Ser	Gly	Ser	Ala	Thr	Ser	Asn	Pro	Phe	Gly	Leu	Gly
				140					145					150
Gly	Leu	Gly	Gly	Leu	Ala	Gly	Leu	Ser	Ser	Leu	Gly	Leu	Asn	Thr
				155					160					165
Thr	Asn	Phe	Ser	Glu	Leu	Gln	Ser	Gln	Met	Gln	Arg	Gln	Leu	Leu
				170					175					180
Ser	Asn	Pro	Glu	Met	Met	Val	Gln	Ile	Met	Glu	Asn	Pro	Phe	Val
				185					190					195
Gln	Ser	Met	Leu	Ser	Asn	Pro	Asp	Leu	Met	Arg	Gln	Leu	Ile	Met
				200					205					210
Ala	Asn	Pro	Gln	Met	Gln	Gln	Leu	Ile	Gln	Arg	Asn	Pro	Glu	Ile
				215					220					225
Ser	His	Met	Leu	Asn	Asn	Pro	Asp	Ile	Met	Arg	Gln	Thr	Leu	Glu
				230					235					240
Leu	Ala	Arg	Asn	Pro	Ala	Met	Met	Gln	Glu	Met	Met	Arg	Asn	Gln
				245					250					255
Asp	Arg	Ala	Leu	Ser	Asn	Leu	Glu	Ser	Ile	Pro	Gly	Gly	Tyr	Asn
				260					265					270
Ala	Leu	Arg	Arg	Met	Tyr	Thr	Asp	Ile	Gln	Glu	Pro	Met	Leu	Ser
				275					280					285
Ala	Ala	Gln	Glu	Gln	Phe	Gly	Gly	Asn	Pro	Phe	Ala	Ser	Leu	Val
				290					295					300
Ser	Asn	Thr	Ser	Ser	Gly	Glu	Gly	Ser	Gln	Pro	Ser	Arg	Thr	Glu
				305					310					315
Asn	Arg	Asp	Pro	Leu	Pro	Asn	Pro	Trp	Ala	Pro	Gln	Thr	Ser	Gln
				320					325					330
Ser	Ser	Ser	Ala	Ser	Ser	Gly	Thr	Ala	Ser	Thr	Val	Gly	Gly	Thr
				335					340					345
Thr	Gly	Ser	Thr	Ala	Ser	Gly	Thr	Ser	Gly	Gln	Ser	Thr	Thr	Ala
				350					355					360
Pro	Asn	Leu	Val	Pro	Gly	Val	Gly	Ala	Ser	Met	Phe	Asn	Thr	Pro
				365					370					375
Gly	Met	Gln	Ser	Leu	Leu	Gln	Gln	Ile	Thr	Glu	Asn	Pro	Gln	Leu
				380					385					390
Met	Gln	Asn	Met	Leu	Ser	Ala	Pro	Tyr	Met	Arg	Ser	Met	Met	Gln
				395					400					405
Ser	Leu	Ser	Gln	Asn	Pro	Asp	Leu	Ala	Ala	Gln	Met	Met	Leu	Asn
				410					415					420
Asn	Pro	Leu	Phe	Ala	Gly	Asn	Pro	Gln	Leu	Gln	Glu	Gln	Met	Arg
				425					430					435
Gln	Gln	Leu	Pro	Thr	Phe	Leu	Gln	Gln	Met	Gln	Asn	Pro	Asp	Thr
				440					445					450
Leu	Ser	Ala	Met	Ser	Asn	Pro	Arg	Ala	Met	Gln	Ala	Leu	Leu	Gln
				455					460					465
Ile	Gln	Gln	Gly	Leu	Gln	Thr	Leu	Ala	Thr	Glu	Ala	Pro	Gly	Leu
				470					475					480
Ile	Pro	Gly	Phe	Thr	Pro	Gly	Leu	Gly	Ala	Leu	Gly	Ser	Thr	Gly
				485					490					495
Gly	Ser	Ser	Gly	Thr	Asn	Gly	Ser	Asn	Ala	Thr	Pro	Ser	Glu	Asn
				500					505					510
Thr	Ser	Pro	Thr	Ala	Gly	Thr	Thr	Glu	Pro	Gly	His	Gln	Gln	Phe
				515					520					525
Ile	Gln	Gln	Met	Leu	Gln	Ala	Leu	Ala	Gly	Val	Asn	Pro	Gln	Leu
				530					535					540
Gln	Asn	Pro	Glu	Val	Arg	Phe	Gln	Gln	Gln	Leu	Glu	Gln	Leu	Ser
				545					550					555
Ala	Met	Gly	Phe	Leu	Asn	Arg	Glu	Ala	Asn	Leu	Gln	Ala	Leu	Ile
				560					565					570

Ala Thr Gly Gly Asp Ile Asn Ala Ala Ile Glu Arg Leu Leu Gly  
 575 580 585  
 Ser Gln Pro Ser

<210> 4  
 <211> 775  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1299481CD1

<400> 4  
 Met Thr Ile Val Asp Lys Ala Ser Glu Ser Ser Asp Pro Ser Ala  
 1 5 10 15  
 Tyr Gln Asn Gln Pro Gly Ser Ser Glu Ala Val Ser Pro Gly Asp  
 20 25 30  
 Met Asp Ala Gly Ser Ala Ser Trp Gly Ala Val Ser Ser Leu Asn  
 35 40 45  
 Asp Val Ser Asn His Thr Leu Ser Leu Gly Pro Val Pro Gly Ala  
 50 55 60  
 Val Val Tyr Ser Ser Ser Ser Val Pro Asp Lys Ser Lys Pro Ser  
 65 70 75  
 Pro Gln Lys Asp Gln Ala Leu Gly Asp Gly Ile Ala Pro Pro Gln  
 80 85 90  
 Lys Val Leu Phe Pro Ser Glu Lys Ile Cys Leu Lys Trp Gln Gln  
 95 100 105  
 Thr His Arg Val Gly Ala Gly Leu Gln Asn Leu Gly Asn Thr Cys  
 110 115 120  
 Phe Ala Asn Ala Ala Leu Gln Cys Leu Thr Tyr Thr Pro Pro Leu  
 125 130 135  
 Ala Asn Tyr Met Leu Ser His Glu His Ser Lys Thr Cys His Ala  
 140 145 150  
 Glu Gly Phe Cys Met Met Cys Thr Met Gln Ala His Ile Thr Gln  
 155 160 165  
 Ala Leu Ser Asn Pro Gly Asp Val Ile Lys Pro Met Phe Val Ile  
 170 175 180  
 Asn Glu Met Arg Arg Ile Ala Arg His Leu Arg Phe Gly Asn Gln  
 185 190 195  
 Glu Asp Ala His Glu Phe Leu Gln Tyr Thr Val Asp Ala Met Gln  
 200 205 210  
 Lys Ala Cys Leu Asn Gly Ser Asn Lys Leu Asp Arg His Thr Gln  
 215 220 225  
 Ala Thr Thr Leu Val Cys Gln Ile Phe Gly Tyr Leu Arg Ser  
 230 235 240  
 Arg Val Lys Cys Leu Asn Cys Lys Gly Val Ser Asp Thr Phe Asp  
 245 250 255  
 Pro Tyr Leu Asp Ile Thr Leu Glu Ile Lys Ala Ala Gln Ser Val  
 260 265 270  
 Asn Lys Ala Leu Glu Gln Phe Val Lys Pro Glu Gln Leu Asp Gly  
 275 280 285  
 Glu Asn Ser Tyr Lys Cys Ser Lys Cys Lys Lys Met Val Pro Ala  
 290 295 300  
 Ser Lys Arg Phe Thr Ile His Arg Ser Ser Asn Val Leu Thr Leu  
 305 310 315  
 Ser Leu Lys Arg Phe Ala Asn Phe Thr Gly Gly Lys Ile Ala Lys  
 320 325 330  
 Asp Val Lys Tyr Pro Glu Tyr Leu Asp Ile Arg Pro Tyr Met Ser  
 335 340 345  
 Gln Pro Asn Gly Glu Pro Ile Val Tyr Val Leu Tyr Ala Val Leu  
 350 355 360  
 Val His Thr Gly Phe Asn Cys His Ala Gly His Tyr Phe Cys Tyr  
 365 370 375  
 Ile Lys Ala Ser Asn Gly Leu Trp Tyr Gln Met Asn Asp Ser Ile  
 380 385 390  
 Val Ser Thr Ser Asp Ile Arg Ser Val Leu Ser Gln Gln Ala Tyr

	395								400						405
Val	Leu	Phe	Tyr	Ile	Arg	Ser	His	Asp	Val	Lys	Asn	Gly	Gly	Glu	
	410								415					420	
Leu	Thr	His	Pro	Thr	His	Ser	Pro	Gly	Gln	Ser	Ser	Pro	Arg	Pro	
	425								430					435	
Val	Ile	Ser	Gln	Arg	Val	Val	Thr	Asn	Lys	Gln	Ala	Ala	Pro	Gly	
	440								445					450	
Phe	Ile	Gly	Pro	Gln	Leu	Pro	Ser	His	Met	Ile	Lys	Asn	Pro	Pro	
	455								460					465	
His	Leu	Asn	Gly	Thr	Gly	Pro	Leu	Lys	Asp	Thr	Pro	Ser	Ser	Ser	
	470								475					480	
Met	Ser	Ser	Pro	Asn	Gly	Asn	Ser	Ser	Val	Asn	Arg	Ala	Ser	Pro	
	485								490					495	
Val	Asn	Ala	Ser	Ala	Ser	Val	Gln	Asn	Trp	Ser	Val	Asn	Arg	Ser	
	500								505					510	
Ser	Val	Ile	Pro	Glu	His	Pro	Lys	Lys	Gln	Lys	Ile	Thr	Ile	Ser	
	515								520					525	
Ile	His	Asn	Lys	Leu	Pro	Val	Arg	Gln	Cys	Gln	Ser	Gln	Pro	Asn	
	530								535					540	
Leu	His	Ser	Asn	Ser	Leu	Glu	Asn	Pro	Thr	Lys	Pro	Val	Pro	Ser	
	545								550					555	
Ser	Thr	Ile	Thr	Asn	Ser	Ala	Val	Gln	Ser	Thr	Ser	Asn	Ala	Ser	
	560								565					570	
Thr	Met	Ser	Val	Ser	Ser	Lys	Val	Thr	Lys	Pro	Ile	Pro	Arg	Ser	
	575								580					585	
Glu	Ser	Cys	Ser	Gln	Pro	Val	Met	Asn	Gly	Lys	Ser	Lys	Leu	Asn	
	590								595					600	
Ser	Ser	Val	Leu	Val	Pro	Tyr	Gly	Ala	Glu	Ser	Ser	Glu	Asp	Ser	
	605								610					615	
Asp	Glu	Glu	Ser	Lys	Gly	Leu	Gly	Lys	Glu	Asn	Gly	Ile	Gly	Thr	
	620								625					630	
Ile	Val	Ser	Ser	His	Ser	Pro	Gly	Gln	Asp	Ala	Glu	Asp	Glu	Glu	
	635								640					645	
Ala	Thr	Pro	His	Glu	Leu	Gln	Glu	Pro	Met	Thr	Leu	Asn	Gly	Ala	
	650								655					660	
Asn	Ser	Ala	Asp	Ser	Asp	Ser	Asp	Pro	Lys	Glu	Asn	Gly	Leu	Ala	
	665								670					675	
Pro	Asp	Gly	Ala	Ser	Cys	Gln	Gly	Gln	Pro	Ala	Leu	His	Ser	Glu	
	680								685					690	
Asn	Pro	Phe	Ala	Lys	Ala	Asn	Gly	Leu	Pro	Gly	Lys	Leu	Met	Pro	
	695								700					705	
Ala	Pro	Leu	Leu	Ser	Leu	Pro	Glu	Asp	Lys	Ile	Leu	Glu	Thr	Phe	
	710								715					720	
Arg	Leu	Ser	Asn	Lys	Leu	Lys	Gly	Ser	Thr	Asp	Glu	Met	Ser	Ala	
	725								730					735	
Pro	Gly	Ala	Glu	Arg	Gly	Pro	Pro	Glu	Asp	Arg	Asp	Ala	Glu	Pro	
	740								745					750	
Gln	Pro	Gly	Ser	Pro	Ala	Ala	Glu	Ser	Leu	Glu	Glu	Pro	Asp	Ala	
	755								760					765	
Ala	Ala	Ser	Leu	Phe	Pro	Phe	Ser	Glu	Gly						
	770								775						

&lt;210&gt; 5

&lt;211&gt; 351

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1873139CD1

&lt;400&gt; 5

Met	Asn	Ala	Ile	Leu	Gln	Ser	Leu	Ser	Asn	Ile	Glu	Gln	Phe	Cys	
1				5					10					25	
Cys	Tyr	Phe	Lys	Glu	Leu	Pro	Ala	Val	Glu	Leu	Arg	Asn	Gly	Lys	
				20					25					30	
Thr	Ala	Gly	Arg	Arg	Thr	Tyr	His	Thr	Arg	Ser	Gln	Gly	Asp	Asn	
				35					40					45	

```

Asn Val Ser Leu Val Glu Glu Phe Arg Lys Thr Leu Cys Ala Leu
50 55 60
Trp Gln Gly Ser Gln Thr Ala Phe Ser Pro Glu Ser Leu Phe Tyr
65 70 75
Val Val Trp Lys Ile Met Pro Asn Phe Arg Gly Tyr Gln Gln Gln
80 85 90
Asp Ala His Glu Phe Met Arg Tyr Leu Leu Asp His Leu His Leu
95 100 105
Glu Leu Gln Gly Gly Phe Asn Gly Val Ser Arg Ser Ala Ile Leu
110 115 120
Gln Glu Asn Ser Thr Leu Ser Ala Ser Asn Lys Cys Cys Ile Asn
125 130 135
Gly Ala Ser Thr Val Val Thr Ala Ile Phe Gly Gly Ile Leu Gln
140 145 150
Asn Glu Val Asn Cys Leu Ile Cys Gly Thr Glu Ser Arg Lys Phe
155 160 165
Asp Pro Phe Leu Asp Leu Ser Leu Asp Ile Pro Ser Gln Phe Arg
170 175 180
Ser Lys Arg Ser Lys Asn Gln Glu Asn Gly Pro Val Cys Ser Leu
185 190 195
Arg Asp Cys Leu Arg Ser Phe Thr Asp Leu Glu Glu Leu Asp Glu
200 205 210
Thr Glu Leu Tyr Met Cys His Lys Cys Lys Lys Lys Gln Lys Ser
215 220 225
Thr Lys Lys Phe Trp Ile Gln Lys Leu Pro Lys Val Leu Cys Leu
230 235 240
His Leu Lys Arg Phe His Trp Thr Ala Tyr Leu Arg Asn Lys Val
245 250 255
Asp Thr Tyr Val Glu Phe Pro Leu Arg Gly Leu Asp Met Lys Cys
260 265 270
Tyr Leu Leu Glu Pro Glu Asn Ser Gly Pro Glu Ser Cys Leu Tyr
275 280 285
Asp Leu Ala Ala Val Val Val His His Gly Ser Gly Val Gly Ser
290 295 300
Gly His Tyr Thr Ala Tyr Ala Thr His Glu Gly Arg Trp Phe His
305 310 315
Phe Asn Asp Ser Thr Val Thr Leu Thr Asp Glu Glu Thr Val Val
320 325 330
Lys Ala Lys Ala Tyr Ile Leu Phe Tyr Val Glu His Gln Ala Lys
335 340 345
Ala Gly Ser Asp Lys Leu
350

```

&lt;210&gt; 6

&lt;211&gt; 136

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1903112CD1

&lt;400&gt; 6

```

Met Ala Leu Met Gln Arg Ser Asp Ile Phe Arg Val Ala Ile Ala
1 5 10 15
Gly Ala Pro Val Thr Leu Trp Ile Phe Tyr Asp Thr Gly Tyr Thr
20 25 30
Glu Arg Tyr Met Gly His Pro Asp Gln Asn Glu Gln Gly Tyr Tyr
35 40 45
Leu Gly Ser Val Ala Met Gln Ala Glu Lys Phe Pro Ser Glu Pro
50 55 60
Asn Arg Leu Leu Leu Leu His Gly Phe Leu Asp Glu Asn Val His
65 70 75
Phe Ala His Thr Ser Ile Leu Leu Ser Phe Leu Val Arg Ala Gly
80 85 90
Lys Pro Tyr Asp Leu Gln Ile Tyr Pro Gln Glu Arg His Ser Ile
95 100 105
Arg Val Pro Glu Ser Gly Glu His Tyr Glu Leu His Leu Leu His

```

	110		115	120
Tyr	Leu	Gln	Glu	Asn
	125	Leu	Gly	Ser
		Arg	Arg	Ile
			Ala	Ala
			Leu	Lys
				Val
				135

Ile

&lt;210&gt; 7

&lt;211&gt; 396

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Inocyte ID No: 1993044CD1

&lt;400&gt; 7

Met	Ser	Leu	Gly	Trp	Leu	Glu	Arg	Pro	Pro	Ala	Leu	Ser	Arg	Ala
1				5					10					15
Ala	Gly	Asp	Gly	Ala	Arg	Arg	Leu	Ser	Gly	Ser	Met	Arg	Gly	Asp
				20					25					30
Val	Trp	Leu	Thr	Ser	Ser	Ala	Ala	Gly	Leu	Leu	Arg	Ser	Val	Ala
				35					40					45
Gly	Gly	Ser	Trp	Cys	Gly	Gly	Gln	Leu	Arg	Ala	Arg	Gly	Gly	Ser
				50					55					60
Gly	Arg	Cys	Val	Ala	Arg	Ala	Met	Thr	Gly	Asn	Ala	Gly	Glu	Trp
				65					70					75
Cys	Leu	Met	Glu	Ser	Asp	Pro	Gly	Val	Phe	Thr	Glu	Leu	Ile	Lys
				80					85					90
Gly	Phe	Gly	Cys	Arg	Gly	Ala	Gln	Val	Glu	Glu	Ile	Trp	Ser	Leu
				95					100					105
Glu	Pro	Glu	Asn	Phe	Glu	Lys	Leu	Lys	Pro	Val	His	Gly	Leu	Ile
				110					115					120
Phe	Leu	Phe	Lys	Trp	Gln	Pro	Gly	Glu	Glu	Pro	Ala	Gly	Ser	Val
				125					130					135
Val	Gln	Asp	Ser	Arg	Leu	Asp	Thr	Ile	Phe	Phe	Ala	Lys	Gln	Val
				140					145					150
Ile	Asn	Asn	Ala	Cys	Ala	Thr	Gln	Ala	Ile	Val	Ser	Val	Leu	Leu
				155					160					165
Asn	Cys	Thr	His	Gln	Asp	Val	His	Leu	Gly	Glu	Thr	Leu	Ser	Glu
				170					175					180
Phe	Lys	Glu	Phe	Ser	Gln	Ser	Phe	Asp	Ala	Ala	Met	Lys	Gly	Leu
				185					190					195
Ala	Leu	Ser	Asn	Ser	Asp	Val	Ile	Arg	Gln	Val	His	Asn	Ser	Phe
				200					205					210
Ala	Arg	Gln	Gln	Met	Phe	Glu	Phe	Asp	Thr	Lys	Thr	Ser	Ala	Lys
				215					220					225
Glu	Glu	Asp	Ala	Phe	His	Phe	Val	Ser	Tyr	Val	Pro	Val	Asn	Gly
				230					235					240
Arg	Leu	Tyr	Glu	Leu	Asp	Gly	Leu	Arg	Glu	Gly	Pro	Ile	Asp	Leu
				245					250					255
Gly	Ala	Cys	Asn	Gln	Asp	Asp	Trp	Phe	Ser	Ala	Val	Arg	Pro	Val
				260					265					270
Ile	Glu	Lys	Arg	Ile	Gln	Lys	Tyr	Ser	Glu	Gly	Glu	Ile	Arg	Phe
				275					280					285
Asn	Leu	Met	Ala	Ile	Val	Ser	Asp	Arg	Lys	Met	Ile	Tyr	Glu	Gln
				290					295					300
Lys	Ile	Ala	Glu	Leu	Gln	Arg	Gln	Leu	Ala	Glu	Glu	Glu	Pro	Met
				305					310					315
Asp	Thr	Asp	Gln	Gly	Asn	Ser	Met	Leu	Ser	Ala	Ile	Gln	Ser	Glu
				320					325					330
Val	Ala	Lys	Asn	Gln	Met	Leu	Ile	Glu	Glu	Glu	Val	Gln	Lys	Leu
				335					340					345
Lys	Arg	Tyr	Lys	Ile	Glu	Asn	Ile	Arg	Arg	Lys	His	Asn	Tyr	Leu
				350					355					360
Pro	Phe	Ile	Met	Glu	Leu	Leu	Lys	Thr	Leu	Ala	Glu	His	Gln	Gln
				365					370					375
Leu	Ile	Pro	Leu	Val	Glu	Lys	Ala	Lys	Glu	Lys	Gln	Asn	Ala	Lys
				380					385					390

Lys Ala Gln Glu Thr Lys  
395

<210> 8  
<211> 246  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2292182CD1

<400> 8  
Met Ala Gly Ala Pro Asp Glu Arg Arg Arg Gly Pro Ala Ala Gly  
1 5 10 15  
Glu Gln Leu Gln Gln Gln His Val Ser Cys Gln Val Phe Pro Glu  
20 25 30  
Arg Leu Ala Gln Gly Asn Pro Gln Gln Gly Phe Phe Ser Ser Phe  
35 40 45  
Phe Thr Ser Asn Gln Lys Cys Gln Leu Arg Leu Leu Lys Thr Leu  
50 55 60  
Glu Thr Asn Pro Tyr Val Lys Leu Leu Leu Asp Ala Met Lys His  
65 70 75  
Ser Gly Cys Ala Val Asn Lys Asp Arg His Phe Ser Cys Glu Asp  
80 85 90  
Cys Asn Gly Asn Val Ser Gly Gly Phe Asp Ala Ser Thr Ser Gln  
95 100 105  
Ile Val Leu Cys Gln Asn Asn Ile His Asn Gln Ala His Met Asn  
110 115 120  
Arg Val Val Thr His Glu Leu Ile His Ala Phe Asp His Cys Arg  
125 130 135  
Ala His Val Asp Trp Phe Thr Asn Ile Arg His Leu Ala Cys Ser  
140 145 150  
Glu Val Arg Ala Ala Asn Leu Ser Gly Asp Cys Ser Leu Val Asn  
155 160 165  
Glu Ile Phe Arg Leu His Phe Gly Leu Lys Gln His His Gln Thr  
170 175 180  
Cys Val Arg Asp Arg Ala Thr Leu Ser Ile Leu Ala Val Arg Asn  
185 190 195  
Ile Ser Lys Glu Val Ala Lys Lys Ala Val Asp Glu Val Phe Glu  
200 205 210  
Ser Cys Phe Asn Asp His Glu Pro Phe Gly Arg Ile Pro His Asn  
215 220 225  
Lys Thr Tyr Ala Arg Tyr Ala His Arg Asp Phe Glu Asn Arg Asp  
230 235 240  
Arg Tyr Tyr Ser Asn Ile  
245

<210> 9  
<211> 262  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2331301CD1

<400> 9  
Met Glu Val Tyr Ile Arg His Leu Glu Lys Val Leu Arg Arg Tyr  
1 5 10 15  
Val Gln Arg Leu Gln Trp Leu Leu Ser Gly Ser Arg Arg Leu Phe  
20 25 30  
Gly Thr Val Leu Glu Ser Lys Val Cys Ile Leu Leu Asp Thr Ser  
35 40 45  
Gly Ser Met Gly Pro Tyr Leu Gln Gln Val Lys Thr Gln Leu Val  
50 55 60  
Leu Leu Ile Trp Glu Gln Leu Arg Lys Cys Cys Asp Ser Phe Asn  
65 70 75  
Leu Leu Ser Phe Ala Glu Ser Leu Gln Ser Trp Gln Asp Thr Leu

	80		85		90
Val Glu Thr Thr Asp	Ala Ala Cys His	Glu Ala Met Gln Trp	Val		
	95		100		105
Thr His Leu Gln Ala	Gln Gly Ser Thr	Ser Ile Leu Gln Ala	Leu		
	110		115		120
Leu Lys Ala Phe Ser	Phe His Asp Leu	Glu Gly Leu Tyr Leu	Leu		
	125		130		135
Thr Asp Gly Lys Pro	Asp Thr Ser Cys	Ser Leu Val Leu Asn	Glu		
	140		145		150
Val Gln Lys Leu Arg	Glu Lys Arg Asp	Val Lys Val His Thr	Ile		
	155		160		165
Ser Leu Asn Cys Ser	Asp Arg Ala Ala	Val Glu Phe Leu Arg	Lys		
	170		175		180
Leu Ala Ser Phe Thr	Gly Gly Arg Tyr	His Cys Pro Val Gly	Glu		
	185		190		195
Asp Thr Leu Ser Lys	Ile His Ser Leu	Leu Thr Lys Gly Phe	Ile		
	200		205		210
Asn Glu Lys Asp Arg	Thr Leu Pro Pro	Phe Glu Gly Asp Asp	Leu		
	215		220		225
Arg Ile Leu Ala Gln	Glu Ile Thr Lys	Ala Arg Ser Phe Leu	Trp		
	230		235		240
Gln Ala Gln Ser Phe	Arg Ser Gln Leu	Gln Lys Lys Asn Asp	Ala		
	245		250		255
Glu Pro Lys Val Thr	Leu Ser				
	260				

&lt;210&gt; 10

&lt;211&gt; 406

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2517512CD1

&lt;400&gt; 10

Met Ala Ala Ala Val	Arg Gln Asp Leu	Ala Gln Leu Met Asn	Ser
1	5	10	15
Ser Gly Ser His Lys	Asp Leu Ala Gly	Lys Tyr Arg Gln Ile	Leu
	20	25	30
Glu Lys Ala Ile Gln	Leu Ser Gly Ala	Glu Gln Leu Glu Ala	Leu
	35	40	45
Lys Ala Phe Val Glu	Ala Met Val Asn	Glu Asn Val Ser Leu	Val
	50	55	60
Ile Ser Arg Gln Leu	Thr Asp Phe Cys	Thr His Leu Pro	Asn
	65	70	75
Leu Pro Asp Ser Thr	Ala Lys Glu Ile	Tyr His Phe Thr	Leu
	80	85	90
Lys Ile Gln Pro Arg	Val Ile Ser Phe	Glu Gln Val Ala	Ser
	95	100	105
Ile Arg Gln His Leu	Ala Ser Ile Tyr	Glu Lys Glu Glu	Asp
	110	115	120
Arg Asn Ala Ala Gln	Val Leu Val Gly	Ile Pro Leu Glu	Thr
	125	130	135
Gln Lys Gln Tyr Asn	Val Asp Tyr Lys	Leu Glu Thr Tyr	Leu
	140	145	150
Ile Ala Arg Leu Tyr	Leu Glu Asp Asp	Asp Pro Val Gln	Ala
	155	160	165
Ala Tyr Ile Asn Arg	Ala Ser Leu Leu	Gln Asn Glu Ser	Thr
	170	175	180
Glu Gln Leu Gln Ile	His Tyr Lys Val	Cys Tyr Ala Arg	Val
	185	190	195
Asp Tyr Arg Arg Lys	Phe Ile Glu Ala	Ala Gln Arg Tyr	Asn
	200	205	210
Leu Ser Tyr Lys Thr	Ile Val His Glu	Ser Glu Arg Leu	Glu
	215	220	225
Leu Lys His Ala Leu	His Cys Thr Ile	Leu Ala Ser Ala	Gly
	230	235	240



Gln	Arg	Ser	Arg	Met	Leu	Ala	Thr	Leu	Phe	Lys	Asp	Glu	Arg	Cys	
				245					250					255	
Gln	Gln	Leu	Ala	Ala	Tyr	Gly	Ile	Leu	Glu	Lys	Met	Tyr	Leu	Asp	
				260					265					270	
Arg	Ile	Ile	Arg	Gly	Asn	Gln	Leu	Gln	Glu	Phe	Ala	Ala	Met	Leu	
				275					280					285	
Met	Pro	His	Gln	Lys	Ala	Thr	Thr	Ala	Asp	Gly	Ser	Ser	Ile	Leu	
				290					295					300	
Asp	Arg	Ala	Val	Ile	Glu	His	Asn	Leu	Leu	Ser	Ala	Ser	Lys	Leu	
				305					310					315	
Tyr	Asn	Asn	Ile	Thr	Phe	Glu	Glu	Leu	Gly	Ala	Leu	Leu	Glu	Ile	
				320					325					330	
Pro	Ala	Ala	Lys	Ala	Glu	Lys	Ile	Ala	Ser	Gln	Met	Ile	Thr	Glu	
				335					340					345	
Gly	Arg	Met	Asn	Gly	Phe	Ile	Asp	Gln	Ile	Asp	Gly	Ile	Val	His	
				350					355					360	
Phe	Glu	Thr	Arg	Glu	Ala	Leu	Pro	Thr	Trp	Asp	Lys	Gln	Ile	Gln	
				365					370					375	
Ser	Leu	Cys	Phe	Gln	Val	Asn	Asn	Leu	Leu	Glu	Lys	Ile	Ser	Gln	
				380					385					390	
Thr	Ala	Pro	Glu	Trp	Thr	Ala	Gln	Ala	Met	Glu	Ala	Gln	Met	Ala	
				395					400					405	

Gln

<210> 11  
 <211> 172  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3489039CD1

Met	Leu	Leu	Pro	Asn	Ile	Leu	Leu	Thr	Gly	Thr	Pro	Gly	Val	Gly	
				5					10					15	
Lys	Thr	Thr	Leu	Gly	Lys	Glu	Leu	Ala	Ser	Lys	Ser	Gly	Leu	Lys	
				20					25					30	
Tyr	Ile	Asn	Val	Gly	Asp	Leu	Ala	Arg	Glu	Glu	Gln	Leu	Tyr	Asp	
				35					40					45	
Gly	Tyr	Asp	Glu	Glu	Tyr	Asp	Cys	Pro	Ile	Leu	Asp	Glu	Asp	Arg	
				50					55					60	
Val	Val	Asp	Glu	Leu	Asp	Asn	Gln	Met	Arg	Glu	Gly	Gly	Val	Ile	
				65					70					75	
Val	Asp	Tyr	His	Gly	Cys	Asp	Phe	Phe	Pro	Glu	Arg	Trp	Phe	His	
				80					85					90	
Ile	Val	Phe	Val	Leu	Arg	Thr	Asp	Thr	Asn	Val	Leu	Tyr	Glu	Arg	
				95					100					105	
Leu	Glu	Thr	Arg	Gly	Tyr	Asn	Glu	Lys	Lys	Leu	Thr	Asp	Asn	Ile	
				110					115					120	
Gln	Cys	Glu	Ile	Phe	Gln	Val	Leu	Tyr	Glu	Glu	Ala	Thr	Ala	Ser	
				125					130					135	
Tyr	Lys	Glu	Glu	Ile	Val	His	Gln	Leu	Pro	Ser	Asn	Lys	Pro	Glu	
				140					145					150	
Glu	Leu	Glu	Asn	Asn	Val	Asp	Gln	Ile	Leu	Lys	Trp	Ile	Glu	Gln	
				155					160					165	
Trp	Ile	Lys	Asp	His	Asn	Ser									
				170											

<210> 12  
 <211> 517  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5432879CD1

&lt;400&gt; 12

Met	Leu	Ser	Ser	Arg	Ala	Glu	Ala	Ala	Met	Thr	Ala	Ala	Asp	Arg
1				5					10					15
Ala	Ile	Gln	Arg	Phe	Leu	Arg	Thr	Gly	Ala	Ala	Val	Arg	Tyr	Lys
				20					25					30
Val	Met	Lys	Asn	Trp	Gly	Val	Ile	Gly	Gly	Ile	Ala	Ala	Ala	Leu
				35					40					45
Ala	Ala	Gly	Ile	Tyr	Val	Ile	Trp	Gly	Pro	Ile	Thr	Glu	Arg	Lys
				50					55					60
Lys	Arg	Arg	Lys	Gly	Leu	Val	Pro	Gly	Leu	Val	Asn	Leu	Gly	Asn
				65					70					75
Thr	Cys	Phe	Met	Asn	Ser	Leu	Leu	Gln	Gly	Leu	Ser	Ala	Cys	Pro
				80					85					90
Ala	Phe	Ile	Arg	Trp	Leu	Glu	Glu	Phe	Thr	Ser	Gln	Tyr	Ser	Arg
				95					100					105
Asp	Gln	Lys	Glu	Pro	Pro	Ser	His	Gln	Tyr	Leu	Ser	Leu	Thr	Leu
				110					115					120
Leu	His	Leu	Leu	Lys	Ala	Leu	Ser	Cys	Gln	Glu	Val	Thr	Asp	Asp
				125					130					135
Glu	Val	Leu	Asp	Ala	Ser	Cys	Leu	Leu	Asp	Val	Leu	Arg	Met	Tyr
				140					145					150
Arg	Trp	Gln	Ile	Ser	Ser	Phe	Glu	Glu	Gln	Asp	Ala	His	Glu	Leu
				155					160					165
Phe	His	Val	Ile	Thr	Ser	Ser	Leu	Glu	Asp	Glu	Arg	Asp	Arg	Gln
				170					175					180
Pro	Arg	Val	Thr	His	Leu	Phe	Asp	Val	His	Ser	Leu	Glu	Gln	Gln
				185					190					195
Ser	Glu	Ile	Thr	Pro	Lys	Gln	Ile	Thr	Cys	Arg	Thr	Arg	Gly	Ser
				200					205					210
Pro	His	Pro	Thr	Ser	Asn	His	Trp	Lys	Ser	Gln	His	Pro	Phe	His
				215					220					225
Gly	Arg	Leu	Thr	Ser	Asn	Met	Val	Cys	Lys	His	Cys	Glu	His	Gln
				230					235					240
Ser	Pro	Val	Arg	Phe	Asp	Thr	Phe	Asp	Ser	Leu	Ser	Leu	Ser	Ile
				245					250					255
Pro	Ala	Ala	Thr	Trp	Gly	His	Pro	Leu	Thr	Leu	Asp	His	Cys	Leu
				260					265					270
His	His	Phe	Ile	Ser	Ser	Glu	Ser	Val	Arg	Asp	Val	Val	Cys	Asp
				275					280					285
Asn	Cys	Thr	Lys	Ile	Glu	Ala	Lys	Gly	Thr	Leu	Asn	Gly	Glu	Lys
				290					295					300
Val	Glu	His	Gln	Arg	Thr	Thr	Phe	Val	Lys	Gln	Leu	Lys	Leu	Gly
				305					310					315
Lys	Leu	Pro	Gln	Cys	Leu	Cys	Ile	His	Leu	Gln	Arg	Leu	Ser	Trp
				320					325					330
Ser	Ser	His	Gly	Thr	Pro	Leu	Lys	Arg	His	Glu	His	Val	Gln	Phe
				335					340					345
Asn	Glu	Phe	Leu	Met	Met	Asp	Ile	Tyr	Lys	Tyr	His	Leu	Leu	Gly
				350					355					360
His	Lys	Pro	Ser	Gln	His	Asn	Pro	Lys	Leu	Asn	Lys	Asn	Pro	Gly
				365					370					375
Pro	Thr	Leu	Glu	Leu	Gln	Asp	Gly	Pro	Gly	Ala	Pro	Thr	Pro	Val
				380					385					390
Leu	Asn	Gln	Pro	Gly	Ala	Pro	Lys	Thr	Gln	Ile	Phe	Met	Asn	Gly
				395					400					405
Ala	Cys	Ser	Pro	Ser	Leu	Leu	Pro	Thr	Leu	Ser	Ala	Pro	Met	Pro
				410					415					420
Phe	Pro	Leu	Pro	Val	Val	Pro	Asp	Tyr	Ser	Ser	Ser	Thr	Tyr	Leu
				425					430					435
Phe	Arg	Leu	Met	Ala	Val	Val	Val	His	His	Gly	Asp	Met	His	Ser
				440					445					450
Gly	His	Phe	Val	Thr	Tyr	Arg	Arg	Ser	Pro	Pro	Ser	Ala	Arg	Asn
				455					460					465
Pro	Leu	Ser	Thr	Ser	Asn	Gln	Trp	Leu	Trp	Val	Ser	Asp	Asp	Thr
				470					475					480
Val	Arg	Lys	Ala	Ser	Leu	Gln	Glu	Val	Leu	Ser	Ser	Ser	Ala	Tyr
				485					490					495

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4993873CB1

<400> 51  
 agaatggacc tgacaaatat tgaccagggtt gctgtgatct tcaaacacca ttttctgtg 60  
 ggtagaggtg atgacgtgct gaagacotgy gccctgctc agtgcccttg ctctagaatg 120  
 ggtccagctt ggttttggt actgggaaca gggatcctgg cctctgtcca ctgtcagccc 180  
 cttcttggcc atggagataa aagtctycag gggcctcaac cccccaggca taagctctca 240  
 gagccagccc ccgctacca cagaatocaa cccaccatta ccacttttgc ttgtggtttg 300  
 tataaagagc tggcagraga cgcctccgga aacatcttct tctcgcaggt gagcatctcc 360  
 accaccctgg cctgtctctc tottngggcc caagctaaca cctcagctct gatcctggag 420  
 ggcttgggat tcaacctcac agaaacccct gaagccgaca tccaccaggg ctcccgaggc 480  
 ctctccacca cccttgccct gccccgccc gaaactcgac taagtagg aaactccctg 540  
 ttctcagaca agcgaactaa gctcggcag cactatttgg acagcatcaa ggagctttat 600  
 ggagobtttg cttttttctg caacttcana gattctgtta caactgggag gcagattaat 660  
 gactatttga gaaggcaaac atacgggcaa gtctgtgact gctcccgga gttcagccag 720  
 gacaggttca tggfllcttgc caattacatc ttcttcnag ccagtgga atgagaggac 780  
 agtcgttacc agaccagaa gcaggcaagt ttctttgtgg atgagaggac ttctctccag 840  
 gtccccatga tgcacccaaa ggaatycac agattcctct atyacaggga ttgtgcttgc 900  
 actgtctccc agatagatca cagaggaaat gcttggcgc tctgtgctct ccttgaccgc 960  
 gggaaatatga agcaggtgga ggctgctcty cagccacaga ccttgagaaa atggggccaa 1020  
 ttgtctctgc ccagtctgtt ggattttgac ttgccagggt tttaaatctc tggaacatat 1080  
 aacctggaag acatacttcc ccaaatttgt ctcaccaaca tactcaactt agaagctgac 1140  
 ttctcaggaag tcactgggca gctcaacaaa accatctcca agglgtcaca caagtcgatg 1200  
 gtggacatga gtgagaaggg gaccgaggcc ggggtctctt caggcctcct ctcccagccc 1260  
 ccatctctga acacccatgt gcacccacat gccaacttca cctctgtctc cctctgtctc 1320  
 ctttgggagg tcaaccacca gagcttactc ttctgggaa aagtgttcaa cctgggacca 1380  
 gggtaacctat ggtggggggc caggagtatt cttatctctat cttgggacca acagataggg 1440  
 cagaaaccagc ctgcacctct ggtctgtat gtggttcagt taatcagttg gccaaagattc 1500  
 taataaagtt gaccttgggt tctgtgaaaa aaaaaaaaag cgtccaaatt taacataagg 1560  
 ggggggttaa ggalgtgaa agcaaatacc cgggtttgac agtctgtttg atggccgtcg actgattggc 1620  
 ggggggggng agagagtttt tacgcatctt tggccgtgtt agcccccggg gggggggggc 1680  
 ccataaggcc ccgggagctg gagggttgtt ttgcgcgggc actcttgggg ggtgttttat 1740  
 gaccgacttt tagtggcaca aactgtgggg cttctctat gtgtgcgcgc gcttctgttt 1800

<210> 52  
 <211> 550  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5208004CB1

<400> 52  
 ggggggtggc tggaggacac ggtgtaggcc ttgcgcgtgl ctgggttctt cggcaactga 60  
 ggagcaggac tgtttctgga aacactgggc tgcgtgtgac ccgtcaaac tctcccaca 120  
 attctgaatc cgagaaagtg aaggaaggat ggtgggggag tggaggaggc gtagcagagg 180  
 ccacagggac ccagcagaga tgcqgtggag acagaggagc ttctttctca ggtgtttct 240  
 gggagacctg agaggcgggc aycaccaccc tccgtcact ctccctcag cctcttcagt 300  
 tcccttctca acccttctc tctctctgca ctcttccctt tottgcctg ttgttagtcc 360  
 ctgtcccaaa actcctggtt cttttgttct gctgcctgg cccccacca ggaggaggtc 420  
 taaggcaaca tccccccac gaggatocaa caaacagggt tcatgtctgg gctgggggag 480  
 ccccgctggg ttcttgatgc cttgtgcaca ggyaagtgtt gcagtcattt ttggactctc 540  
 ctgaatgtgt

<210> 53  
 <211> 2597  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5267783CB1

gttaatatctc	ctgctgttgag	aaacttttaa	gtttcaataa	cccaagatgc	ttcagtgctc	660
atagtggatt	actatgagcc	aaggagacag	goggtgagaa	gttacaactc	tgaagtgaag	720
ctgtctctct	gtgacotttg	cagtgatgtc	cagggctguc	gtccttgtga	ggatggagct	780
ccaggctccc	atcctcactc	ttcagtcatt	tttattttct	gtttcaagct	tctgtacttt	840
atggaccttt	ggctgtgatt	tcttttttaa	ggactctgtg	taacactaac	atttccagta	900
gtcacatgtg	attgttttgt	tttcttagaa	gaatactgct	tttatttttg	aaaaagagtt	960
ttttttcttt	ctatgggggt	gcagggatgg	tgtacaacag	gtcctagcat	gtatagctgc	1020
atagatttct	tcacctgctc	tttgtgtgga	agalcagaa	gaatgcagtt	gtgtgtctat	1080
attttccctc	ctcaaaatct	tttagaattt	ttttggaggt	gtttgttttc	tcacagaata	1140
aggtattact	ttagaatagg	tattctctct	atttttgtga	agaaatgaac	ctagattctt	1200
aagcttattt	acacatccat	gtttgtttaa	agatggattt	ccctgggaat	gggagaaaac	1260
agccagcagg	aggagcttca	tctgttctct	tcccactctc			1300

<210> 50  
 <211> 2241  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3883861CB1

<400> 50	ccgggctgca	ccggaggcgg	cgagatggtc	gcgcgcgtcg	gootctctgt	60
gctcccagag	cagctgtctac	tttggggcra	cttggaagcc	agcccgccgg	agcccgccgg	120
ggcgcccttg	cgcaaggagg	cggaggcatt	cctagagaa	taaggatacc	tcactgaaca	180
ggtctccaaa	gctcccacct	ccactcgatt	cagcgatgcc	atcagagcgt	ttcagtgggt	240
gtcccagcta	ctgtctagcg	gogtcttqga	ccgcgccacc	ctgcgccaga	tgactctctc	300
cgcctggggg	gttacagata	ccaacagtta	tgccgcttgg	gctgagagga	tcagtgaact	360
gtttgtctaga	ccacgggacc	aaatgagggc	laagaaacgc	tttgcacagg	aaagtaacaa	420
atggtacaa	cagcaacctct	ectaccgctc	gggtgaactg	cctgagcctc	tgccggagcc	480
ggcagtttgg	ggcgcctgtg	gcgcgcctct	ccagtttctg	agcaagctct	cagcgtctga	540
gttctggggg	gccccagcca	caggccccgc	tgcaatccgg	ctcaactctc	tcacagggga	600
ccacacagat	gggctgggca	atgcctttga	tgcccccagg	ggcgccctgg	cgccgcctct	660
cctgccccgc	cgccggggaag	cgcaactctg	ccaagatgag	cgctggtccc	lgagcccgcc	720
ccgcgggggc	aaactgtttc	tggtgcttgc	gcacagagtc	ggtcacacgc	ttggtctcac	780
ccactcggcc	gcgcggggcg	cgctccttgc	gcctcctctc	aaagggctgg	ggcgggagcc	840
gctgctcagg	tgggacagag	tgctggccgt	gcagagcctg	tatgggaagg	ccctaggggg	900
ctcagtgccc	gtccagctcc	caggaaagct	gttcaactgac	tttgagacct	gggactctta	960
cagcccccaa	ggaaggccgc	ctgaaacgca	gggccccaaa	tactgccaat	cttctcttga	1020
tgccatcact	gtagacagge	aacagcaact	gtacattttt	aaagggagcc	atttctggga	1080
ggtggcagct	gatgggaacg	lctcagagcc	ccgtccactg	caggaaagat	gggtcgggct	1140
gccccccaac	attgaggtct	cggcagtgct	attgaatgat	ggagatttct	acttcttcaa	1200
aggggggtcg	tgctggaggt	tcgggggccc	caagccagtg	tggggtctct	caacagctgt	1260
ccgggcaggg	ggcctgcctc	gcaatcttga	gcgcgcctct	ttcttccctc	ctctgcgcgc	1320
cctctctctc	ttcaaggggt	cccgctacta	cgtgctggcc	cgagggggac	tgcaagtggg	1380
gcctactctc	ccccgaagtc	tgccagagct	ggggggccat	cctgagggag	tcagccagcc	1440
cctgcccagg	cccgatggct	ccatctctct	cttccagagat	gacccgtact	ggcgccctcg	1500
ccaggccaaa	ctgcaggcaa	ccacctcgag	cccgctggcc	accagagctg	cctggatggg	1560
ctgctggcat	gccaactcgg	ggagcgccct	gttctgaagg	caactctctc	cctcagaaac	1620
tggttggtgt	ctcaggggca	aatcalttct	cccccccccc	gggcagaaac	cctctllaga	1680
gcctctgagt	ccctctgcag	aagaccgggc	agcaaaagct	ccatctggaa	gtctgtctgc	1740
ctttgttctc	tgaggaatgc	agcatttgtt	ttgtctgtct	ccaccacatg	gaagtggggg	1800
tgggatcaat	cttaggaaaa	gcaaaaaagg	gtccagatct	ccttggccct	ttctccagag	1860
gacttctatc	ctcccagggc	ctttgttctc	tgggtanaag	gtacagtctc	tttcagaggg	1920
taacagcact	gggatcccaag	cagggggatg	aaaaactcag	cagagaaatt	cgagaccatt	1980
ttgcaagact	gtgcctctct	cctcaggga	ccctgggtct	gttcttgaaa	aaaggtgtct	2040
tatttagtca	gagggccccc	cccgaggaag	catggatggg	gatgaaggca	caggcgtctc	2100
caactctaga	ggcccttctg	gggtctagga	ccacagagtg	gagggagact	gatgcaggcc	2160
taccagctcc	tggttttttg	tctggggctg	gaataaagag	gtgccttcag	ctggctgggc	2220
gagaggccag	aaaaaaaaaa	a				2241

<210> 51  
 <211> 1860  
 <212> DNA  
 <213> Homo sapiens

```

tgcccagggg agctgacgag cagaggagaa ggtcccttt ccagatgctt ctgclgctgg 780
agdagatgca ggacagccgg cagaaagcag tgcggccctt ggagctggcc tactgctgc 810
agaagtgcna cgtgcccttg ttgtgccaa cgtgctgtgc ccaactgtac ctcaaaactct 900
ggaacctgat taaggaccag atcaactgat tgcacttggg ggaagagactg caggccctgt 960
atcagatccg ggtgaaggaa tcccttgaat gcttggactg tgccatggag agtagcagaa 1020
acagcagcat gctcaacctc ccactttctc ttttggatgt ggaactcaag cccclgaaga 1080
cactggagga cgccttgac gaactgtggg aagaagacc cgttctgtga aaaagcaagt 1140
gcttctgtga cctgacactt caactcatgc gttgggaaca gttcttgaa ctgacccatt 1200
tgccccagac cctgacactt bccccccaga gattctccat caggaattca cagacagaa 1260
agatctgcca ctccctgtac gaggagcagt gcttggatbt cagccagatc ctccaatga 1320
agcyagagtc ttgtgatgct gactccggtc ctggagggca gtatgagctt ttgtctgtga 1380
ttgggcacgt ggaatggaa gactccggtc attactgtgt ctacatccgg aatyctgtgg 1440
atggaaaetg gttctgtctc aatgaactca atatttgcct ggtgtctcgg gaagacatcc 1500
agtgtaccta cggaaatcct aactaccact aatgggaaac tgcatactct caggtttaca 1560
tgaagtggga gtgctaattg aaatgcccaa aaccttcaga gattgacacg ctgtcatttt 1620
ccatttccgt tccctggatc aaggagtctt ctgaagagatt ttgcaattgag gaggagcatt 1680
gttttcaaac tatataactg agccttattt ataattaggg atattatcaa aatattgtaa 1740
catgaggccc ctccaggtct tgcagtcag aatggatgct tccaccagca gaccctggca 1800
tgttggctgt cggctcctgg tgctcgtgc tggcaagac attagccctt gcagtgggag 1860
cctgtgggaa cttcaggggg tcccagtggt gagagcagty clgctgtgac cagacttgta 1920
gcaaaaggtc agtggcaggg ggtatttcag tattatacaa clgctgtgac cagacttgta 1980
tactggctga atatcagtyc tgtttgtaac atttacttt gagaaacca 2040
takgaatcaa gtattttgta actgctatct ttttacttt caaatattta ttgatcatct 2100
cttctccata agatagtgtg ataaccacag tcatgaataa agttattttc caaaaaaaaa 2160
aa 2162

```

<210> 48  
 <211> 578  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3092341CB1

```

<400> 48
gaaaggccca cctgtgtcct ggttgagggt ctccagggtt ctttggggcc cgaggaaatt 60
cagacctgaa caatcaggtg gacaacttct tgtgactttg ggtattctga aattactcat 120
cttgactcta ctgtgecttg atgtagctgt tgtttttgtg tgatccctgt tcatggccaa 180
tggtggcaga gtctacatag aactatgctt cgttgtgttc tggggaaaac ctttggactt 240
gttggctata ctattcaata tggctgtata gctcattgtg cttttgaata cgttggtgtt 300
gttctcatgg tgcgaatggg loatgttttg llagaagggtg acaatctaca gaattctaca 360
gattccaggt gctatggacc tattccatat ggactaataa gaggaacgat cttcttttag 420
atttggcltc tgaatgattt tggattttta cgtggcagcc ctcaatggcc cagattttct 480
gatgattagt aagcatttat tcttttgaat tgattattgt ctccttttca tgtgaattta 540
tactccctgt tgaaccctgt tacttaccaa taactctt 578

```

<210> 49  
 <211> 1300  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3658034CR1

```

<400> 49
agggaaatcc aattatgagg cggctaagca ggcaaaagaa tagtttgggt ggttttgcatt 60
ctactccagg taccactgtg gcttlaaagg ctgtctgaat ttgcagccct aatgaataca 120
gaaaggacaa atatccaagt gaccgtgacg gggcctagct caccagctcc tgtaaaagttt 180
ctgattgaca cacacaaccg cttaactnct cagacagcag gcttgcctgt ggtacagcca 240
acggcaggtta atatttccgc aatgggtttt ggatttgcta agacttgcta caatgttgta 300
tataatgtga aggtttcttg gtctctctga agaagaagat ctatccaaa taaagaagcc 360
tttgatttag ctgttgcctg aaagatgctc tcaatcatgt ggatttgaat 420
gtgtgtacaa gcttttgggg cccgggtagg agtggcatgg ctcttatgga agttaaacta 480
ttaagtgggt ttatgggtgc ttaagaagca atttctctga gcgagacagt gaagaaagt 540
gaatatgato atggaaaact caactcttat ttagattctg taaatgaac ccagtttllgt 600

```

atatttatca ataaagccct ttgctctctc aaaaaaaaaa a

1661

<210> 46  
 <211> 1910  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2323917CB1

<400> 46  
 gtactccggc cgcagccctg ggggtccagc ggaatctctc agaatctggg ctccagaaac 60  
 attattctaa tctgtgtatg gaaaaatccc agaaatttaa tccatttata ttgcattatc 120  
 tccaagaagt ggatgaagaa attaaaaggg ggcctagcagc agaatccacn ttaaacattg 180  
 ctggtaacaa tgccttagtg ccagtagaaa gagttacagg tgaagatttt tggatttttt 240  
 ccaaaatttt aaagaattgt ctgtatatta alggctctggg tgttggatat aaccttttat 300  
 gtgatgttgg tgcatactat gctgcgaaac tgcttcagaa acaacttaac ctcatctact 360  
 taaacctcat gtttaattgt attgggcccg aaggctggagc attgatttgc aaagtgttac 420  
 ataagaatcg gactctgaaa taccataaga tgactggaaa canaattgaa aataaggttg 480  
 gaattttttt tgtgtcaalg ctgcaattta attcatcatt agagaaatta gatctgggtg 540  
 actgtgatct gggaatgcaa agtgtgatag catttgcctc agtaactaac caaaacccag 600  
 caattaaagg cactaaacct aacagacctc tactgtacgg tgaacaggaa gagtctacag 660  
 tccatgtagg cctcatgttg aaagaaaatc actgtctgtb tgcaactaac atgtgtlaagc 720  
 ttgatataaa aaacagtggt atacaacctg tatgtgatgc actgtatctg aacagtagcc 780  
 tgcgtacctb tgcaacaaa accctggagc taactaatga tggaaatggg tatttggctg 840  
 atgtactgaa aagcaaacct accctggagc taatagatct ttcccttaac agaatagaac 900  
 atgcaggcgc aaactatctc agtgaacctc ttacttcaac caacaggagt cttaagcgt 960  
 tgcagtagt cagcaacaac atagaggag aaggacttgt tgcacttcca caatcaatga 1020  
 aaacaaatct cactttctct catatctaca ttgggggaaa caaatttgat gaggtctact 1080  
 gtatagcata ttcagaacta attcaaatgg gttgtctaaa accagcaat acagatgtgg 1140  
 agccctttgt ggtgatgga cytgtatata ttgcagaagt ctccaatggo cttaaaagc 1200  
 atbattattg gacatcaact batggagaat tatggagaa ctcatctaat gtaggttttg 1260  
 ctcttgttcc agtaggtcaa cagccatgaa aaagtatctg taaatctatt ttccttttat 1320  
 tgbtttattg tttcaaatgt atlaafatcc tatagccatb tttcttttat ttaacttcca 1380  
 aagttacttt tttcaaatgt gtaaaagata atgaactgat taaactttgt atnaacttcca 1440  
 aaaaaaaaaa aaaggggggc gcccggatta gtgagctcgt ccccggggt atttactccc 1500  
 ggggcccggc ccttggggg ggggttaacca gtctttccgc caagggttgt gtcctgtata 1560  
 tccagggctc ttggaaagag aagtcgtgct gtccttgggc tcttctcttg gtcggagggc 1620  
 atttlaactg tctgtcccgc ctggtgtact atagtaacgc caccgctgtg gtcctgtgta 1680  
 tgttggaaat aagtgtttgg tggagtctcg ctgggtgttg gaagttaggt gteectogca 1740  
 gaagtctctc tatcgggttg tggctggctg tcccgctggt gtgttgcga 1800  
 ggtagtgggt ttgcacctgt atttatattg gtccctccgc cggcgagggt ggaagcgtg 1860  
 ttggtttagt agtcctgtgc ggttgttgtg ttctcaacc cacacacaat 1910

<210> 47  
 <211> 2162  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2754960CB1

<400> 47  
 gacagcgcaa cgcactcca gctggacga cagaalaga cccgtctcca aaaaaaaaaa 60  
 aaaaaaaa aagcggaggc tggaaegtct gagcaggtgc ggggtgcaa cggcagccgc 120  
 gggagctcg ggcgggcagg ccgtctccgc ccgtctccgc ccagctccc cggcggaggg 180  
 cgtctgaagt ttctgtttcc aaacgaaagc tggggggggg gccacgagcg 240  
 cggggccaga ccaaggcggg ccgggagcgg aacttcggtc ccagctccgt ccagctccgt 300  
 gtcctgaagt ggaactcagg ctacataag atggcgcttg agagattcca 360  
 caggcagctg cggcctgggg cgttctctgg aagtgaagtc gtgtgtctcc gaacgcgggc 420  
 tggagcaaat ctgtcagctc atcttggagt gttctcggat gacgaaggcg ttgggctcc 480  
 aaaaagagga agaaagacgc aacctgaaga gtttgggttg agtcctgca gtcccgggca gatcttgag 540  
 gggactaccc tcatggcctg gttgggttac ggttggttgg gatcaggaat gtagagagcc cccaggcgct 600  
 ccttgatcca ggtgttcgta atgaalgtgg aactcaacc gatattgag aggtatccgg 660  
 720

ggcagcgatg	gatttllgggt	ggggangtcc	teettgogga	gtlgyngcca	gclagctctt	1980
gaacagctgg	atgagcaaga	tggtgatgca	gaacaaagca	acggaaagat	gaacggtagc	2040
accttaata	agatgaatc	aaaygaagaa	agaaaggaay	aggaggaatt	aatatttaat	2100
gaagatattc	tgtgtccaca	tggtgagtta	tgcatactct	aaaktgaag	agggcttgtt	2160
tctaaagagg	cltggagcaa	actgcagcag	tactttccaa	aggtccctga	gtttccaaag	2220
tacaaagagt	gctgttcaca	gtgcaagatt	ttagaagag	aagggggaag	aatgaagcc	2280
ttacataaga	tgattgceaa	cgagcaaaag	acttctctcc	caaatttgtt	ccaggataaa	2340
aacagaccgt	gtctcagtaa	ctggccagag	galacggatg	teetctacat	cytgtctcag	2400
ttctllgtag	aaaggtggcg	gaattttgtt	agaaagccta	caagatgcag	ccctgtgtca	2460
tcagttggga	acagtgctct	tttgtgtccn	caaggggggc	tcatgtttac	atttgcctcc	2520
atgacccaaag	aaagtctctaa	ecttatagct	ctcatatggc	ccagttagtg	gcaaatgata	2580
caaaagctct	ttgtttgtga	toatqlaatt	aaatctacga	gaattgaagt	gggagatgta	2640
aaccccttcag	aaacacagta	tattttctgag	cccaaactct	gtccagaatg	caaggaaggc	2700
ttattgtgtc	agcagcagag	gyacctgctt	gaatacactc	aagccaccat	ctatgtccat	2760
aaagtctgtg	ataataaaaa	ggtgatgaag	gattcggtct	cggaactgaa	tgtgaqtagt	2820
tctgaacacg	agggagacaa	ggaggaagct	aaaccagatg	gagaaaaaga	tccagatttt	2880
aatcaaatca	tgcattgcatt	ttcagttgct	ccctttgacc	agaatttgtc	aatgatgga	2940
aaagtctttaa	gtgatgactg	tgcacacctc	ggcacctctg	gcgtcattcc	tgaatctgtc	3000
attctattga	aggtgatgta	accaattgca	gattatgctg	caattgagta	tgcatgcaa	3060
gtttgtatgc	caggaagag	gtttaaaggt	actggctctc	ttggacatta	atctttgaat	3120
acttgctgac	tgttaagaaa	tgaccagagg	ggagagagg	tttgacatgt	taaggcatm	3180
aagcaaaaggt	ggatttaaga	attaaacctt	tacatgcccc	ttccaaaagg	cagaatcca	3240
ttcaaacctg	actgtcccaa	algccttatg	tcasataaag	cagattgcac	tgatggacat	3300
cagacttgaa	ggaaatgtct	ccaattttat	atttaagggg	gggtggtggg	gggagggggc	3360
aaqtaaaagac	ggaaacagct	taqtaagagt	aatagtaaat	catgtttaca	tatgagattt	3420
atagtctgtg	gagggggaata	aaqttctgtt	atatthcctt	gctogagttt	cataccagat	3480
gcttltgttc	atcaaggtat	gtatcaagta	gatgggacaa	cattctgtct	tgaacgaaaa	3540
gtaatttttag	agacataacc	tgtttaccaa	tgcctgtctt	tgattcatat	letantttca	3600
ataaagcatg	aaagtgaaga	acttgtaaaa	aaaa			3634

<210> 45  
 <211> 1661  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1973875CB1

<400> 45						
ctcagagccga	ttcggctcga	ggcggactgg	tctcggcccg	ggcaccgtcg	cgggccccc	60
tggcccgccc	acctgggaac	gtgctgggga	gtctgcccct	tccctctctc	ccctggcccg	120
caaaagtttg	gggyaccat	cgctggggct	gagcgccccc	ccggggggag	atcggggagc	180
gcccagatct	gggggggggg	agccattgac	ccgggagcgc	gcctgctctg	gagcagccga	240
ccaccccgcc	goccccggtg	catggggaet	ggctgagggg	ccagcatggg	caactgcttg	300
ggggagacag	gcccgggag	gcccggcagcc	ccgggacacc	cccgcaagcg	agcaggagcg	360
aatgagccccc	tgaagaaaga	gcggcttaag	tggaaagagc	actaccocat	gaatgaaggg	420
cagctgcggga	gcaaacggga	tgagtctctg	gacacagcgc	ctgcccctga	gggcccgaag	480
gagatctggg	atgcctccaa	ggtgtccggc	tatgctgctg	aaqccaaag	ccacagctcg	540
gcccaggcca	tccctggatg	agccagcacc	acctgcccct	atggcaccct	ctgtgaatgc	600
taagatgagc	tgggcaatcg	ctaccagctg	cccatctact	gcctgtacc	gcccgtgaac	660
ctgctgctgg	agcacacgga	ggaggagagc	ctggagcccc	ccgagccctc	accagagctg	720
cgccgtgaat	tcccgtgaa	ggtgcgctcg	tccacgggca	aggacgtgag	gctcagcgcc	780
agcctgccc	acacagtggt	gcagctcaag	aggcagctgc	acgcccagga	gggaatcgag	840
ccatcgctgg	agcgtggtt	cttctccggg	aaqctgctca	cagaccgcac	acggctccag	900
gagaccaaga	tccagaaaga	ttttgtcatc	agggtcatca	tcaaccagcc	cccaccacc	960
caggaactgat	gggcccacgg	acccctggga	agagggcccg	ccctggagca	taggcccaca	1020
ccctgctgct	gccttccagt	gctgtcattt	tcttcagggg	ccctccccc	ggtgtggctg	1080
gtgggtgagc	cgtaagggga	ccctgctctt	aaqggcacta	cgcccccaca	gttccgggta	1140
cccaggggagc	aggcagccac	acacgggccc	tgcacacttg	tcaagaaaaa	ggcgaacag	1200
gcccacccc	tgcctgtctc	ccgaagagag	tlcagagccac	aaqggccccc	caagagggcc	1260
ctggaagccc	gctctgtcat	ctgggtgctg	cagctgtgct	cactctgggt	ttctgctcag	1320
ggtctgaagc	agctgctgtc	tcccctctct	gcccccatcc	ccctggctctc	ccctggggac	1380
agtgccactc	ccctgggaag	gaggggaacca	cccgtyagcc	ccagggtctg	ggagagctga	1440
ggcggggcctc	tgcctctccc	tgcctccagc	aaacttggca	gaatagagag	aaqtaagggg	1500
cggttgggct	gtcgtggcag	ggtctgcaca	ggcccatgtc	ctggctgtaa	ccagggcagt	1560
gggaggtctg	cagcctggctc	atggccctcc	cagcaggtcc	ctgtggagag	acatctctgc	1620

&lt;400&gt; 43

gcacaaatltg	tttcaaaaaa	tttgtgagga	ttttgattcc	tatctcgttc	naattttttgg	60
ttcatcgtca	cctcaacottg	gattttgggat	gtcagggggaa	gaactcagtg	aatctactcc	120
agagccctcaa	aaagaaatttt	ctgagtcatt	gaptgtccac	agagaccagg	atgaagatga	180
taaggctcca	gagcccaactg	gggctgatga	cttgccctgcc	actacotcat	ctgaggccac	240
caacacccccc	aguccactgc	tcagcaacccc	tytggatggg	gcagagggatc	ctagatgttt	300
ggaagccttg	aagcctggaa	actgtggtga	atatgtggtt	cgatgggtatt	atgacaanca	360
ggtaaatct	tgtgcccgat	tttggttcag	tggclgtaat	ggctcaggaa	ctagattcaa	420
cagtgaanaag	gaatgtcaag	aaacctgcac	tcaaggatga	gcaagttaat	tggcctgtct	480
ctatcaaaag	catagaacclt	cctaatllcc	acatattccr	ccaatacaca	tacagcacta	540
tattttgagt	tatactgagt	atttacaact	tatacatgta	attgaattct	caclacagcc	600
ctaggatgta	catattatta	accacttata	tagytaagaa	agctgaggct	ctgagaagtt	660
tagtaacttg	tcaactgtca	cccaactaaa	aaqtttcaqa	gctgaggatt	tagacttaga	720
gctgtgttaa	ttcaatacau	agactctate	tacttcaaaa	cctgcaatgt	gattctgatt	780
cttttaattc	ctgtttgatg	tactatgtca	gtccaaaacc	ctacccctgt	ccttgcccat	840
acctcaaccc	actcaacctc	ctaacctctc	tatgtccctc	acagtaccaa	gatgtaggtg	900
atagggaagg	cttcgggtgtg	agaatttaga	atgatgtaaa	tggttaacga	ggagtycttg	960
gataaggatc	gggatggtga	gggtagttag	atttttgect	cacttgccct	gaaagtgtta	1020
ataggagaa	accaatctga	attacaatta	cttaaatgta	tcaagactgt	tcacttttga	1080
ttctcccaac	atgttttgta	acaagtgttt	aatgtatgtt	aaaataaaga	aggtttttat	1140
acccctccat	taaaatlatgt	cagtgggccc	ttccatttta	tygagtygaa	tgggaaggcc	1200
cttgacagcc	aggaaccact	tgaagctggc	atccaactct	gaacagagag	tattaaagac	1260
aggattccca	ctgaanaagt	egccaccaaa	ttgcaatggc	agtaatggag	agtaacggag	1320
aatcgtctga	ccttggggaag	cagaggttgc	agagagccgg	gattgccccg	g	1371

&lt;210&gt; 44

&lt;211&gt; 3634

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Inocyte ID No: 1911808CB1

&lt;400&gt; 44

gggggggag	tgggggggac	aggtgattga	ctggccagct	gactgaagga	goyccaggtc	60
ctccttfgctg	gcaggttggcg	aagccccctg	gggggggggt	gcagaccggc	gggggggctg	120
ggggggtctg	gtccgggagg	cgttccclgg	gcccaggcca	tygcccngcg	gtgtgcagctg	180
gagcaggcgg	cctggcgctg	ggcggagacg	gtggggcccg	aggagggtgc	gcaggagcat	240
atcagagccg	nttaacggcat	ctgggtggag	ccttgccttc	guyggtgtgtg	cagagcgaac	300
tgcaaaaggaa	atccgaattg	cttgggttgt	atctgtgagc	atatttggct	agggagaaata	360
gatgaaaata	gttttcaata	catcgatgat	cccaactgtg	aggggggaaa	aaagaactca	420
tttgtgggoc	tgaactaacct	tggagccact	tgttatgtca	acacatttct	tcaagtgtgtg	480
tttctcaact	tyggagcttcg	gcaggcactc	tacttatgtc	caagcacttg	tagtgactac	540
atgctgggag	acggcatcca	agaagaaaata	gacttatgag	ctcaaaacaat	tttgtggcat	600
ctccagtact	tgtttgcctt	gttgcaaaac	agtaatagga	gatacattga	tccatcaggga	660
tttgittaaag	ccttgggccc	ggaactgtga	caacagccagg	atgtctcaaga	atttccaag	720
ctcttttatgt	ctctattgga	agataactttg	tctaaacaaa	agaatccaga	tgtgcgcaat	780
attgttcaac	agcagllclg	tyyayaatat	gactatgtaa	ctgttttgcaa	ccagtgtggc	840
agagagctca	agcttttctg	aaaattttat	gagctggagt	tcaatatcca	agggccacaaa	900
cagttaacag	attgtatctc	ggaattttttg	aaggaagaaa	aattagaagg	agacaatcgc	960
tattttttgcg	agaactgtca	aagcaaacca	aattgcaaaa	gaaagattcg	acttctcagc	1020
cttcccltga	ctctgaactt	gcagctaattg	cgttttctct	ttgacaggca	aactggacat	1080
aagaaaaagc	tgaataccta	cattggcttc	tcagaaattt	tggatatgga	gacttatgtg	1140
gaaatataag	gtgggtctctc	cgtgtatgaa	ctcagccag	tccatataca	cagaggagtg	1200
agtgcttatt	ctggccacta	catcgccccc	gtgaaagatc	cacagctctgg	tgaattgttat	1260
aagtttaattg	atgaagacat	agaaaagatg	gaggggaaga	aattacaact	agggatttgag	1320
gaagactctag	cagaacottc	taagtctcag	acacgtaaaac	ccaagtgtgg	caagggaaact	1380
callgtcttc	gaaatgcata	tatgttygtt	tatagactgc	aaactcaaga	aaagcccaac	1440
actactgttc	aagttccagc	ctttcttcaa	gagctggtag	atcgggatag	ttccaaatttt	1500
gaggagtngt	gtattgaamt	ggctgagatg	cgtzaagcaa	gtgtggatag	aggaaaagca	1560
aaacacgaag	aggttaagga	gctgtaccaa	aggttaactg	ctggagcttg	gocclatgag	1620
ttgtctcttc	tggaaatggcl	gcaaaagtgg	ttggatgaat	caacacctac	caaacctatt	1680
gataatcacg	cttgccctgtg	ttcccatgac	aagcttcacc	cggtataaat	atcaatttatg	1740
aagaggatat	ctgaatatgc	agctgaactt	ttctatatga	gatlatggag	aggtccaaga	1800
ctaactytga	aagccctgtg	taaggaatgt	gtagtagaac	gttgctgcac	attggtctctg	1860
aagaaacaaa	laaatgaaga	ttataaaact	gttaataatc	tycctgaagc	agcagtaaaag	1920



```

tgttttaaca aaagagtgtg gacattttta ttttanaatt taggcaaaag tcaactatcaa 1660
atgggttgctt atttgttice cacagccata tagtttttcc tggagggttt tgttttgyttg 1740
ttgttgaana gattttgtct aagcttagat gaaactttct atagaaaaaa aaaattgtttg 1800
aaaggtccag ttctcagtac catgtgagtt aatgatacta aaactaagtt ctttttaaaa 1860
agtgaattat gtattttata aattaccitt tcacatatgc aaactctgtt tctactacaa 1920
tgttattttt actaatgect battgttgna ctctttttga aacatctctg agtgaatata 1980
tgaatcaatt tgggcttaaa actgaaagcc agttgggctg aaagggtttga aatacgtacc 2040
coagtaaaac cattcaatcc ataatttgta aataatcttt taasaatggt tttaatccgt 2100
atagatgaca tttttagact ttggacatgg tggtaattaa ggggcattaa ttltacactc 2160
caaggtz

```

<210> 42  
 <211> 1826  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1256390CB1

<220>  
 <221> unsure  
 <222> 1755  
 <223> a, t, c, g, or other

```

<400> 42
ccccggcctt tctatagcat agttctcttt aggtgggaatg attcctatna gatttctcat 60
tattaaatca tgcatttttc aagatggnaa caatcttlln tttatctaa gctgatattc 120
tcatttgytt gaagaacac acacatgcta gagagagagg aggaatatata cccacgaccc 180
cacagccagt tagtalccay ttggtgctgg actccagcca ggttctctgc ctcaggttag 240
ctaaatgata tatagaaaag gtaaattttt aaagaactat ttalttaalt atlccataaa 300
aacallttta aggtAACCAc ttaaaaaatg ttaatttttc cattccaaag taaatgotaa 360
gcattgtttat taatgaagca gtactctctg ttagtatatg acattctgaa gtttaataaa 420
ctcaattgca taaatgtgtc tctcttggtt tagtggaggga tttagaggatt ttttggggat 480
gtagaggtgt gatattgtag atccttagcc caacacagac aagagattgt ggcaggtcc 540
ggaggtctgg gtagcttggg gactattcc tatcacctat accaccccat ggcagagato 600
gtgaatccat ggagcttggg cagtgagagat taccaggaaq tggtagacaa ggcattctct 660
tatgagtgga tgagagagat cagtgagaaq ccccatgtat tatctgaaga tcagccaacc atctggtaat 720
ggagtgaact atgagacccc ggaactgllga attcagccca gagaatggat tgcctctgct 780
cccaagaaaa tcaattggat agaaattcta caaaaccata aagacaactc aagtataccc 840
ttttgcraat ggttcgtcaa ottctatgto totttggagg tggaaetttc aatgcctctt ggtgtagtat 900
aagctccctt ggaactggg totttggagg aatgcctctt ggtgtagtat tgtgttctct 960
tacacttggg caactgatcg tggaaetttc attctgtggg acagggccag tgtgttctct 1020
tgtlltggga ggaatcnaa agaatcaaac gagcaagaag gatgatattt tttgttctct 1080
agaaactgoc gtttcataga agttaattct acacacttac ggcacacca aaatataatc 1140
gctgtttgca gtttcataga agttaattct acacacttac ggcacacca aaatataatc 1200
tcttatgggc agttaattct acacacttac ggcacacca aaatataatc 1260
caggaatgga ttcaagttgg tggatcgag tgcagatatt tctatgctt gctcaggttc 1320
aattatagag ttggatcgag tgcagatatt tctatgctt gctcaggttc agctgaggga 1380
tgggcccagag acattgggat tcccttctca agctcagatc cagccacact gtagggagac 1440
tatgggtttg ttctgcccga tcttgatga tcttgatgag aacactggc actcggaacg 1500
gtgctgtcag tcttgatga tcttgatgag aacactggc actcggaacg tctctaatgt 1560
gtgacatctg ccactatgct aggttttttt ctgcccagaa ttttctctcc ttttctctcc 1620
cattctgccc tttctttgat actaattctg acattacctg tggaaatttt gctgacttcc 1680
tttctctctc tctcttaact gtttctacac actttaatgt aataatttac 1740
agtgaatag gattttacat gcccctg

```

<210> 43  
 <211> 1371  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1786774CB1

```

tccgtacaga taaaggaaga gatccaggtt ctgatanata tyaggeagat tctgagttgt 900
tgctccagat acgaatggt gtgcttgact caotgggtat taglccotgac ctgcttccly 960
aggactttgt caggtactgc ttctccyaga tggccctagt gtgtgcyytg gttggaggga 1020
ttttggcaca ggaattgtg aagccctgt ctccagggga ccttccctac aacaccttct 1080
tetctcttga tggcctgaag gggcaatggga ttgtggagt ccttgccccc aagtgaacte 1140
aagatttggc agccccagag atgccaactg cagcatgccc acctgtalte cctgtcccc 1200
tccttcatga agcatctcc gtgcccaggt gtgcccaggt gctgcttccc atccaccgca yctgctcgac 1260
ttctgcacac gaaggaggtg gtgcccaggt gactgttcca ggctgctgt atcccccggc 1320
aagggggcga ggggtggtgt ctttgttcca gaactgttca ccttgcctgt cctgtcctt 1380
ctgcccagct ccttgagtg ggcctcgcca gacctgtggg ccttcttggc attgtcctt 1440
atgctgtccc ggcctcgcca gacctgtggg ccttcttggc attgtcctt caggaatgag 1500
caagctctgt tgcctgggag ccttcttggc attgtcctt ccttcttggc attgtcctt 1560
ttatcgagaa aagtgtgaat tcaggtggag agtaggccc ggccccatga ggcaccagtg 1620
gaagcaccgc tccaggttca gacaggtgac cttagagagg aaaaaccatg caggcaaatg 1680
catttctctt ggagtttgag acctgacaa accacaggtg gcattctgtg tgnlytlett 1740
gagtttctgt ttgagttcca ttgagttcca gctgggtttt gggagaaagg agatgtctac 1800
aagttcttga tgttagggcg agacctgca agttgagtat tagagagctt gtcttctaat 1860
gcaggttctt ggggcttcag ggttaggagg ttgaggttcc aggtatgtt cctttttaac 1920
tcacatcggt ctccagtcac tccagaggtg tccatttca ccttcttctt ggtctatgtt 1980
ttactctggg aaccacagct gtttgtgttt cctcatgtt ttgagtttcc gtgaatttct 2040
tttaaggaa tgataataaa gttacttgtt accagttgac ttttccacc ctttcttctt 2100
gaagcttctg agaggggaat ggtatgacct ccttccacc tctggttcca ttatataata 2160
actccagaga tcttctctct ccttccacc tctggttcca ttatataata ttatataata 2220
caagagttca attgttccac ggtatgttcc tctggttcca ttatataata ttatataata 2280
ggataattca gttatattgt ggtatgttcc tctggttcca ttatataata ttatataata 2340
tcaaaagtga tgcctcatat ggcctcagtg ccacagccgc agttcaccct 2400
ggagagctag gtttgccttg atgcagccaa tcttcttctt ccttcttctt 2460
gtgttctctc tttatgttaa ttgtgtgaly tcttcttctt ccttcttctt 2520
aa

```

&lt;210&gt; 41

&lt;211&gt; 2167

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 411344CB1

&lt;400&gt; 41

```

cgccggtctt ggggggatgg ttccatcatg gcgtcaatgc agaaacgact acagaaggaa 60
ctgtttgctt tgcanaatga cccacctctt ggaatgaact taaatgagaa gagtgttcaa 120
aattcaatta cacagtggat tctagacatg gaaggtgcac caatgagaa alataaggga 180
gaanaalltc aacttctatt taattttagt agtcgatata cttttgactc tctcaggtc 240
atgtttactg gtgaaaatat tctgttctat cctcatgtct atagcaatgg tcatatctgt 300
ttatctattc taacagaaga ctggctccca gctctctcag tccaatcagt ttgtottago 360
attattagca tgccttccag ctgcaaggaa aagagacgac caccygates ttctttttat 420
gttgcgaacat gtaacaagaa tccaaagaaa acaaaatggt ggtatcatga tgatacttgt 480
tgatgcccac gttatcalcc tccaaagaaa agctaglyctt actyagaaaa ttagcacctt 540
gactattcag tctttgaact ttaacctttg actggaagtg acctatagge atgagagact 600
acttcccttt actgcatttt tacttctgtg cttcttgggc ccatgttget gcaggttget 660
gtccaggcaa ctgacatgct tttattagtc atacagtatt aatgcaggtg tcttggaaaag 720
tcaaatatca ttcactcttt tatttttatt tttttaaget tttggaaaag ctccaggtcc 780
tcatgtattg tgcaataaca ttgacttctt tggcggtttt ggtacgltca ttgcaggcaa 840
tggngtltgt aacaygaaaa gtttctatta actcttgcca ttcaatgatt ttgcaggcaa 900
agggcctatg aatgaactt actgattata gtgggaatat aatataagtg agggalccaa 960
cattacttct aagllcaccct caactgttta tttttggatt ctatgcactg tgatccttaag 1020
gttaacagca ttttgaattg acttcaagtt gttttgacat atgaaagato attgnaattt 1080
tggcagagag ttatccctat ggtgaaata ygaagatttt caagtaacat 1140
ttctccaatc tctgaagtta aggtatgagt ggtgaaata aattaatttt ttgtatatag 1200
tttatttttg aagtaagaaa aatggcagtg aatataaggt atyaatggtt taatcaaggc 1260
gottgttatt tgtacatlaa aatggcagtg atgattaaat tgotgcagtc cataatttgg 1320
tgcacatcag catgagccct agatcttctt caagttaagt aactctctgc 1380
cogtaaacat tttgttttgc aatcttcttca cttatttctg ttggnaacc 1440
aatactttta ttgaaacttt tgtaagtttt ctttctgtat tgaatgttcc 1500
tcatcttctc ctttaactct gtaacctaaa atagaaata cttttttgac attgtgagta 1560

```

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5432879CB1

<400> 39

gtccggccgc	ggcggcgccg	gtagcggagg	agacgggttc	aggccctccg	tgccggtgca	60
atgctgagct	cccgggccga	ggcggcgatg	acccggcccg	acagggccat	ccagcgcttc	120
clcgguaccc	ggcggcgccg	ccgatatata	gtcatgaaga	actggggagt	tataggtgga	180
attgctgctg	ctcttgcagc	aggaatatat	gttatttggg	gtcccatcac	agaagaagag	240
aagcgtagaa	aagggcttgt	gcctggcctt	gttcaattag	ggaacacctg	cttcattgac	300
tccctgctac	aagggctgtc	tgctgtctct	gttttccata	ggtggcttga	agagttccac	360
tccagtaact	ccaggatata	gaaggagccc	ccctccaccc	agtatttata	cttcaacatc	420
ttgcaccttc	tgaagccctt	gtcctgcca	gaagttactg	atgatgaggt	cttagatgca	480
agctgcttgt	tggatgtctt	aagaatgtac	agatggcaga	tctcatcatc	tgaagaaacg	540
gatgctcagc	aattattcca	tgctcattacc	tgctcattgg	aagatgagcg	agaccgccag	600
cctcgggttc	cacatttgtt	tgaatgtcat	tccctggagc	agcagtcaga	aataactccc	660
aaacaaatta	cctgcgcgac	aagagggtca	ccctccccc	catccaatca	ctggaggtct	720
caacatactt	tctcattgga	actcactagt	aatatgtctt	gcaaacactg	tgaacaccag	780
agtcctgttc	gatttgatcc	ctttgataga	ctttccactaa	gtattccagc	gtccacactg	840
ggtcacccat	tgaccttggg	ccactgcctt	caccacttca	tctcatcaga	ctcagtgcgg	900
gatgttgtgt	gtgacaaact	tacaaagatt	gaagccaaag	gaacgttga	ccgggaaag	960
gtggaaacac	agaggaccac	ttttgtttaa	cagttaaaac	taggggaagt	ccctcagttt	1020
ctctgcatcc	acctaacgag	gclgaactgg	ccagggcagc	gcaagcctct	gaagcggcat	1080
gagcacgtgc	agttcaatga	gttccctgat	atggacattt	acaagtacca	ccctccttga	1140
cataaacctc	gtcaaacaca	ccctaaactg	aacaaagacc	cagggcctac	actggagctg	1200
caggatgggc	cgggagcccc	caacccagtt	ctgaatcagc	caggggcccc	caaaacacag	1260
atttttatga	atggcgccctg	ctcccatctt	ttattgccaa	cgtgtgcagc	gccgatgcc	1320
ttccctctcc	cagttgttcc	cgaactacgc	tcctccacat	acctctcccg	gctgatggca	1380
gttgctgtcc	accatggaga	catgcactct	ggacactttg	tcacttaccc	acggtcccca	1440
ccttctgcca	ggaacccctc	ctccactagc	natcagtguc	tgtaggtctc	cgalgacact	1500
gtccgcaagg	ccagcctgca	ggaggtccct	tcctccagcg	cctacctgct	gttctacagc	1560
cggctccttt	ccaggatgca	gcaaccagcg	caggagtgc	agtctgaaga	atgaactgtc	1620
cctcctgcaa	ggctagagct	gatggcactg	tcctcactgt	ccaggcaaaa	agtcacactg	1680
tactgttgcc	tgtcgaagcg	gcccacttag	agccttccag	ccttctgggt	tgttctaaag	1740
gcaggctcca	ccggggagcc	agccccagt	caacacaaac	caggtcctcc	gaacagtctt	1800
gtctatgtct	gtaggttggt	ctgttgtgtt	aagaagcat	tcattatgtc	cggagtgtct	1860
ttttactcct	ctgatacagg	taattaaaag	aatcagatt	cttgaagcca	ccgttttcat	1920
attgtaactg	taggtgttct	cagaggggag	gtacctttgt	ctcatcaacg	tttccactta	1980
gactttttat	ttttaataag	cagggccata	aaaattgttg	acaagaatta	atgaatttat	2040
taaaaggaaa	aattttagag	aaaaagtgc	tttcaacttc	gatttcttct	gtagcacgtc	2100
cattgtgaaa	tattccttcc	aggctactca	aaggatagca	agagaacagg	taaatgatgc	2160
ctaaagaaca	ccttcccttt	ctatag				2186

<210> 40  
 <211> 2522  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5853753CB1

<400> 40

ctccggggct	gctgcggcg	gggttaggtg	gcgcgggggt	ccggcggggg	gttggcttga	60
gagggaacgg	agctgagaca	ggagagagcc	ggccatgggt	ggagaaggag	gaggctggcg	120
gcggcattag	cgaggaggag	gcggcacagt	atgacccgca	gataccgctg	tggggactgg	180
agggccagaa	acggctggcg	gcctctcggg	tgcttcttgt	cggcttgaaa	ggaacttggg	240
ctgaaaattg	caagaatctc	atcttggcag	gagtgaaagg	actgaccatg	ctggatccag	300
aaaggttaa	tccagaagat	cccgagagct	agtctctgat	tcgtactggg	tctgttggcc	360
gaaatagggc	tgaagcctct	ttggagcgag	ctcagaatct	caaccccatg	gtggatgtga	420
aggtggacac	tgaggatata	gagaagaaac	cagagtcact	tttccctcaa	ttcagatgtg	480
tggtcttgac	ttgctgtctc	agggatgtca	tagttaaagt	tgaccagatc	tgctacaaaa	540
atagcatcaa	gttctttaca	ggagatgttt	ttggtacca	tggaacaca	tttgccaaac	600
tayggagayca	tgagttttga	gaggagaaba	ctaaagtgtc	caaagttagc	caaggagtag	660
aagatgggac	cgaacccag	agagcaaaac	ttgattcttc	tgagccaacg	atggtcaaaa	720
agaaggttgt	cttctgcctt	gttaagaag	ccctggaggt	ggactggagc	agtgagaaag	780
caaaaggtgc	tctgaagcgc	acgacctccg	actactttct	ccttcaagtg	ctcttaaggt	840

tggtgaataa	tcagatata	atgagacaaa	ngttggaaat	tgccagggaat	ccaggcaatga	1080
tgcaaggagat	gatgaggaaac	cagggaacag	ctttgagcaa	octagaaagc	atcccagggg	1140
qatataatgc	tttaaggagc	atgtacacag	atattcagga	accaatgctg	agtgetgcac	1200
aagagcagtt	tggtggtaat	ccatttgcct	cccttggtgag	caataacatcc	cttggtgag	1260
gtagtcacac	ttcccgtaaa	gaaaatagag	atccactacc	caatccatgg	gtccacaga	1320
ctccacagag	ttcatcagct	tcacagcgga	ctgcagcga	tgtyggtggc	actactggtg	1380
gtactgcag	tgacactctc	gggcagagta	ctactgcgc	aaatttgggtg	ctggagtag	1440
gagctagtat	gttcaacaaa	ccagysatgc	agagnttgtt	gcaacaaata	actgaaagc	1500
cacaacttat	gcacaacatg	ttgtctgccc	ctacactgag	agccatgatg	cagtcactaa	1560
gcacagaatcc	tgaccttgc	gcacagatga	tgtgaataa	ccccctattt	gtgggaate	1620
ctcagcttca	agaaacaaatg	agacaaacagc	ccccactttt	octccaaaca	atgcagaate	1680
ctgatacaact	atcagcaatg	tcaaacctta	gagcaatgca	ggccttggtta	cagattcagc	1740
aggytttacc	gacatcagca	acggagagccc	caggccctcat	cccagggttt	actccctgct	1800
tgggggaact	aggaagcaact	ggaggtctct	caggaaactaa	tggtatcaac	gccacaccta	1860
gtgcaaacac	aggtcccaaa	gcaggaaccca	ctgaaactgg	aatccagcag	tttattcaga	1920
agatgcttca	ggctcttgc	ggagtaaatc	ctcagctaca	gaatccagaa	gtcagatttc	1980
agcaacbaact	ggacacacctc	agtgcacatg	gatttttgaa	ccgtgagcca	aaactgcaag	2040
ctctaataagc	aacaggaggt	gatatcaatg	cagctattga	aagggttactg	ggctccacgc	2100
catcatagac	gcatttctgt	attctgaaaa	aatgttaattt	atttttgata	agggcctta	2160
aaacttataaa	tacctgcttt	atttcatttt	gactcttggg	attctgtgtc	gttataaaca	2220
aaaccaatat	gatgcatttt	aaggtggagt	acagttaagat	gtgtgggttt	ttctgtattt	2280
ttcttttctg	gaacagctgg	aattaaggct	actgcactgca	tcacttctgc	atttattgta	2340
atttttttaa	aacatcacct	tttatagttg	ggtgaccaga	ttttgtccctg	catctgtcra	2400
gtttatttgc	tttttaaaaa	ttagccctatg	gtatgaaatt	atgtagaata	aaagcattaa	2460
aaagaagcaa	atcattttgca	ctctataatt	tgtggtacag	tattgcttat	tgtgactttg	2520
gcctgcatttt	ttgcaaacaa	tycttgaaga	tttatactac	tgataatttt	gttttatctg	2580
latacaatat	agagtatgca	catttgggac	tgcatttctg	gaaacataact	gcaataggct	2640
ctctgagcaa	aacacctgta	actaaaaaag	tgaagataag	aaaatactct	aaagctgag	2700
tatttccctaa	ttgtatagaa	tcctaacgca	tccttgacaa	acatctccca	gcaaaagtgc	2760
cggttagtca	ggtttgttga	aaatacagta	gaaaagctga	ttctggttat	ctctttaagg	2820
acaattaatc	gtacagacac	ataatgtaac	attgtctcaa	catctattca	cagattgact	2880
gtaaattact	ttaatctttg	tgcagactga	aggaacactg	tagtatccc	caaagtgcac	2940
ttccttagga	cttctcagct	tctccactga	gtagtctaac	aggtattaaa	atttctaat	3000
gaaatgttgc	tttcaactgaa	aaagtgtctt	gabgttccag	ttatttttaa	tcgcaataaa	3060
aaaatagaac	tatcttttgg	gttttatctgt	ttctcactgc	acaggcaata	cacaatttta	3120
aaatgagctg	tgagccactt	gtttctgga	gggttggga	gltctattaa	gaatatagta	3180
aatatttgtc	tttccagagc	ctcagagaca	ggggagcgcg	gtgcgggggt	ggggcagcgg	3240
aatctgtctc	ggatggyggc	agcttaataa	atcttgccaa	cnaagat		3287

<210> 31  
 <211> 2412  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1299481CB1

<400> 31						
gggtgtgtgt	cggtgggggc	gggucgggac	gaggggagatg	gagcgagagc	cgagccgggt	60
cagagttgaa	caatgaccat	agttgacaaa	gcttctgaat	cttcagccc	atcagcttat	120
cagaatcagc	ctggcagctc	cgagucagtc	tcacctggag	acatggatgc	aggttctgcc	180
agctgggggtg	ctgtgtcttc	attgaatgat	gtgtcaaatc	acacacttcc	tttaggacca	240
gtactctggtg	ctgtagtcta	ttcaggttca	tblgtacctg	ataaatcaaa	acctcacc	300
caaaaggatc	aagccctagg	tgatggcacc	gtcctccac	agaaagtctc	tttcccatct	360
gagaagattt	gtcttaagtg	gcaacaaaact	catagagttg	gagctgggcl	ccagaatttg	420
ggcaataacg	gttttgccaa	tgcagcactg	cagtgtttaa	ctacacacc	acctcttgcc	480
aattacatgc	tatcacatga	acactccaaa	acatglaaly	cagaaggctt	ctgtatgatg	540
tgtacaatgc	aagcacatat	taccacaggca	ctcagtaate	ctggggagct	tattaaacca	600
atgttttgc	tcaatganat	ggggcgtata	gclaggccac	tcogttttgg	aaaccaagaa	660
gatgcccatg	aattccctca	atacactggt	gatgctatgc	agaaagcatg	cttgastggc	720
agcaataaat	tayacagaca	cacccaggcc	accactcttg	tttctcagat	atttggaggc	780
tacctaagat	ctagagtcac	atgtttcaat	tgaaggggcg	tttcagctac	tttctgacaa	840
tatcttgata	taacattggc	gataaaggct	gtccagagtg	tcaacaaggc	attggagcag	900
tttctgaaag	cggaacagct	tgetyggaa	aactcgtaca	agtgcagcaa	gtgtaaaaag	960
atggtlccag	cttcnaagag	gttcaactatc	catagatcct	ctaattgtct	tacactttct	1020
ctgaaacggt	ctgcaaat	taccygtgga	aaaattgcta	aggatgtgaa	alacccctgag	1080

&lt;400&gt; 29

```

gtgcttgaggc cctgcgcctgg acgactcggc cggtagtgga gatgtccggc cggctctaagc 60
gggagtgctcg cggttccact cggcgggaagc gagagtcctga gtcgcggggc agctccgggta 120
ggcgcgcagcg ggagcgagat ggagcgagat ggagcgagat ggagcgagat ggagcgagat 180
cgtgcgcctgt gaagcgggag ggagcgagat ggagcgagat ggagcgagat ggagcgagat 240
tcccgcttctgt ggagcgggag ggagcgagat ggagcgagat ggagcgagat ggagcgagat 300
agggtgcgagc aaagaatggc cggagtgagat cggagtgagat cggagtgagat cggagtgagat 360
acctgggcacac cattacacagc agtggtgctgg agtggtgctgg agtggtgctgg agtggtgctgg 420
ccctctcaca catcaatgct tatgctctgt attcagctgt attcagctgt attcagctgt 480
gtttgaagtc tccagcctac attcagctgt attcagctgt attcagctgt attcagctgt 540
tccacacccct caagttttac tgccttccag agggccacat tgccttccag agggccacat 600
aggatatcac gttatgtgtg agggccacat tgccttccag agggccacat tgccttccag 660
agcaagccaa attgtccggg gatttatgca ctggaagaag gtcacagcgtt gtcacagcgtt 720
tgaaacacat aaaggccaa ctggaagaag gtcacagcgtt gtcacagcgtt gtcacagcgtt 780
ctcctctccg gaactacttt gttcttctgt caaggcactt ttttcagatc gcaactcagct 840
gggatatcat gttcttctgt caaggcactt ttttcagatc gcaactcagct ggggtccatg 900
ctcgaaatbt caaggcactt ttttcagatc gcaactcagct ggggtccatg agagcagctg 960
tgaatgcctb gcaactcagct ggggtccatg agagcagctg agagcagctg agagcagctg 1020
atgttttcca ggggtccatg agagcagctg agagcagctg agagcagctg agagcagctg 1080
caggaagaaa agagcagctg agagcagctg agagcagctg agagcagctg agagcagctg 1140
agtcacacat lcccccagtg taagacttac aatcttttgt tgtcaatttc aatccacatg 1200
zgaaggaata taagacttac aatcttttgt tgtcaatttc aatccacatg aatccacatg 1260
ctccatctct aatcttttgt tgtcaatttc aatccacatg aatccacatg aatccacatg 1320
atccacatct aatcttttgt tgtcaatttc aatccacatg aatccacatg aatccacatg 1380
aagtaacagc agagcagctg agagcagctg agagcagctg agagcagctg agagcagctg 1440
gcaagccctc agagcagctg agagcagctg agagcagctg agagcagctg agagcagctg 1500
atgaattaca agagcagctg agagcagctg agagcagctg agagcagctg agagcagctg 1560
cttacattca gatttggaag agggccttgc tgaacacagc accagagcac tgatcatttt 1620
ggagcgctgc agggccttgc tgaacacagc accagagcac tgatcatttt cagcagggca 1680
aaagcttagc accagagcac tgatcatttt cagcagggca cagcagggca cagcagggca 1740
lbtggtctac accagagcac tgatcatttt cagcagggca cagcagggca cagcagggca 1800
agctgttctt cagcagggca cagcagggca cagcagggca cagcagggca cagcagggca 1860
ctgtgagtag ttgaagaaca ggcagtaact tttgtatttt tagatggatt 1920
caagcagaaa ggcagtaact tttgtatttt tagatggatt tagatggatt tagatggatt 1980
catttccctga aaaaaa 2040

```

&lt;210&gt; 30

&lt;211&gt; 3287

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1261376CB1

&lt;400&gt; 30

```

cagggcgctg acccgcgctg ctcgcgcggc cgcgcgcgaa gggagcgtgg ggaaggggca 60
gggagggagga agcggtgctg gctgcggatg tgggtgtgag cggagcggcg ctgaacacac 120
ggcggctgac gagcgctga cccgggcctg cggcagagcc tgcacccagc tccggggccc 180
cacacccgct accggtgccc tgcgcgcgtt gctactgagg cggcgtgctc tgcattcttc 240
gctgtccagg cctgcgggct ctgggtgtct cttggtcctc cttgcctccc tgcctccctc 300
tgcttgcttg agtcacccgc ggcgcgcgcg ccacagccat ggccgagagt ggtgaaagcg 360
ggcgtctctc gggctccagc agggagcaga aggttactgg gcccccggg gcccgcggg 420
ccgctgcctc cgggagcgc aaaatcatga aagtcacagt gaagacccc gaagaaaaag 480
aggaattcgc cgtgcccagc aatagctcgc tccagcagt taaagaaaga atctctaaac 540
gttttaaatc acatactgac caacttgtgt tgatatttgc tggaaaaatt ttgaagatc 600
aagatacctc gattcagat atggacttac ttttcaact ttctcaact ttgtaataa 660
cacaaaaagc gctcaggat cttcagctc agcaaacaaa tacagcttga agcatttga 720
ctacatctac aactctant agtaactcta catctggtc tgcctactag aaccttttg 780
gttttaggtg cettggggga cttgcaggct tggtagctt ggggtttgat actacacac 840
ctctctgaat ccagagtcag attcttttgc aactcttcaa tccctgactg atgatggct 900
agatcatgga aaatcccttt gttcagagca tgcctctcaa tccctgactg atgatggct 960
caattatggc caatccacaa atgcaagcgt tcatcacaga aatccagaa attagtcata 1020

```

Asp	Gly	Ile	Asn	His	Asp	Val	Thr	Thr	Ala	Trp	Gly	Gly	Asp	Tyr
			650						655					660
Trp	Arg	Leu	Leu	Thr	Pro	Gly	Asp	Tyr	Met	Val	Thr	Ala	Ser	Ala
			665						670					675
Glu	Gly	Tyr	His	Ser	Val	Thr	Arg	Asn	Cys	Arg	Val	Thr	Phe	Glu
			680						685					690
Glu	Gly	Pro	Phe	Pro	Cys	Asn	Phe	Val	Leu	Thr	Lys	Thr	Pro	Lys
			695						700					705
Gln	Arg	Leu	Arg	Glu	Leu	Leu	Ala	Ala	Gly	Ala	Lys	Val	Pro	Pro
			710						715					720
Asp	Leu	Arg	Arg	Arg	Leu	Glu	Arg	Leu	Arg	Gly	Gln	Lys	Asp	
			725						730					

&lt;210&gt; 28

&lt;211&gt; 2080

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: O88715CB1

&lt;400&gt; 28

tgaaggactt	tccagggacc	caaggccaca	cactggaagt	cttgcaagctg	aaggaggcca	60
ctccttgcc	tccgcagccg	atcacatgaa	ggtaggtgcca	agtcctctgc	tctccgtctt	120
cctggcacag	gtgtggctgg	laccoggett	ggccccaggt	cttcagtcgc	cagagacccc	180
agccctctag	aaccagacca	gcagggtagt	gcaggctccc	agggagggaag	aggaagatga	240
gcaggaggcc	agcggggaga	aggtctgtga	ggaagagaaa	gcttggttga	tggtccagcag	300
gcagcagctt	gccaaggaga	cttcaaacct	cggtctcagc	ctgctgcgaa	agatctccat	360
gaggcagcat	ggcaacatgg	tctctctctc	atttgccatg	tccttggcca	tgacaggtct	420
gatgctgggg	gccaacaggc	cgactgaaac	ccagatcaag	agagggtctc	acttgcaggc	480
cctgaagccc	accaagcccc	ggctctctgc	tccctctctt	aagggaactc	gagagacctt	540
ctcccgcaac	ctggaaclgg	gctctctaca	ggggagtttt	gcttctctcc	acaaggattt	600
tgatgtccaa	gagactttct	tcaattttat	caagaggtat	tttgatacag	agtgcgtgoc	660
tatgaatttt	cgcaatgect	cacaggccca	aaggctctatg	aatcattaca	ttcaacagga	720
gactgggggg	aaaaatcccc	aactgtttga	tgagatlaat	cctgaaccca	aattaattct	780
tgttgattac	atcttgttca	aagggaatg	gttgacccca	tttgacctg	cttccaccca	840
agtgcacact	tcccaactgg	acaaatlaaa	gaccttlaag	gtgcccata	tgtnccgttc	900
aggcagtttt	gcttccacct	ttgacaagaa	ttttcgttgt	catgtctcca	aactgcctta	960
ccaaggaaat	gccaccatgc	tgggtgtctt	catggagaaa	atgggtgacc	acctcgcctt	1020
tnagaactac	ctgaccacag	aottggtgga	gacatggctc	agaaacatga	aaaccagaaa	1080
catggaagtt	ttctttccga	agttccagct	agatcagaag	tatgagctgc	atgagctgct	1140
tggcagatg	ggaalccagaa	gaatctcttc	accttttgc	gaccttctg	aactctcagc	1200
tactggagga	aatctccaa	tatccagggt	tttacaagaa	acagtgtattg	aagttgatga	1260
aaggggccct	gaggaagtgg	caggaaatct	gtcagaaatt	actgcttatt	ccatgcctcc	1320
tctcatccaa	gtggacccgc	catttccatt	catgatctat	gaagaaacct	ctggaatgct	1380
tctgtttctg	ggcaggytgg	tgaatccgac	tctcctataa	ttcaggacat	gcatagcaca	1440
cllctgtctg	tagtagatgc	tgaatctgag	gtatcaaaaca	cacacaggat	accagcaatg	1500
gatggcaggg	gagagtgttc	cttttgttct	taactagttt	aggggtgtct	caaatcaata	1560
cagtagtccc	caotttatctg	agggggtatc	atcaaaagac	ccccagcaga	tgcctgaaac	1620
ggtggacagt	gctgaacctt	atatataatt	tttctctcac	atccatacct	atgatesagt	1680
ctaatttata	aattagggcac	aglaagagat	taacaataat	anacaactta	agtaaatgta	1740
gttacttgaa	cgcaagcact	gcaataccat	aacagtcaaa	ctgattatag	agaaggctac	1800
caagtgaact	atgggcaggg	agcaatagaca	gtgtggaagac	atlyggcag	gggaggaatc	1860
acatccctgg	tgggacagag	caggacgatg	caagattcca	tcccactact	cagaatggca	1920
tgctgcttaa	gactttttaga	ttgttttatt	ctggaalctt	tcttttctatg	tttttggacc	1980
atgggttgac	atgggttaact	gagactgcag	aaagcaaaac	catggataag	ggagggaactac	2040
tacaaaagca	ctaaallgnt	acatattttt	taaaaaaaa			2080

&lt;210&gt; 29

&lt;211&gt; 2225

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 114551CB1

	140		145		150
Ser Gly Leu Glu Asp Gly Asp Leu Tyr	145	Asp Gly Ala Trp Cys Ala	155		
	155		160		165
Glu Glu Gln Asp Ala Asp Pro Trp Phe	170	Gln Val Asp Ala Gly His	175		180
Pro Thr Arg Phe Ser Gly Val Ile Thr	185	Gln Gly Ser Asn Ser Val	190		195
	190		205		210
Trp Arg Tyr Asp Trp Val Thr Ser Tyr	200	Lys Val Gln Phe Ser Asn	215		225
	215		220		225
Asp Ser Arg Thr Trp Trp Gly Ser Arg	230	Asn His Ser Ser Gly Met	235		240
	235		250		255
Asp Ala Val Phe Pro Ala Asn Ser Asp	245	Pro Glu Thr Pro Val Leu	265		270
	250		275		285
Asn Leu Leu Pro Glu Pro Gln Val Ala	260	Arg Phe Ile Arg Leu Leu	280		285
	265		295		300
Pro Gln Thr Trp Leu Gln Gly Gly Ala	275	Pro Cys Leu Arg Ala Glu	305		315
	280		310		315
Ile Leu Ala Cys Pro Val Ser Asp Pro	290	Asn Asp Leu Phe Leu Glu	325		330
	295		335		345
Ala Pro Ala Ser Gly Ser Ser Asp Pro	305	Leu Asp Phe Gln His His	340		350
	310		355		360
Asn Tyr Lys Ala Met Arg Lys Leu Met	320	Lys Gln Val Gln Glu Gln	370		375
	325		385		390
Cys Pro Asn Ile Thr Arg Ile Tyr Ser	335	Ile Gly Lys Ser Tyr Gln	395		405
	340		415		420
Gly Leu Lys Leu Val Met Glu Met	350	Ser Asp Lys Pro Gly Glu	425		435
	355		430		440
His Glu Leu Gly Glu Pro Glu Val Arg	365	Tyr Val Ala Gly Met His	445		455
	370		460		465
Gly Asn Glu Ala Leu Gly Arg Glu Leu	380	Leu Leu Leu Met Gln	475		480
	385		490		495
Phe Leu Cys His Glu Phe Leu Arg Gly	395	Asn Pro Arg Val Thr Arg	505		510
	400		515		525
Leu Leu Ser Glu Met Arg Ile His Leu	410	Leu Pro Ser Met Asn Pro	520		530
	415		535		545
Asp Gly Tyr Glu Ile Ala Tyr His Arg	425	Gly Ser Glu Leu Val Gly	540		555
	430		550		565
Trp Ala Glu Gly Arg Trp Asn Asn Gln	440	Ser Ile Asp Leu Asn His	565		570
	445		580		585
Asn Phe Ala Asp Leu Asn Thr Pro Leu	455	Trp Glu Ala Gln Asp Asp	595		600
	460		610		615
Gly Lys Val Pro His Ile Val Pro Asn	470	His His Leu Pro Leu Pro	625		630
	475		635		645
Thr Tyr Tyr Thr Leu Pro Asn Ala Thr	485	Val Ala Pro Glu Thr Arg	640		
	490				
Ala Val Ile Lys Trp Met Lys Arg Ile	500	Pro Phe Val Leu Ser Ala			
	505				
Asn Leu His Gly Gly Glu Leu Val Val	515	Ser Tyr Pro Phe Asp Met			
	520				
Thr Arg Thr Pro Trp Ala Ala Arg Glu	530	Leu Thr Pro Thr Pro Asp			
	535				
Asp Ala Val Phe Arg Trp Leu Ser Thr	545	Val Tyr Ala Gly Ser Asn			
	550				
Leu Ala Met Gln Asp Thr Ser Arg Arg	560	Pro Cys His Ser Gln Asp			
	565				
Phe Ser Val His Gly Asn Ile Ile Asn	575	Gly Ala Asp Trp His Thr			
	580				
Val Pro Gly Ser Met Asn Asp Phe Ser	590	Tyr Leu His Thr Asn Cys			
	595				
Phe Glu Val Thr Val Glu Leu Ser Cys	605	Asp Lys Phe Pro His Glu			
	610				
Asn Glu Leu Pro Gln Glu Trp Glu Asn	620	Asn Lys Asp Ala Leu Leu			
	625				
Thr Tyr Leu Glu Gln Val Arg Met Gly	635	Ile Ala Gly Val Val Arg			
	640				
Asp Lys Asp Thr Glu Leu Gly Ile Ala	645	Asp Ala Val Ile Ala Val			

Gly	Leu	Met	Val	Ser	Gly	Tyr	Ala	Val	Thr	Gly	Ala	Val	Leu	Gly	470	475
Gln	Asp	Arg	Leu	Ile	Asn	Tyr	Ala	Thr	Asn	Gly	Ala	Lys	Phe	Leu	480	485
Lys	Arg	His	Met	Phe	Asp	Val	Ala	Ser	Gly	Arg	Leu	Met	Arg	Thr	490	495
Cys	Tyr	Thr	Gly	Pro	Gly	Gly	Thr	Val	Glu	His	Ser	Asn	Pro	Pro	500	505
Cys	Trp	Gly	Phe	Leu	Glu	Asp	Tyr	Ala	Phe	Val	Val	Arg	Gly	Leu	510	515
Leu	Asp	Leu	Tyr	Glu	Ala	Ser	Gln	Glu	Ser	Ala	Trp	Leu	Glu	Trp	520	525
Ala	Leu	Arg	Leu	Gln	Asp	Thr	Gln	Asp	Arg	Leu	Phe	Trp	Asp	Ser	530	535
Gln	Gly	Gly	Gly	Tyr	Phe	Cys	Ser	Glu	Ala	Glu	Leu	Gly	Ala	Gly	540	545
Leu	Pro	Leu	Arg	Leu	Lys	Asp	Asp	Gln	Asp	Gly	Ala	Glu	Pro	Ser	550	555
Ala	Asn	Ser	Val	Ser	Ala	His	Asn	Leu	Leu	Arg	Leu	His	Gly	Phe	560	565
Thr	Gly	His	Lys	Asp	Trp	Met	Asp	Lys	Cys	Val	Cys	Leu	Leu	Thr	570	575
Ala	Phe	Ser	Glu	Arg	Met	Arg	Arg	Val	Pro	Val	Ala	Leu	Pro	Glu	580	585
Met	Val	Arg	Ala	Leu	Ser	Ala	Gln	Gln	Gln	Thr	Leu	Lys	Gln	Ile	590	595
Val	Ile	Cys	Gly	Asp	Arg	Gln	Ala	Lys	Asp	Thr	Lys	Ala	Leu	Val	600	605
Gln	Cys	Val	His	Ser	Val	Tyr	Ile	Pro	Asn	Lys	Val	Leu	Ile	Leu	610	615
Ala	Asp	Gly	Asp	Pro	Ser	Ser	Phe	Leu	Ser	Arg	Gln	Leu	Pro	Phe	620	625
Leu	Ser	Thr	Leu	Arg	Arg	Leu	Glu	Asp	Gln	Ala	Thr	Ala	Tyr	Val	630	635
Cys	Glu	Asn	Gln	Ala	Cys	Ser	Val	Pro	Ile	Thr	Asp	Pro	Cys	Glu	640	645
Leu	Arg	Lys	Leu	Leu	His	Pro			730						650	655

&lt;210&gt; 27

&lt;211&gt; 734

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5583922CD1

&lt;400&gt; 27

Met	Trp	Gly	Leu	Leu	Leu	Ala	Leu	Ala	Ala	Phe	Ala	Pro	Ala	Val	1	5
Gly	Pro	Ala	Leu	Gly	Ala	Pro	Arg	Asn	Ser	Val	Leu	Gly	Leu	Ala	10	15
Gln	Pro	Gly	Thr	Thr	Lys	Val	Pro	Gly	Ser	Thr	Pro	Ala	Leu	His	20	25
Ser	Ser	Pro	Ala	Gln	Pro	Pro	Ala	Glu	Thr	Ala	Asn	Gly	Thr	Ser	30	35
Glu	Gln	His	Val	Arg	Ile	Arg	Val	Ile	Lys	Lys	Lys	Lys	Val	Ile	40	45
Met	Lys	Lys	Arg	Lys	Lys	Leu	Thr	Leu	Thr	Arg	Pro	Thr	Pro	Leu	50	55
Val	Thr	Ala	Gly	Pro	Leu	Val	Thr	Pro	Thr	Pro	Ala	Gly	Thr	Leu	60	65
Asp	Pro	Ala	Glu	Lys	Gln	Glu	Thr	Gly	Cys	Pro	Pro	Leu	Gly	Leu	70	75
Glu	Ser	Leu	Arg	Val	Ser	Asp	Ser	Arg	Leu	Glu	Ala	Ser	Ser	Ser	80	85
Gln	Ser	Phe	Gly	Leu	Gly	Pro	His	Arg	Gly	Arg	Leu	Asn	Ile	Gln	90	95



&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Inocyte ID No: 5267783CD1

&lt;400&gt; 26

Met	Pro	Ala	Gly	Gly	Lys	Gly	Ser	His	Pro	Ser	Ser	Thr	Pro	Gln
1				5					10					15
Arg	Val	Pro	Asn	Arg	Leu	Ile	His	Glu	Lys	Ser	Pro	Tyr	Leu	Leu
				20					25					30
Gln	His	Ala	Tyr	Asn	Pro	Val	Asp	Trp	Tyr	Pro	Trp	Gly	Glu	Glu
				35					40					45
Ala	Phe	Asp	Lys	Ala	Arg	Lys	Glu	Asn	Lys	Pro	Ile	Phe	Leu	Ser
				50					55					60
Val	Gly	Tyr	Ser	Thr	Cys	His	Trp	Cys	His	Met	Met	Glu	Glu	Glu
				65					70					75
Ser	Phe	Gln	Asn	Glu	Glu	Ile	Gly	Arg	Leu	Leu	Ser	Glu	Asp	Phe
				80					85					90
Val	Ser	Val	Lys	Val	Asp	Arg	Glu	Glu	Arg	Pro	Asp	Val	Asp	Lys
				95					100					105
Val	Tyr	Met	Thr	Phe	Val	Gln	Ala	Thr	Ser	Ser	Gly	Gly	Gly	Trp
				110					115					120
Pro	Met	Asn	Val	Trp	Leu	Thr	Pro	Asn	Leu	Gln	Pro	Phe	Val	Gly
				125					130					135
Gly	Thr	Tyr	Phe	Pro	Pro	Glu	Asp	Gly	Leu	Thr	Arg	Val	Gly	Phe
				140					145					150
Arg	Thr	Val	Leu	Leu	Arg	Ile	Arg	Glu	Gln	Trp	Lys	Gln	Asn	Lys
				155					160					165
Asn	Thr	Leu	Leu	Glu	Asn	Ser	Gln	Arg	Val	Thr	Thr	Ala	Leu	Leu
				170					175					180
Ala	Arg	Ser	Glu	Ile	Ser	Val	Gly	Asp	Arg	Gln	Leu	Pro	Pro	Ser
				185					190					195
Ala	Ala	Thr	Val	Asn	Asn	Arg	Cys	Phe	Gln	Gln	Leu	Asp	Glu	Cly
				200					205					210
Tyr	Asp	Glu	Glu	Tyr	Gly	Gly	Phe	Ala	Glu	Ala	Pro	Lys	Phe	Pro
				215					220					225
Thr	Pro	Val	Ile	Leu	Ser	Phe	Leu	Phe	Ser	Tyr	Trp	Leu	Ser	His
				230					235					240
Arg	Leu	Thr	Gln	Asp	Gly	Ser	Arg	Ala	Gln	Gln	Met	Ala	Leu	His
				245					250					255
Thr	Leu	Lys	Met	Met	Ala	Asn	Gly	Gly	Ile	Arg	Asp	His	Val	Gly
				260					265					270
Gln	Gly	Phe	His	Arg	Tyr	Ser	Thr	Asp	Arg	Gln	Trp	His	Val	Pro
				275					280					285
His	Phe	Glu	Lys	Met	Leu	Tyr	Asp	Gln	Ala	Gln	Leu	Ala	Val	Ala
				290					295					300
Tyr	Ser	Gln	Ala	Phe	Gln	Leu	Ser	Gly	Asp	Glu	Phe	Tyr	Ser	Asp
				305					310					315
Val	Ala	Lys	Gly	Ile	Leu	Gln	Tyr	Val	Ala	Arg	Ser	Leu	Ser	His
				320					325					330
Arg	Ser	Gly	Gly	Phe	Tyr	Ser	Ala	Glu	Asp	Ala	Asp	Ser	Pro	Pro
				335					340					345
Glu	Arg	Gly	Gln	Arg	Pro	Lys	Glu	Gly	Ala	Tyr	Tyr	Val	Trp	Thr
				350					355					360
Val	Lys	Glu	Val	Gln	Gln	Leu	Leu	Pro	Glu	Pro	Val	Leu	Gly	Ala
				365					370					375
Thr	Glu	Pro	Leu	Thr	Ser	Gly	Gln	Leu	Leu	Met	Lys	His	Tyr	Gly
				380					385					390
Leu	Thr	Glu	Ala	Gly	Asn	Ile	Ser	Pro	Ser	Gln	Asp	Pro	Lys	Gly
				395					400					405
Glu	Leu	Gln	Gly	Gln	Asn	Val	Leu	Thr	Val	Arg	Tyr	Ser	Leu	Glu
				410					415					420
Leu	Thr	Ala	Ala	Arg	Phe	Gly	Leu	Asp	Val	Glu	Ala	Val	Arg	Thr
				425					430					435
Leu	Leu	Asn	Ser	Gly	Leu	Gln	Lys	Leu	Phe	Gln	Ala	Arg	Lys	His
				440					445					450
Arg	Pro	Lys	Pro	His	Leu	Asp	Ser	Lys	Met	Leu	Ala	Ala	Trp	Asn
				455					460					465

Phe	Ser	Ala	Asn	Phe	Thr	Asp	Ser	Val	Thr	Thr	Gly	Arg	Gln	Ile	
					170				175					180	
Asn	Asp	Tyr	Leu	Arg	Arg	Gln	Thr	Tyr	Gly	Gln	Val	Val	Asp	Cys	
					185				190					195	
Leu	Pro	Glu	Phe	Ser	Gln	Asp	Thr	Phe	Met	Val	Leu	Ala	Asn	Tyr	
					200				205					210	
Ile	Phe	Phe	Lys	Ala	Lys	Trp	Lys	His	Pro	Phe	Ser	Arg	Tyr	Gln	
					215				220					225	
Thr	Gln	Lys	Gln	Ala	Ser	Phe	Phe	Val	Asp	Glu	Arg	Thr	Ser	Leu	
					230				235					240	
Gln	Val	Pro	Met	Met	His	Gln	Lys	Glu	Met	His	Arg	Phe	Leu	Tyr	
					245				250					255	
Asp	Gln	Asp	Leu	Ala	Cys	Thr	Val	Leu	Gln	Ile	Glu	Tyr	Arg	Gly	
					260				265					270	
Asn	Ala	Leu	Ala	Leu	Leu	Val	Leu	Pro	Asp	Pro	Gly	Lys	Met	Lys	
					275				280					285	
Gln	Val	Glu	Ala	Ala	Leu	Gln	Pro	Gln	Thr	Leu	Arg	Lys	Trp	Gly	
					290				295					300	
Gln	Leu	Leu	Leu	Pro	Ser	Leu	Leu	Asp	Leu	His	Leu	Pro	Arg	Phe	
					305				310					315	
Ser	Ile	Ser	Gly	Thr	Tyr	Asn	Leu	Glu	Asp	Ile	Leu	Pro	Gln	Ile	
					320				325					330	
Gly	Leu	Thr	Asn	Ile	Leu	Asn	Leu	Glu	Ala	Asp	Phe	Ser	Gly	Val	
					335				340					345	
Thr	Gly	Gln	Leu	Asn	Lys	Thr	Ile	Ser	Lys	Val	Ser	His	Lys	Ala	
					350				355					360	
Met	Val	Asp	Met	Ser	Glu	Lys	Gly	Thr	Glu	Ala	Gly	Ala	Ala	Ser	
					365				370					375	
Gly	Leu	Leu	Ser	Gln	Pro	Pro	Ser	Leu	Asn	Thr	Met	Ser	Asp	Pro	
					380				385					390	
His	Ala	His	Phe	Asn	Arg	Pro	Phe	Leu	Leu	Leu	Leu	Trp	Glu	Val	
					395				400					405	
Thr	Thr	Gln	Ser	Leu	Leu	Phe	Leu	Gly	Lys	Val	Val	Asn	Pro	Val	
					410				415					420	

Ala Gly

&lt;210&gt; 25

&lt;211&gt; 114

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5208004CD1

&lt;400&gt; 25

Met	Arg	Trp	Arg	Gln	Arg	Ser	Phe	Leu	Leu	Arg	Leu	Phe	Leu	Gly	
1				5					10					15	
Ser	Leu	Arg	Gly	Gly	Gln	His	His	Pro	Pro	Leu	Thr	Leu	Pro	Ser	
				20					25					30	
Ala	Ser	Ser	Leu	Pro	Phe	Ser	Thr	Leu	Ser	Leu	Leu	Leu	Ala	Ser	
				35					40					45	
Ser	Leu	Ser	Cys	Cys	Leu	Val	Ser	Pro	Cys	Pro	Lys	Thr	Pro	Gly	
				50					55					60	
Ser	Phe	Val	Leu	Leu	Pro	Trp	Pro	Pro	Pro	Arg	Arg	Arg	Ser	Gln	
				65					70					75	
Ala	Pro	Ser	Pro	Pro	Arg	Gly	Ile	His	Thr	Thr	Gly	Ser	Cys	Trp	
				80					85					90	
Gly	Trp	Gly	Ser	Pro	Ala	Gly	Phe	Leu	Met	Pro	Cys	Ala	Gln	Gly	
				95					100					105	
Ser	Ala	Ala	Val	Ile	Phe	Gly	Leu	Ser							
				110											

&lt;210&gt; 26

&lt;211&gt; 742

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

Lys Glu Asn Lys Asp Asp Leu Asn His Val Asp Leu Asn Val Cys  
 95 100 105  
 Thr Ser Phe Ser Gly Pro Gly Arg Ser Gly Met Ala Leu Met Glu  
 110 115 120  
 Val Asn Leu Leu Ser Gly Phe Met Val Pro Ser Glu Ala Ile Ser  
 125 130 135  
 Leu Ser Glu Thr Val Lys Lys Val Glu Tyr Asp His Gly Lys Leu  
 140 145 150  
 Asn Leu Tyr Leu Asp Ser Val Asn Glu Thr Gln Phe Cys Val Asn  
 155 160 165  
 Ile Pro Ala Val Arg Asn Phe Lys Val Ser Asn Thr Gln Asp Ala  
 170 175 180  
 Ser Val Ser Ile Val Asp Tyr Tyr Glu Pro Arg Arg Gln Ala Val  
 185 190 195  
 Arg Ser Tyr Asn Ser Glu Val Lys Leu Ser Ser Cys Asp Leu Cys  
 200 205 210  
 Ser Asp Val Gln Gly Cys Arg Pro Cys Glu Asp Gly Ala Ser Gly  
 215 220 225  
 Ser His His His Ser Ser Val Ile Phe Ile Phe Cys Phe Lys Leu  
 230 235 240  
 Leu Tyr Phe Met Glu Leu Trp Leu  
 245

<210> 23  
 <211> 520  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3883861CD1

<400> 23  
 Met Val Ala Arg Val Gly Leu Leu Leu Arg Ala Leu Gln Leu Leu  
 1 5 10 15  
 Leu Trp Gly His Leu Asp Ala Gln Pro Ala Glu Arg Gly Gly Gln  
 20 25 30  
 Glu Leu Arg Lys Glu Ala Glu Ala Phe Leu Glu Lys Tyr Gly Tyr  
 35 40 45  
 Leu Asn Glu Gln Val Pro Lys Ala Pro Thr Ser Thr Arg Phe Ser  
 50 55 60  
 Asp Ala Ile Arg Ala Phe Gln Trp Val Ser Gln Leu Pro Val Ser  
 65 70 75  
 Gly Val Leu Asp Arg Ala Thr Leu Arg Gln Met Thr Arg Pro Arg  
 80 85 90  
 Cys Gly Val Thr Asp Thr Asn Ser Tyr Ala Ala Trp Ala Glu Arg  
 95 100 105  
 Ile Ser Asp Leu Phe Ala Arg His Arg Thr Lys Met Arg Arg Lys  
 110 115 120  
 Lys Arg Phe Ala Lys Gln Gly Asn Lys Trp Tyr Lys Gln His Leu  
 125 130 135  
 Ser Tyr Arg Leu Val Asn Trp Pro Glu His Leu Pro Glu Pro Ala  
 140 145 150  
 Val Arg Gly Ala Val Arg Ala Ala Phe Gln Leu Trp Ser Asn Val  
 155 160 165  
 Ser Ala Leu Glu Phe Trp Glu Ala Pro Ala Thr Gly Pro Ala Asp  
 170 175 180  
 Ile Arg Leu Thr Phe Phe Gln Gly Asp His Asn Asp Gly Leu Gly  
 185 190 195  
 Asn Ala Phe Asp Gly Pro Gly Gly Ala Leu Ala His Ala Phe Leu  
 200 205 210  
 Pro Arg Arg Gly Glu Ala His Phe Asp Gln Asp Glu Arg Trp Ser  
 215 220 225  
 Leu Ser Arg Arg Arg Gly Arg Asn Leu Phe Val Val Leu Ala His  
 230 235 240  
 Glu Ile Gly His Thr Leu Gly Leu Thr His Ser Pro Ala Pro Arg  
 245 250 255  
 Ala Leu Met Ala Pro Tyr Tyr Lys Arg Leu Gly Arg Asp Ala Leu

Leu Ser Trp Asp	260	Val Leu Ala Val	265	Gln Ser Leu Tyr Gly	270
Asp	275	Val Ala Val Gln	280	Lys	285
Pro Leu Gly Gly	290	Leu	295	Pro Gly Lys Leu	300
Thr Asp Phe Glu	305	Trp Asp Ser Tyr	310	Pro Gln Gly Arg	315
Pro Glu Thr Gln	320	Pro Lys Tyr Cys	325	His Ser Ser Phe Asp	330
Ile Thr Val Asp	335	Gln Gln Gln Leu	340	Tyr Ile Phe Lys Gly	345
His Phe Trp Glu	350	Val Ala Ala Asp Gly	355	Asn Val Ser Glu Pro	360
Pro Leu Gln Glu	365	Trp Val Gly Leu	370	Pro Asn Ile Glu	375
Ala Ala Val Ser	380	Leu Asn Asp Gly Asp	385	Phe Tyr Phe Phe Lys	390
Gly Arg Cys Trp	395	Arg Phe Arg Gly Pro	400	Lys Pro Val Trp Gly	405
Pro Gln Leu Cys	410	Arg Ala Gly Gly Leu	415	Pro Arg His Pro Asp	420
Ala Leu Phe Phe	425	Pro Leu Arg Arg	430	Leu Ile Leu Phe Lys	435
Ala Arg Tyr Tyr	440	Val Leu Ala Arg Gly	445	Gly Leu Gln Val Glu	450
Tyr Tyr Pro Arg	455	Ser Leu Gln Asp Trp	460	Gly Ile Pro Glu	465
Val Ser Gly Ala	470	Leu Pro Arg Pro Asp	475	Gly Ser Ile Ile Phe	480
Arg Asp Asp Arg	485	Tyr Trp Arg Leu Asp	490	Gln Ala Lys Leu Gln	495
Thr Thr Ser Gly	500	Arg Trp Ala Thr Glu	505	Leu Pro Trp Met Gly	510
Trp His Ala Asn	515	Ser Gly Ser Ala Leu	520	Phe	

&lt;210&gt; 24

&lt;211&gt; 422

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4993873CD1

&lt;400&gt; 24

Met Gly Pro Ala Trp	1	Leu Trp Leu Leu Gly	10	Thr Gly Ile Leu Ala	15
Ser Val His Cys Gln	20	Pro Leu Leu Ala His	25	Gly Asp Lys Ser Leu	30
Gln Gly Pro Gln Pro	35	Pro Arg His Gln Leu	40	Ser Glu Pro Ala Pro	45
Ala Tyr His Arg Ile	50	Thr Pro Thr Ile Thr	55	Asn Phe Ala Leu Arg	60
Leu Tyr Lys Glu Leu	65	Ala Ala Asp Ala Pro	70	Gly Asn Ile Phe Phe	75
Ser Pro Val Ser Ile	80	Ser Thr Thr Leu Ala	85	Leu Leu Ser Leu Gly	90
Ala Gln Ala Asn Thr	95	Ser Ala Leu Ile Leu	100	Glu Gly Leu Gly Phe	105
Asn Leu Thr Glu Thr	110	Pro Glu Ala Asp Ile	115	His Gln Gly Phe Arg	120
Ser Leu Leu His Thr	125	Leu Ala Leu Pro Ser	130	Pro Lys Leu Glu Leu	135
Lys Val Gly Asn Ser	140	Leu Phe Leu Asp Lys	145	Arg Leu Lys Pro Arg	150
Gln His Tyr Leu Asp	155	Ser Ile Lys Glu Leu	160	Tyr Gly Ala Phe Ala	165

Ser	Lys	Cys	Phe	Cys	Glu	Asn	Cys	Gly	Lys	Lys	Thr	Arg	Gly	Lys	225
				230					235						240
Gln	Val	Leu	Lys	Leu	Thr	His	Leu	Pro	Gln	Thr	Leu	Thr	Ile	His	255
				245					250						255
Leu	Met	Arg	Phe	Ser	Ile	Arg	Asn	Ser	Gln	Thr	Arg	Lys	Ile	Cys	270
				260					265						270
His	Ser	Leu	Tyr	Phe	Pro	Gln	Ser	Leu	Asp	Phe	Ser	Gln	Ile	Leu	285
				275					280						285
Pro	Met	Lys	Arg	Glu	Ser	Cys	Asp	Ala	Glu	Glu	Gln	Ser	Gly	Gly	300
				290					295						300
Gln	Tyr	Glu	Leu	Phe	Ala	Val	Ile	Ala	His	Val	Gly	Met	Ala	Asp	315
				305					310						315
Ser	Gly	His	Tyr	Cys	Val	Tyr	Ile	Arg	Asn	Ala	Val	Asp	Gly	Lys	330
				320					325						330
Trp	Phe	Cys	Phe	Asn	Asp	Ser	Asn	Ile	Cys	Leu	Val	Ser	Trp	Glu	345
				335					340						345
Asp	Ile	Gln	Cys	Thr	Tyr	Gly	Asn	Pro	Asn	Tyr	His	Trp	Gln	Glu	360
				350					355						360
Thr	Ala	Tyr	Leu	Leu	Val	Tyr	Met	Lys	Met	Glu	Cys				
				365					370						

&lt;210&gt; 21

&lt;211&gt; 94

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3092341CD1

&lt;400&gt; 21

Met	Leu	Arg	Gly	Val	Leu	Gly	Lys	Thr	Phe	Arg	Leu	Val	Gly	Tyr	
1				5					10					15	
Thr	Ile	Gln	Tyr	Gly	Cys	Ile	Ala	His	Cys	Ala	Phe	Glu	Tyr	Val	
				20					25					30	
Gly	Gly	Val	Val	Met	Val	Pro	Met	Gly	His	Val	Trp	Leu	Glu	Gly	
				35					40					45	
Asp	Asn	Leu	Gln	Asn	Ser	Thr	Asp	Ser	Arg	Cys	Tyr	Gly	Pro	Ile	
				50					55					60	
Pro	Tyr	Gly	Leu	Ile	Arg	Gly	Arg	Ile	Phe	Phe	Lys	Ile	Trp	Leu	
				65					70					75	
Leu	Ser	Asp	Phe	Gly	Phe	Leu	Arg	Ala	Ser	Pro	Asn	Gly	His	Arg	
				80					85					90	
Phe	Ser	Asp	Asp												

&lt;210&gt; 22

&lt;211&gt; 248

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3658034CD1

&lt;400&gt; 22

Met	Asn	Thr	Glu	Arg	Thr	Asn	Ile	Gln	Val	Thr	Val	Thr	Gly	Pro	
1				5					10					15	
Ser	Ser	Pro	Ser	Pro	Val	Lys	Phe	Leu	Ile	Asp	Thr	His	Asn	Arg	
				20					25					30	
Leu	Leu	Leu	Gln	Thr	Ala	Glu	Leu	Ala	Val	Val	Gln	Pro	Thr	Ala	
				35					40					45	
Val	Asn	Ile	Ser	Ala	Asn	Gly	Phe	Gly	Phe	Ala	Ile	Cys	Gln	Leu	
				50					55					60	
Asn	Val	Val	Tyr	Asn	Val	Lys	Ala	Ser	Gly	Ser	Ser	Arg	Arg	Arg	
				65					70					75	
Arg	Ser	Ile	Gln	Asn	Gln	Glu	Ala	Phe	Asp	Leu	Asp	Val	Ala	Val	
				80					85					90	

His	Val	Gly	Leu	Met	Leu	Lys	Glu	Asn	His	Cys	Leu	Val	Ala	Leu
			200						205					210
His	Met	Cys	Lys	His	Asp	Ile	Lys	Asn	Ser	Gly	Ile	Gln	Gln	Leu
			215						220					225
Cys	Asp	Ala	Leu	Tyr	Leu	Asn	Ser	Ser	Leu	Arg	Tyr	Leu	Asp	Val
			230						235					240
Ser	Cys	Asn	Lys	Ile	Thr	His	Asp	Gly	Met	Val	Tyr	Leu	Ala	Asp
			245						250					255
Val	Leu	Lys	Ser	Asn	Thr	Thr	Leu	Glu	Val	Ile	Asp	Leu	Ser	Phe
			260						265					270
Asn	Arg	Ile	Glu	Asn	Ala	Gly	Ala	Asn	Tyr	Leu	Ser	Glu	Thr	Leu
			275						280					285
Thr	Ser	His	Asn	Arg	Ser	Leu	Lys	Ala	Leu	Ser	Val	Val	Ser	Asn
			290						295					300
Asn	Ile	Glu	Gly	Glu	Gly	Leu	Val	Ala	Leu	Ser	Gln	Ser	Met	Lys
			305						310					315
Thr	Asn	Leu	Thr	Phe	Ser	His	Ile	Tyr	Ile	Trp	Gly	Asn	Lys	Phe
			320						325					330
Asp	Glu	Ala	Thr	Cys	Ile	Ala	Tyr	Ser	Asp	Leu	Ile	Gln	Met	Gly
			335						340					345
Cys	Leu	Lys	Pro	Asp	Asn	Thr	Asp	Val	Glu	Pro	Phe	Val	Val	Asp
			350						355					360
Gly	Arg	Val	Tyr	Leu	Ala	Glu	Val	Ser	Asn	Gly	Leu	Lys	Lys	His
			365						370					375
Tyr	Tyr	Trp	Thr	Ser	Thr	Tyr	Gly	Glu	Ser	Tyr	Asp	His	Ser	Ser
			380						385					390
Asn	Ala	Gly	Phe	Ala	Leu	Val	Pro	Val	Gly	Gln	Gln	Pro		
			395						400					

&lt;210&gt; 20

&lt;211&gt; 372

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2754960CD1

&lt;400&gt; 20

Met	Ser	Lys	Ala	Phe	Gly	Leu	Leu	Arg	Gln	Ile	Cys	Gln	Ser	Ile
1				5					10					15
Leu	Ala	Glu	Ser	Ser	Gln	Ser	Pro	Ala	Asp	Leu	Glu	Glu	Lys	Lys
				20					25					30
Glu	Glu	Asp	Ser	Asn	Met	Lys	Arg	Glu	Gln	Pro	Arg	Glu	Arg	Pro
				35					40					45
Arg	Ala	Trp	Asp	Tyr	Pro	His	Gly	Leu	Val	Gly	Leu	His	Asn	Ile
				50					55					60
Gly	Gln	Thr	Cys	Cys	Leu	Asn	Ser	Leu	Ile	Gln	Val	Phe	Val	Met
				65					70					75
Asn	Val	Asp	Phe	Thr	Arg	Ile	Leu	Lys	Arg	Ile	Thr	Val	Pro	Arg
				80					85					90
Gly	Ala	Asp	Glu	Gln	Arg	Arg	Ser	Val	Pro	Phe	Gln	Met	Leu	Leu
				95					100					105
Leu	Leu	Glu	Lys	Met	Gln	Asp	Ser	Arg	Gln	Lys	Ala	Val	Arg	Pro
				110					115					120
Leu	Glu	Leu	Ala	Tyr	Cys	Leu	Gln	Lys	Cys	Asn	Val	Pro	Leu	Phe
				125					130					135
Val	Gln	His	Asp	Ala	Ala	Gln	Leu	Tyr	Leu	Lys	Leu	Trp	Asn	Leu
				140					145					150
Ile	Lys	Asp	Gln	Ile	Thr	Asp	Val	His	Leu	Val	Glu	Arg	Leu	Gln
				155					160					165
Ala	Leu	Tyr	Thr	Ile	Arg	Val	Lys	Asp	Ser	Leu	Ile	Cys	Val	Asp
				170					175					180
Cys	Ala	Met	Glu	Ser	Ser	Arg	Asn	Ser	Ser	Met	Leu	Thr	Leu	Pro
				185					190					195
Leu	Ser	Leu	Phe	Asp	Val	Asp	Ser	Lys	Pro	Leu	Lys	Thr	Leu	Glu
				200					205					210
Asp	Ala	Leu	His	Cys	Phe	Phe	Gln	Pro	Arg	Glu	Leu	Ser	Ser	Lys

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 411344CD1

&lt;400&gt; 14

Met	Ala	Ser	Met	Gln	Lys	Arg	Leu	Gln	Lys	Glu	Leu	Leu	Ala	Leu	
1				5					10					15	
Gln	Asn	Asp	Pro	Pro	Pro	Gly	Met	Thr	Leu	Asn	Glu	Lys	Ser	Val	
				20					25					30	
Gln	Asn	Ser	Ile	Thr	Gln	Trp	Ile	Val	Asp	Met	Glu	Gly	Ala	Pro	
				35					40					45	
Gly	Thr	Leu	Tyr	Gln	Gly	Glu	Lys	Phe	Gln	Leu	Leu	Phe	Lys	Phe	
				50					55					60	
Ser	Ser	Arg	Tyr	Pro	Phe	Asp	Ser	Pro	Gln	Val	Met	Phe	Thr	Gly	
				65					70					75	
Glu	Asn	Ile	Pro	Val	His	Pro	His	Val	Tyr	Ser	Asn	Gly	His	Ile	
				80					85					90	
Cys	Leu	Ser	Ile	Leu	Thr	Glu	Asp	Trp	Ser	Pro	Ala	Leu	Ser	Val	
				95					100					105	
Gln	Ser	Val	Cys	Leu	Ser	Ile	Ile	Ser	Met	Leu	Ser	Ser	Cys	Lys	
				110					115					120	
Glu	Lys	Arg	Arg	Pro	Pro	Asp	Asn	Ser	Phe	Tyr	Val	Arg	Thr	Cys	
				125					130					135	
Asn	Lys	Asn	Pro	Lys	Lys	Thr	Lys	Trp	Trp	Tyr	His	Asp	Asp	Thr	
				140					145					150	

Cys

&lt;210&gt; 15

&lt;211&gt; 362

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1256390CD1

&lt;400&gt; 15

Met	Leu	Val	Pro	Gly	Gly	Leu	Gly	Tyr	Asp	Arg	Ser	Leu	Ala	Gln	
1				5					10					15	
His	Arg	Gln	Glu	Ile	Val	Asp	Lys	Ser	Val	Ser	Pro	Trp	Ser	Leu	
				20					25					30	
Glu	Thr	Tyr	Ser	Tyr	Asn	Ile	Tyr	His	Pro	Met	Gly	Glu	Ile	Tyr	
				35					40					45	
Glu	Trp	Met	Arg	Glu	Ile	Ser	Glu	Lys	Tyr	Lys	Glu	Val	Val	Thr	
				50					55					60	
Gln	His	Phe	Leu	Gly	Val	Thr	Tyr	Glu	Thr	His	Pro	Met	Tyr	Tyr	
				65					70					75	
Leu	Lys	Ile	Ser	Gln	Pro	Ser	Gly	Asn	Pro	Lys	Lys	Ile	Ile	Trp	
				80					85					90	
Met	Asp	Cys	Gly	Ile	His	Ala	Arg	Glu	Trp	Ile	Ala	Pro	Ala	Phe	
				95					100					105	
Cys	Gln	Trp	Phe	Val	Lys	Glu	Ile	Leu	Gln	Asn	His	Lys	Asp	Asn	
				110					115					120	
Ser	Ser	Ile	Arg	Lys	Leu	Leu	Arg	Asn	Leu	Asp	Phe	Tyr	Val	Leu	
				125					130					135	
Pro	Val	Leu	Asn	Ile	Asp	Gly	Tyr	Ile	Tyr	Thr	Trp	Thr	Thr	Asp	
				140					145					150	
Arg	Leu	Trp	Arg	Lys	Ser	Arg	Ser	Pro	His	Asn	Asn	Gly	Thr	Cys	
				155					160					165	
Phe	Gly	Thr	Asp	Leu	Asn	Arg	Asn	Phe	Asn	Ala	Ser	Trp	Cys	Ser	
				170					175					180	
Ile	Gly	Ala	Ser	Arg	Asn	Cys	Gln	Asp	Gln	Thr	Phe	Cys	Gly	Thr	
				185					190					195	
Gly	Pro	Val	Ser	Glu	Pro	Glu	Thr	Lys	Ala	Val	Ala	Ser	Phe	Ile	
				200					205					210	
Glu	Ser	Lys	Lys	Asp	Asp	Ile	Leu	Cys	Phe	Leu	Thr	Met	His	Ser	
				215					220					225	

Leu Leu Phe Tyr Glu Arg Val Leu Ser Arg Met Gln His Gln Ser  
 500 505 510  
 Gln Glu Cys Lys Ser Glu Glu  
 515

&lt;210&gt; 13

&lt;211&gt; 346

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5853753CD1

&lt;400&gt; 13

Met Val Glu Lys Glu Glu Ala Gly Gly Gly Ile Ser Glu Glu Glu  
 1 5 10 15  
 Ala Ala Gln Tyr Asp Arg Gln Ile Arg Leu Trp Gly Leu Glu Ala  
 20 25 30  
 Gln Lys Arg Leu Arg Ala Ser Arg Val Leu Leu Val Gly Leu Lys  
 35 40 45  
 Gly Leu Gly Ala Glu Ile Ala Lys Asn Leu Ile Leu Ala Gly Val  
 50 55 60  
 Lys Gly Leu Thr Met Leu Asp His Glu Gln Val Thr Pro Glu Asp  
 65 70 75  
 Pro Gly Ala Gln Phe Leu Ile Arg Thr Gly Ser Val Gly Arg Asn  
 80 85 90  
 Arg Ala Glu Ala Ser Leu Glu Arg Ala Gln Asn Leu Asn Pro Met  
 95 100 105  
 Val Asp Val Lys Val Asp Thr Glu Asp Ile Glu Lys Lys Pro Glu  
 110 115 120  
 Ser Phe Phe Thr Gln Phe Asp Ala Val Cys Leu Thr Cys Cys Ser  
 125 130 135  
 Arg Asp Val Ile Val Lys Val Asp Gln Ile Cys His Lys Asn Ser  
 140 145 150  
 Ile Lys Phe Phe Thr Gly Asp Val Phe Gly Tyr His Gly Tyr Thr  
 155 160 165  
 Phe Ala Asn Leu Gly Glu His Gln Phe Val Glu Glu Lys Thr Lys  
 170 175 180  
 Val Ala Lys Val Ser Gln Gly Val Glu Asp Gly Pro Asp Thr Lys  
 185 190 195  
 Arg Ala Lys Leu Asp Ser Ser Glu Thr Thr Met Val Lys Lys Lys  
 200 205 210  
 Val Val Phe Cys Pro Val Lys Glu Ala Leu Glu Val Asp Trp Ser  
 215 220 225  
 Ser Glu Lys Ala Lys Ala Ala Leu Lys Arg Thr Thr Ser Asp Tyr  
 230 235 240  
 Phe Leu Leu Gln Val Leu Leu Lys Phe Arg Thr Asp Lys Gly Arg  
 245 250 255  
 Asp Pro Ser Ser Asp Thr Tyr Glu Glu Asp Ser Glu Leu Leu Leu  
 260 265 270  
 Gln Ile Arg Asn Asp Val Leu Asp Ser Leu Gly Ile Ser Pro Asp  
 275 280 285  
 Leu Leu Pro Glu Asp Phe Val Arg Tyr Cys Phe Ser Glu Met Ala  
 290 295 300  
 Pro Val Cys Ala Val Val Gly Gly Ile Leu Ala Gln Glu Ile Val  
 305 310 315  
 Lys Ala Leu Ser Gln Arg Asp Pro Pro His Asn Asn Phe Phe Phe  
 320 325 330  
 Phe Asp Gly Met Lys Gly Asn Gly Ile Val Glu Cys Leu Gly Pro  
 335 340 345

Lys

&lt;210&gt; 14

&lt;211&gt; 151

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens



Tyr	Gly	Gln	Leu	Ile	Leu	Thr	Pro	Tyr	Gly	Tyr	Thr	Lys	Asn	Lys	
				230					235					240	
Ser	Ser	Asn	His	Pro	Glu	Met	Ile	Gln	Val	Gly	Gln	Lys	Ala	Ala	
				245					250					255	
Asn	Ala	Leu	Lys	Ala	Lys	Tyr	Gly	Thr	Asn	Tyr	Arg	Val	Gly	Ser	
				260					265					270	
Ser	Ala	Asp	Ile	Leu	Tyr	Ala	Ser	Ser	Gly	Ser	Ser	Arg	Asp	Trp	
				275					280					285	
Ala	Arg	Asp	Ile	Gly	Ile	Pro	Phe	Ser	Tyr	Thr	Phe	Glu	Leu	Arg	
				290					295					300	
Asp	Ser	Gly	Thr	Tyr	Gly	Phe	Val	Leu	Pro	Glu	Ala	Gln	Ile	Gln	
				305					310					315	
Pro	Thr	Cys	Glu	Glu	Thr	Met	Glu	Ala	Val	Leu	Ser	Val	Leu	Asp	
				320					325					330	
Asp	Val	Tyr	Ala	Lys	His	Trp	His	Ser	Asp	Ser	Ala	Gly	Arg	Val	
				335					340					345	
Thr	Ser	Ala	Thr	Met	Leu	Leu	Gly	Leu	Leu	Val	Ser	Cys	Met	Ser	
				350					355					360	

Ileu Leu

<210> 16  
 <211> 123  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1786774CD1

Met	Ser	Gly	Glu	Glu	Leu	Ser	Glu	Ser	Thr	Pro	Glu	Pro	Gln	Lys	
1				5					10					15	
Glu	Ile	Ser	Glu	Ser	Leu	Ser	Val	Thr	Arg	Asp	Gln	Asp	Glu	Asp	
				20					25					30	
Asp	Lys	Ala	Pro	Glu	Pro	Thr	Trp	Ala	Asp	Asp	Leu	Pro	Ala	Thr	
				35					40					45	
Thr	Ser	Ser	Glu	Ala	Thr	Thr	Thr	Pro	Arg	Pro	Leu	Leu	Ser	Thr	
				50					55					60	
Pro	Val	Asp	Gly	Ala	Glu	Asp	Pro	Arg	Cys	Leu	Glu	Ala	Leu	Lys	
				65					70					75	
Pro	Gly	Asn	Cys	Gly	Glu	Tyr	Val	Val	Arg	Trp	Tyr	Tyr	Asp	Lys	
				80					85					90	
Gln	Val	Asn	Ser	Cys	Ala	Arg	Phe	Trp	Phe	Ser	Gly	Cys	Asn	Gly	
				95					100					105	
Ser	Gly	Asn	Arg	Phe	Asn	Ser	Glu	Lys	Glu	Cys	Gln	Glu	Thr	Cys	
				110					115					120	

Ile Gln Gly

<210> 17  
 <211> 983  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1911808CD1

Met	Ala	Pro	Arg	Leu	Gln	Leu	Glu	Lys	Ala	Ala	Trp	Arg	Trp	Ala	
1				5					10					15	
Glu	Thr	Val	Arg	Pro	Glu	Glu	Val	Ser	Gln	Glu	His	Ile	Glu	Thr	
				20					25					30	
Ala	Tyr	Arg	Ile	Trp	Leu	Glu	Pro	Cys	Ile	Arg	Gly	Val	Cys	Arg	
				35					40					45	
Arg	Asn	Cys	Lys	Gly	Asn	Pro	Asn	Cys	Leu	Val	Gly	Ile	Gly	Glu	
				50					55					60	
His	Ile	Trp	Leu	Gly	Glu	Ile	Asp	Glu	Asn	Ser	Phe	His	Asn	Ile	

Tyr	Gly	Gln	Leu	Ile	Leu	Thr	Pro	Tyr	Gly	Tyr	Thr	Lys	Asn	Lys	
				230					235					240	
Ser	Ser	Asn	His	Pro	Glu	Met	Ile	Gln	Val	Gly	Gln	Lys	Ala	Ala	
				245					250					255	
Asn	Ala	Leu	Lys	Ala	Lys	Tyr	Gly	Thr	Asn	Tyr	Arg	Val	Gly	Ser	
				260					265					270	
Ser	Ala	Asp	Ile	Leu	Tyr	Ala	Ser	Ser	Gly	Ser	Ser	Arg	Asp	Trp	
				275					280					285	
Ala	Arg	Asp	Ile	Gly	Ile	Pro	Phe	Ser	Tyr	Thr	Phe	Glu	Leu	Arg	
				290					295					300	
Asp	Ser	Gly	Thr	Tyr	Gly	Phe	Val	Leu	Pro	Glu	Ala	Gln	Ile	Gln	
				305					310					315	
Pro	Thr	Cys	Glu	Glu	Thr	Met	Glu	Ala	Val	Leu	Ser	Val	Leu	Asp	
				320					325					330	
Asp	Val	Tyr	Ala	Lys	His	Trp	His	Ser	Asp	Ser	Ala	Gly	Arg	Val	
				335					340					345	
Thr	Ser	Ala	Thr	Met	Leu	Leu	Gly	Leu	Leu	Val	Ser	Cys	Met	Ser	
				350					355					360	

Ileu Leu

<210> 16  
 <211> 123  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1786774CD1

<400> 16  
 Met Ser Gly Glu Glu Leu Ser Glu Ser Thr Pro Glu Pro Gln Lys  
 1 5 10 15  
 Glu Ile Ser Glu Ser Leu Ser Val Thr Arg Asp Gln Asp Glu Asp  
 20 25 30  
 Asp Lys Ala Pro Glu Pro Thr Trp Ala Asp Asp Leu Pro Ala Thr  
 35 40 45  
 Thr Ser Ser Glu Ala Thr Thr Thr Pro Arg Pro Leu Leu Ser Thr  
 50 55 60  
 Pro Val Asp Gly Ala Glu Asp Pro Arg Cys Leu Glu Ala Leu Lys  
 65 70 75  
 Pro Gly Asn Cys Gly Glu Tyr Val Val Arg Trp Tyr Tyr Asp Lys  
 80 85 90  
 Gln Val Asn Ser Cys Ala Arg Phe Trp Phe Ser Gly Cys Asn Gly  
 95 100 105  
 Ser Gly Asn Arg Phe Asn Ser Glu Lys Glu Cys Gln Glu Thr Cys  
 110 115 120  
 Ile Gln Gly

<210> 17  
 <211> 983  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1911808CD1

<400> 17  
 Met Ala Pro Arg Leu Gln Leu Glu Lys Ala Ala Trp Arg Trp Ala  
 1 5 10 15  
 Glu Thr Val Arg Pro Glu Glu Val Ser Gln Glu His Ile Glu Thr  
 20 25 30  
 Ala Tyr Arg Ile Trp Leu Glu Pro Cys Ile Arg Gly Val Cys Arg  
 35 40 45  
 Arg Asn Cys Lys Gly Asn Pro Asn Cys Leu Val Gly Ile Gly Glu  
 50 55 60  
 His Ile Trp Leu Gly Glu Ile Asp Glu Asn Ser Phe His Asn Ile

1	5	10	15
Pro Gly His Pro Arg	Lys Arg Ala Gly Arg	Asn Glu Pro Leu Lys	
20	25	30	
Lys Glu Arg Leu Lys	Trp Lys Ser Asp Tyr	Pro Met Thr Asp Gly	
35	40	45	
Gln Leu Arg Ser Lys	Arg Asp Glu Phe Trp	Asp Thr Ala Pro Ala	
50	55	60	
Phe Glu Gly Arg Lys	Glu Ile Trp Asp Ala	Leu Lys Ala Ala Ala	
65	70	75	
Tyr Ala Ala Glu Ala	Asn Asp His Glu Leu	Ala Gln Ala Ile Leu	
80	85	90	
Asp Gly Ala Ser Ile	Thr Leu Pro His Gly	Thr Leu Cys Glu Cys	
95	100	105	
Tyr Asp Glu Leu Gly	Asn Arg Tyr Gln Leu	Pro Ile Tyr Cys Leu	
110	115	120	
Ser Pro Pro Val Asn	Leu Leu Leu Glu His	Thr Glu Glu Glu Ser	
125	130	135	
Leu Glu Pro Pro Glu	Pro Pro Pro Ser Val	Arg Arg Glu Phe Pro	
140	145	150	
Leu Lys Val Arg Leu	Ser Thr Gly Lys Asp	Val Arg Leu Ser Ala	
155	160	165	
Ser Leu Pro Asp Thr	Val Gly Gln Leu Lys	Arg Gln Leu His Ala	
170	175	180	
Gln Glu Gly Ile Glu	Pro Ser Trp Gln Arg	Trp Phe Phe Ser Gly	
185	190	195	
Lys Leu Leu Thr Asp	Arg Thr Arg Leu Gln	Glu Thr Lys Ile Gln	
200	205	210	
Lys Asp Phe Val Ile	Gln Val Ile Ile Asn	Gln Pro Pro Pro Pro	
215	220	225	

Gln Asp

&lt;210&gt; 19

&lt;211&gt; 403

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2323917CD1

&lt;400&gt; 19

Met Glu Lys Ser Gln	Lys Ile Asn Pro Phe	Ile Leu His Ile Leu	
1	5	10	15
Gln Glu Val Asp Glu	Glu Ile Lys Lys Gly	Leu Ala Ala Gly Ile	
20	25	30	
Thr Leu Asn Ile Ala	Gly Asn Asn Arg Leu	Val Pro Val Glu Arg	
35	40	45	
Val Thr Gly Glu Asp	Phe Trp Ile Leu Ser	Lys Ile Leu Lys Asn	
50	55	60	
Cys Leu Tyr Ile Asn	Gly Leu Asp Val Gly	Tyr Asn Leu Leu Cys	
65	70	75	
Asp Val Gly Ala Tyr	Tyr Ala Ala Lys Leu	Leu Gln Lys Gln Leu	
80	85	90	
Asn Leu Ile Tyr Leu	Asn Leu Met Phe Asn	Asp Ile Gly Pro Glu	
95	100	105	
Gly Gly Glu Leu Ile	Ala Lys Val Leu His	Lys Asn Arg Thr Leu	
110	115	120	
Lys Tyr Leu Arg Met	Thr Gly Asn Lys Ile	Glu Asn Lys Gly Gly	
125	130	135	
Met Phe Phe Ala Ala	Met Leu Gln Ile Asn	Ser Ser Leu Glu Lys	
140	145	150	
Leu Asp Leu Gly Asp	Cys Asp Leu Gly Met	Gln Ser Val Ile Ala	
155	160	165	
Phe Ala Thr Val Leu	Thr Gln Asn Gln Ala	Ile Lys Ala Ile Asn	
170	175	180	
Leu Asn Arg Pro Ile	Leu Tyr Gly Glu Gln	Glu Glu Ser Thr Val	
185	190	195	

Leu	Asn	Glu	Asp	Tyr	Lys	Thr	Val	Asn	Asn	Leu	Leu	Lys	Ala	Ala
				575					580					585
Val	Lys	Gly	Ser	Asp	Gly	Phe	Trp	Val	Gly	Lys	Ser	Ser	Leu	Arg
				590					595					600
Ser	Trp	Arg	Gln	Leu	Ala	Leu	Glu	Gln	Leu	Asp	Glu	Gln	Asp	Gly
				605					610					615
Asp	Ala	Glu	Gln	Ser	Asn	Gly	Lys	Met	Asn	Gly	Ser	Thr	Leu	Asn
				620					625					630
Lys	Asp	Glu	Ser	Lys	Glu	Glu	Arg	Lys	Glu	Glu	Glu	Glu	Leu	Asn
				635					640					645
Phe	Asn	Glu	Asp	Ile	Leu	Cys	Pro	His	Gly	Glu	Leu	Cys	Ile	Ser
				650					655					660
Glu	Asn	Glu	Arg	Arg	Leu	Val	Ser	Lys	Glu	Ala	Trp	Ser	Lys	Leu
				665					670					675
Gln	Gln	Tyr	Phe	Pro	Lys	Ala	Pro	Glu	Phe	Pro	Ser	Tyr	Lys	Glu
				680					685					690
Cys	Cys	Ser	Gln	Cys	Lys	Ile	Leu	Glu	Arg	Glu	Gly	Glu	Glu	Asn
				695					700					705
Glu	Ala	Leu	His	Lys	Met	Ile	Ala	Asn	Gln	Gln	Lys	Thr	Ser	Leu
				710					715					720
Pro	Asn	Leu	Phe	Gln	Asp	Lys	Asn	Arg	Pro	Cys	Leu	Ser	Asn	Trp
				725					730					735
Pro	Glu	Asp	Thr	Asp	Val	Leu	Tyr	Ile	Val	Ser	Gln	Phe	Phe	Val
				740					745					750
Glu	Glu	Trp	Arg	Lys	Phe	Val	Arg	Lys	Pro	Thr	Arg	Cys	Ser	Pro
				755					760					765
Val	Ser	Ser	Val	Gly	Asn	Ser	Ala	Leu	Leu	Cys	Pro	His	Gly	Gly
				770					775					780
Leu	Met	Phe	Thr	Phe	Ala	Ser	Met	Thr	Lys	Glu	Asp	Ser	Lys	Leu
				785					790					795
Ile	Ala	Leu	Ile	Trp	Pro	Ser	Glu	Trp	Gln	Met	Ile	Gln	Lys	Leu
				800					805					810
Phe	Val	Val	Asp	His	Val	Ile	Lys	Ile	Thr	Arg	Ile	Glu	Val	Gly
				815					820					825
Asp	Val	Asn	Pro	Ser	Glu	Thr	Gln	Tyr	Ile	Ser	Glu	Pro	Lys	Leu
				830					835					840
Cys	Pro	Glu	Cys	Arg	Glu	Gly	Leu	Leu	Cys	Gln	Gln	Gln	Arg	Asp
				845					850					855
Leu	Arg	Glu	Tyr	Thr	Gln	Ala	Thr	Ile	Tyr	Val	His	Lys	Val	Val
				860					865					870
Asp	Asn	Lys	Lys	Val	Met	Lys	Asp	Ser	Ala	Pro	Glu	Leu	Asn	Val
				875					880					885
Ser	Ser	Ser	Glu	Thr	Glu	Glu	Asp	Lys	Glu	Glu	Ala	Lys	Pro	Asp
				890					895					900
Gly	Glu	Lys	Asp	Pro	Asp	Phe	Asn	Gln	Ile	Met	His	Ala	Phe	Ser
				905					910					915
Val	Ala	Pro	Phe	Asp	Gln	Asn	Leu	Ser	Ile	Asp	Gly	Lys	Ile	Leu
				920					925					930
Ser	Asp	Asp	Cys	Ala	Thr	Leu	Gly	Thr	Leu	Gly	Val	Ile	Pro	Glu
				935					940					945
Ser	Val	Ile	Leu	Leu	Lys	Ala	Asp	Glu	Pro	Ile	Ala	Asp	Tyr	Ala
				950					955					960
Ala	Met	Asp	Asp	Val	Met	Gln	Val	Cys	Met	Pro	Glu	Glu	Gly	Phe
				965					970					975
Lys	Gly	Thr	Gly	Leu	Leu	Gly	His							
				980										

<210> 18

<211> 227

<212> PRT

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 1973875CD1

<400> 18

Met Gly Asn Cys Val Gly Arg Gln Arg Arg Glu Arg Pro Ala Ala

&lt;400&gt; 53

```

gootgggttg  gcoqngtcoo  tetgetyccc  cngdcoggtg  caggcoctgc  cggagagccc  60
agggtgtcccg  gngtctggcc  caggacctgg  cccacacagg  gtcccagcag  gggtagctcc  120
tcccgggaca  aggacogag  tgogacggto  agtngttcag  tccccatgco  tgctggaggg  180
aaaggaagcc  atcttccatc  tacaccccag  aggggtccca  accgctgat  ccagagagag  240
tcccatacc  tccatcaaca  tgootacaat  cctgtggtct  ggtacccctg  gggagagaaa  300
goottcgaca  aggccagga  ggaagagggg  cggatttcc  tctcagtcgg  gtactccacc  360
tyccactggt  gccacatgat  ggaagagggg  tcttccaga  atgaggagat  tggccgctg  420
ctcagtgagg  actttgtgag  tgtgaaggta  gaccgtgagg  agcggcoctg  cgtggacaaq  480
gtgtacatga  cgttccgtga  ggcoac<ag>  agcggcgggg  gctggg<cat  gaatgtgtgg  540
ctgactccca  acctccagcc  otttgtcggg  ggcaectatt  tccctccga  ggttggcttg  600
acc<gggtcg  gcttccgca  agtggttgct  agaatacag  aacagtggaa  acagaaacag  660
aacacccctg  tagaaaatag  ccagcgtgtc  accactgccc  tgetggcccc  atcagagato  720
agcgtgggtg  accgccagct  gccgcctct  gccgccaccg  tgaacaatcg  ctgcttccag  780
cagctggatg  agggcataga  tgaggaatac  ggtgctctg  ctgaggcccc  caagtctccc  840
acgcgggtga  tcttgagctt  cctgttctcc  tactggctca  gccatcgact  gactcaggat  900
ggctctcggg  cccagcagat  ggcocttgc  accctgaaa  tgatggctaa  cgggggcatc  960
cgggaccatg  tggggcaggg  ctttccacgc  tactccacag  accgccagtg  gcacgtccct  1020
caclttaga  agatgtctta  lgaacaggca  cagtctgtg  tggcctatto  gnaggtcttc  1080
cagctcactg  gtgatgaatt  ctactctgac  gtggccaaag  gcactcctga  gtacgtgggt  1140
cggagcctga  gccacoggtc  cggaggcttc  tatacgccag  aagatgcaga  ctgcgcccca  1200
gagcggggcc  agcggcccaa  agagggcgcc  tactatgtgt  ggaaggtcaa  agaggttcag  1260
cagctcctcc  cggagcctgt  gttggttgc  aacagccgc  tgactccagg  ccagctctcc  1320
atgaagcaat  accgctccac  agaggtggt  aacatcagcc  ccagtcagg  ccccaagggt  1380
gagctgcagg  gccagaatgt  gctgaccgtc  cggctactgc  tggagctgac  tgcctgccgc  1440
tttggtcttg  atgtggaggc  cgtgogggac  ttgctcaatt  cagggttga  gaaatcttc  1500
caggcccgga  agcatcgccc  caagccgcac  ctggacagca  agatgctggc  tgcctgggat  1560
ggcttgatgg  tglcaggtta  tgcctgtgac  ggggtctgtc  tgggccaaga  caggtctgac  1620
aactatgca  ccaatggtgc  caagttcccg  aagcggcaca  tgtttgatgt  ggcagtggc  1680
cgcctggtgc  ggacctgcta  caccggcct  ggggggactg  tggagcacag  caacccccc  1740
tgctggggct  tcttgaggga  ctacgccttc  gttgtccggg  gctctgcggc  tgcaggacac  1800
gcctccaggg  agagtgcgtg  gctcgagtg  ttctgcagty  aggtgagct  ggggctggc  1860
ctcttttggg  actccaggg  tggggctaac  ggagcagagc  ccagcgccaa  aggactggat  1920
ctgcccctgc  gtctgaagga  cgaccaggat  acgggccaca  agatcgctgt  ctgtggagac  1980
gcccacacac  tgcctccggt  gcattgcttc  ttccagagcc  tcccggtggc  ggtggagac  2040
gtgtgcctat  tgacccgctt  tccagagcc  accctcaagc  agatcgctgt  ctgtggagac  2100
atgttccggc  ccttccagc  ggccctgggt  cagtggllcc  acttctgtct  caltcttacc  2160
cgtcaggcca  ttctggctga  tggggacccc  tccagcttcc  tgtcccgcca  gctgcttcc  2220
ctgagtlccc  tccagcaggt  ggaagagcag  gccantgcac  atgtgtgtga  gactcaggcc  2280
tgcagcttcc  ccaatcactga  tccctggcaa  ttacgaaaac  tactacatcc  atgactgcc  2340
caaccccttc  ggggtggggc  agaaggtgae  gcatcccaac  tgactagaga  ctcaggccct  2400
gcaggggccct  atagaacctg  tggccatccc  tgaagcaacct  gccaccaggt  gaactcggcc  2460
atctcactg  cccccccttg  gcacccactc  accctagaa  aaacttaaca  gtgtcccggt  2520
gtaaaaaaa  aaaaaaa  2597

```

&lt;210&gt; 54

&lt;211&gt; 2385

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5583922CB1

&lt;400&gt; 54

```

gaccgacccg  ccacccgcgc  tagcccggc  gccctgggca  ctcaatcccc  gccatgtggg  60
ggctcctgct  cgccttgccc  gcccttcgcc  cggccctcgg  cccagctctg  ggggccccca  120
ggaaactcgt  gctgggcttc  gccagcccg  ggaccaccaa  ggtcccaggc  tccgcccgg  180
cctgtcatag  cagcccgcca  cagcccgccg  cggagacagc  taacgggacc  tcagacagc  240
atgtccggat  tccaggtuac  aagaagaaaa  aggtcattat  gaagagaggg  aagagctaa  300
ctctaaactg  cccaccccca  ctggtgactg  cggggccctt  tgtgaccccc  actccagcag  360
ggacccctga  ccccgctgag  aaacaaagaa  caggtcttcc  tcccttgggt  ctggagctcc  420
tgagagtttc  agatagccgg  ettgagggat  ccagcagcca  gtcctttggg  ctggagccac  480
accagggagc  gctaaacatt  cagtccaggg  tggaggacgg  cgalctatat  galggagcct  540
ggtgtgctga  ggagcaggac  gccgatccat  ggtttcagggt  ggacgctggg  cacccccacc  600
gcttctcgga  tgttatccaa  cagggcagta  actctgtctg  gagglatgac  tgggtccact  660

```

catacaaggt	ccagttcagc	aatgacagtc	ggacctggty	gggaagtagg	amccacagca	720
gtgggatgga	cgcagtattt	cotgccaatt	cagccccaga	aactccagtg	ctgaacctcc	780
tgccggagcc	ccaggtggcc	cgttccatcc	gcctgctgcc	ccagacctgg	ctccagggag	840
gcgcgccttg	cctccgggca	gagatccctg	cctgcccagt	ctcagacccc	aalgacctat	900
tccttgaggg	ccttgctgct	ggatccctct	acctctctaga	ctttcagcat	cacaattaca	960
gggcctatgag	gaagctgagt	aagcaggtac	aagagcaalg	ccccacatc	acccgcctct	1020
acagcattgg	gaagagctac	cagggcctga	agctgtatgt	gatggaaatg	tcggacaagc	1080
ctggggagca	tgagctgugc	gagcctgagc	tgctctacgt	ggctggcagt	catggggacg	1140
agggcctggg	gogggagttg	cttctgctcc	tgatgcagtt	cctgtgccat	gagttcctgc	1200
gagggaaacc	acgggtgacc	cggctgctct	ctgagatggc	cattccacct	ctggcctcca	1260
tgaaacctga	tggtatgag	atgcctaac	acgggggttc	agagctggty	ggctgggccc	1320
agggccgctg	gaacaacctg	agcctcgatc	ttaccctaaa	ttttgctgac	ctcaacacac	1380
actgtgggga	agcacaggac	gatgggaaag	tgccccacat	cgtccccaac	catcacctgc	1440
cattgcccac	ttactacacc	ctgcccacat	ccacctgggc	tcctgaaacg	cgggcagtaa	1500
traaagtggat	gaaggagatc	ccclltgtgc	taagtgcaca	cctccacggg	ggtagctctg	1560
tggtgtccta	cccattcgac	altgactcgca	ccccgtgggc	tgcccgcyag	ctnaccacca	1620
caccagatga	tgatgltgtt	cgctggctca	gcactgtcta	tgctggcagt	aactctggca	1680
tgccaggacac	cagctgcaga	ccctgccaca	gccaggactt	ctccgtgcac	ggcaacatca	1740
tcacaggygc	tgacttgcac	acggtccccc	ggagcatgaa	tgacttcagc	taacctacaa	1800
ccactgctt	tgaggtcact	gtggagctgt	cctgtgacaa	gttccctcac	gagaatgaat	1860
tgcccacagga	gtgggagaaac	aacaaagacy	ccctctctac	ctacctggag	caggtgcgca	1920
tgggcattgc	aggaagtgtg	agggacaagg	acacggagct	tgsgattgct	gacgctgtca	1980
ctgcctggga	tgggattaac	catgacgtga	ccacggcctg	ggcgggggat	tattggcgtc	2040
tgctgacccc	aggggactac	atggtgaact	ccagtgcoga	gggtaccat	tcagtgcacac	2100
ggaactgtcg	ggtcaccttt	gaagaggggc	ccttccctcg	caetttcgtg	ctcaccacga	2160
ctcccaaac	gaggtctgcgc	gagctgctgg	cagctggggc	caaggtgccc	coggaccttc	2220
gcaggcgcc	ggagcggcta	aggggacaga	aggattgata	cctgcggttt	aagagcccta	2280
gggcaggctg	gacctgtcaa	gacgggaagc	ggagagctag	aggggtgagg	acaaagtgaq	2340
gaaaxggtgc	tcattcaagc	taccgggcac	cttaaaaaaa	aaaaa		2385